



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:22 AM GMT

PDB ID : 2A0P
Title : Crystal structure of RNA oligomer containing 4'-thioribose
Authors : Haeberli, P.; Berger, I.; Pallan, P.S.; Egli, M.
Deposited on : 2005-06-16
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1953 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)
RNA backbone	2183	1002 (2.70-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	8	
1	B	8	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 466 atoms, of which 59 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*CP*CP*(S4C)P*CP*GP*GP*GP*G)-3'.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	8	Total	C	H	N	O	P	S	0	0	0
			198	76	29	32	53	7	1			
1	B	8	Total	C	H	N	O	P	S	0	0	0
			199	76	30	32	53	7	1			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	35	Total	O	0	0
			35	35		
2	B	34	Total	O	0	0
			34	34		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-R(*CP*CP*(S4C)P*CP*GP*GP*GP*G)-3'

Chain A: 



- Molecule 1: 5'-R(*CP*CP*(S4C)P*CP*GP*GP*GP*G)-3'

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	42.44Å 42.44Å 127.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 1.95 21.28 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (25.00-1.95) 88.5 (21.28-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.96Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.202 , (Not available) 0.215 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.360	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.13 , 59.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	5 of 3080 reflections (0.162%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	466	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: S4C

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.46	24/165 (14.5%)	3.37	34/254 (13.4%)
1	B	3.45	21/165 (12.7%)	3.52	34/254 (13.4%)
All	All	3.45	45/330 (13.6%)	3.44	68/508 (13.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	10	C	N3-C4	-11.79	1.25	1.33
1	A	8	G	N1-C2	-10.78	1.29	1.37
1	A	6	G	N7-C5	10.65	1.45	1.39
1	B	10	C	N1-C2	-10.20	1.29	1.40
1	A	8	G	C5-C4	-8.82	1.32	1.38
1	A	8	G	C2'-C1'	-8.79	1.43	1.53
1	B	15	G	C6-N1	-7.86	1.34	1.39
1	B	15	G	N9-C8	-7.86	1.32	1.37
1	B	10	C	C3'-C2'	-7.83	1.44	1.52
1	A	7	G	P-O5'	7.81	1.67	1.59
1	A	7	G	C2'-C1'	-7.77	1.44	1.53
1	A	7	G	N3-C4	7.68	1.40	1.35
1	A	5	G	C5-C6	-7.58	1.34	1.42
1	B	13	G	N7-C5	7.46	1.43	1.39
1	A	2	C	O4'-C1'	7.30	1.51	1.41
1	B	13	G	C5-C6	-7.27	1.35	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	16	G	P-O5'	-7.16	1.52	1.59
1	A	2	C	C2'-O2'	6.96	1.50	1.41
1	A	5	G	C3'-C2'	-6.91	1.45	1.52
1	B	12	C	N1-C6	-6.64	1.33	1.37
1	A	1	C	C4'-C3'	-6.63	1.45	1.53
1	B	10	C	N1-C6	-6.53	1.33	1.37
1	B	16	G	O4'-C1'	6.52	1.50	1.41
1	A	5	G	C5-C4	-6.48	1.33	1.38
1	B	16	G	C3'-C2'	-6.37	1.45	1.52
1	A	2	C	N3-C4	-6.25	1.29	1.33
1	A	8	G	C6-N1	-6.21	1.35	1.39
1	A	1	C	C3'-C2'	-6.21	1.46	1.52
1	B	12	C	P-O5'	6.09	1.65	1.59
1	A	8	G	N7-C5	6.05	1.42	1.39
1	A	8	G	P-O5'	-6.00	1.53	1.59
1	B	12	C	O3'-P	5.80	1.68	1.61
1	B	10	C	C4-C5	-5.78	1.38	1.43
1	A	4	C	C4'-C3'	-5.54	1.47	1.52
1	B	15	G	C2'-C1'	-5.46	1.47	1.53
1	A	5	G	O3'-P	5.38	1.67	1.61
1	A	4	C	C3'-C2'	-5.34	1.46	1.52
1	A	6	G	N9-C8	5.34	1.41	1.37
1	A	1	C	C2'-C1'	-5.28	1.47	1.53
1	B	15	G	C8-N7	-5.24	1.27	1.30
1	A	6	G	O4'-C1'	5.24	1.48	1.41
1	B	14	G	C8-N7	-5.20	1.27	1.30
1	B	13	G	C5-C4	-5.12	1.34	1.38
1	B	13	G	N9-C8	5.09	1.41	1.37
1	B	14	G	C5-C4	-5.04	1.34	1.38

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	12	C	O4'-C1'-N1	15.01	120.21	108.20
1	B	13	G	C5-C6-O6	-13.02	120.79	128.60
1	B	9	C	O4'-C4'-C3'	-12.30	91.69	104.00
1	B	10	C	O4'-C4'-C3'	-11.49	92.51	104.00
1	A	7	G	O4'-C4'-C3'	-10.00	94.00	104.00
1	B	13	G	N1-C6-O6	9.86	125.81	119.90
1	B	13	G	C4-C5-N7	9.61	114.64	110.80
1	A	7	G	N9-C1'-C2'	-9.49	101.56	112.00
1	A	6	G	C5-N7-C8	-9.27	99.67	104.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	9	C	P-O3'-C3'	9.18	130.71	119.70
1	A	5	G	N1-C6-O6	9.09	125.36	119.90
1	A	5	G	C5-C6-O6	-8.55	123.47	128.60
1	A	8	G	O4'-C4'-C3'	-8.50	95.50	104.00
1	B	15	G	O4'-C4'-C3'	-8.30	95.70	104.00
1	A	1	C	C1'-O4'-C4'	8.30	116.54	109.90
1	B	12	C	C5'-C4'-C3'	-8.07	103.08	116.00
1	A	1	C	O4'-C4'-C3'	-8.05	95.95	104.00
1	A	7	G	C5-C6-O6	-7.85	123.89	128.60
1	A	2	C	O4'-C4'-C3'	-7.69	96.31	104.00
1	B	14	G	N1-C6-O6	-7.47	115.42	119.90
1	B	16	G	N1-C6-O6	-7.45	115.43	119.90
1	A	2	C	N1-C2-O2	7.42	123.35	118.90
1	B	9	C	O4'-C1'-C2'	-7.40	98.40	105.80
1	A	1	C	N3-C2-O2	-7.39	116.73	121.90
1	A	5	G	O4'-C4'-C3'	-7.37	96.63	104.00
1	B	15	G	N3-C2-N2	-7.34	114.76	119.90
1	B	10	C	O4'-C1'-C2'	-7.33	98.47	105.80
1	A	7	G	O4'-C1'-N9	7.29	114.03	108.20
1	A	7	G	C1'-O4'-C4'	7.26	115.71	109.90
1	A	6	G	P-O3'-C3'	7.25	128.41	119.70
1	A	8	G	C1'-O4'-C4'	7.20	115.66	109.90
1	A	6	G	O4'-C4'-C3'	-7.14	96.86	104.00
1	B	15	G	C4-C5-N7	-7.10	107.96	110.80
1	B	14	G	O4'-C1'-N9	-7.05	102.56	108.20
1	A	1	C	O4'-C1'-C2'	-6.97	98.83	105.80
1	A	8	G	C5'-C4'-O4'	6.83	117.30	109.10
1	A	6	G	O4'-C1'-C2'	-6.79	99.02	105.80
1	B	12	C	N1-C1'-C2'	-6.59	104.75	112.00
1	B	12	C	C5'-C4'-O4'	6.58	117.00	109.10
1	A	5	G	O4'-C1'-N9	-6.57	102.94	108.20
1	B	13	G	C5-N7-C8	-6.54	101.03	104.30
1	B	16	G	C5-N7-C8	6.53	107.56	104.30
1	B	14	G	O4'-C4'-C3'	-6.47	97.53	104.00
1	B	12	C	N1-C2-O2	6.38	122.72	118.90
1	A	6	G	C1'-O4'-C4'	6.32	114.96	109.90
1	B	9	C	C1'-O4'-C4'	6.20	114.86	109.90
1	A	7	G	N3-C2-N2	6.13	124.19	119.90
1	A	7	G	C5-N7-C8	-6.12	101.24	104.30
1	B	12	C	N3-C2-O2	-6.10	117.63	121.90
1	A	1	C	N1-C2-O2	6.07	122.54	118.90
1	A	8	G	N1-C6-O6	-5.96	116.33	119.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	13	G	O4'-C1'-C2'	-5.90	99.90	105.80
1	B	14	G	OP1-P-OP2	5.87	128.41	119.60
1	A	7	G	O5'-P-OP2	5.80	117.66	110.70
1	B	9	C	C5-C4-N4	-5.78	116.15	120.20
1	A	4	C	OP1-P-O3'	-5.64	92.79	105.20
1	A	7	G	C4-C5-N7	5.52	113.01	110.80
1	B	14	G	C4-C5-N7	-5.51	108.59	110.80
1	A	8	G	C5'-C4'-C3'	-5.39	107.37	116.00
1	B	14	G	C5-C6-N1	5.34	114.17	111.50
1	B	12	C	O3'-P-O5'	5.32	114.10	104.00
1	B	13	G	C6-C5-N7	-5.29	127.23	130.40
1	A	6	G	C4-C5-N7	5.16	112.86	110.80
1	B	9	C	N3-C4-N4	5.13	121.59	118.00
1	A	5	G	N1-C2-N2	5.09	120.78	116.20
1	B	16	G	C6-N1-C2	-5.06	122.06	125.10
1	B	14	G	C6-N1-C2	-5.05	122.07	125.10
1	A	7	G	P-O3'-C3'	5.02	125.72	119.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	8	G	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	169	29	89	0	0
1	B	169	30	90	0	0
2	A	35	0	0	0	0
2	B	34	0	0	0	0
All	All	407	59	179	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	6/8 (75%)	0	0
1	B	6/8 (75%)	0	0
All	All	12/16 (75%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	S4C	A	3	1	13,21,22	1.75	3 (23%)	10,30,33	2.41	4 (40%)
1	S4C	B	11	1	13,21,22	2.82	1 (7%)	10,30,33	4.45	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	S4C	A	3	1	-	0/2/25/26	0/2/2/2
1	S4C	B	11	1	-	0/2/25/26	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3	S4C	C3'-C2'	-2.31	1.47	1.53
1	A	3	S4C	O2'-C2'	-2.09	1.38	1.43
1	A	3	S4C	C6-N1	3.88	1.41	1.35
1	B	11	S4C	C3'-C4'	9.30	1.59	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	11	S4C	C3'-C4'-S4'	-12.99	90.10	105.73
1	A	3	S4C	C3'-C4'-S4'	-4.31	100.55	105.73
1	B	11	S4C	C5'-C4'-S4'	-2.33	107.89	111.45
1	A	3	S4C	O3'-C3'-C2'	-2.15	104.85	111.83
1	A	3	S4C	C5'-C4'-S4'	3.25	116.42	111.45
1	B	11	S4C	C2-N3-C4	4.33	121.72	115.61
1	A	3	S4C	C2-N3-C4	4.83	122.43	115.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	7/8 (87%)	-0.38	0 100 100	18, 19, 22, 24	0
1	B	7/8 (87%)	-0.22	0 100 100	12, 17, 22, 25	0
All	All	14/16 (87%)	-0.30	0 100 100	12, 19, 24, 25	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	S4C	A	3	20/21	0.95	0.09	-	15,18,28,29	0
1	S4C	B	11	20/21	0.94	0.10	-	11,15,25,28	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.