



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:22 AM GMT

PDB ID : 2A0U  
Title : Crystal structure of the eukaryotic initiation factor 2B from Leishmania major at 2.1 Å resolution  
Authors : Bosch, J.; Hol, W.G.J.; Structural Genomics of Pathogenic Protozoa Consortium (SGPP)  
Deposited on : 2005-06-17  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	383	<div> <div>12%</div> <div>90%</div> <div>7% ..</div> </div>
1	B	383	<div> <div>9%</div> <div>88%</div> <div>7% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	401	-	-	-	X
2	SO4	A	404	-	-	-	X
2	SO4	B	409	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5968 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called initiation factor 2B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	Se	0	1	0
			2831	1782	494	543	8	4			
1	B	367	Total	C	N	O	S	Se	0	1	0
			2779	1750	482	535	8	4			

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	159	Total	O	0	0
			159	159		
3	B	149	Total	O	0	0
			149	149		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: initiation factor 2B



#### • Molecule 1: initiation factor 2B



## 4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.73Å 164.57Å 172.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.30 – 2.10 15.25 – 2.10	Depositor EDS
% Data completeness (in resolution range)	85.4 (15.30-2.10) 85.7 (15.25-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.85 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.214 , 0.264 0.218 , 0.270	Depositor DCC
$R_{free}$ test set	1851 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.8	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 43.4	EDS
Estimated twinning fraction	0.007 for -h,l,k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 36707 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5968	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/2876	0.51	0/3907
1	B	0.32	0/2821	0.49	0/3832
All	All	0.32	0/5697	0.50	0/7739

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	16	ALA	Peptide
1	A	371	ALA	Peptide
1	B	17	THR	Peptide
1	B	327	VAL	Peptide



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2831	0	2870	22	0
1	B	2779	0	2824	22	0
2	A	25	0	0	1	0
2	B	25	0	0	0	0
3	A	159	0	0	0	0
3	B	149	0	0	0	0
All	All	5968	0	5694	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:LEU:HA	1:A:88:LYS:CB	1.78	1.11
1:B:371:ALA:HB1	1:B:372:PRO:HD2	1.31	1.06
1:A:87:LEU:CA	1:A:88:LYS:HB2	1.90	1.01
1:A:87:LEU:HA	1:A:88:LYS:HB2	0.97	0.97
1:B:252:MSE:HE1	1:B:260:VAL:CG2	2.06	0.85
1:B:371:ALA:HB1	1:B:372:PRO:CD	2.12	0.79
1:B:182:LEU:HD23	1:B:252:MSE:HE3	1.64	0.79
1:A:371:ALA:HB1	1:A:372:PRO:HD2	1.67	0.77
1:B:18:LEU:O	1:B:18:LEU:HG	1.95	0.67
1:B:252:MSE:HE1	1:B:260:VAL:HG21	1.75	0.66
1:B:371:ALA:CB	1:B:372:PRO:HD2	2.19	0.65
1:A:371:ALA:HB1	1:A:372:PRO:CD	2.26	0.65
1:A:61:ARG:HH12	1:A:191:ALA:HB2	1.64	0.62
1:A:87:LEU:CA	1:A:88:LYS:CB	2.61	0.61
1:B:252:MSE:CE	1:B:260:VAL:CG2	2.78	0.59
1:B:31:LEU:HD21	1:B:36:LEU:HD11	1.88	0.55
1:A:82:ALA:HA	1:A:87:LEU:O	2.08	0.54
1:B:18:LEU:HD13	1:B:231:TYR:CE2	2.44	0.53
1:A:133:ALA:O	1:A:135:ALA:N	2.41	0.52
1:B:63:ALA:HB3	1:B:64:PRO:HD3	1.92	0.52
1:B:252:MSE:CE	1:B:260:VAL:HG22	2.39	0.52

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:213:ARG:HD3	1:B:241:THR:OG1	2.09	0.52
1:A:213:ARG:HD3	1:A:241:THR:OG1	2.11	0.51
1:A:33:GLN:O	1:A:227:ARG:NH2	2.45	0.49
1:A:61:ARG:HD2	2:A:402:SO4:O1	2.13	0.48
1:A:63:ALA:HB3	1:A:64:PRO:HD3	1.94	0.48
1:B:299:THR:HA	1:B:302:LEU:HD13	1.96	0.47
1:A:145:ALA:O	1:A:148:VAL:HG22	2.16	0.46
1:B:162:HIS:CG	1:B:363:GLY:HA2	2.51	0.46
1:A:61:ARG:NH1	1:A:191:ALA:HB2	2.29	0.45
1:A:371:ALA:CB	1:A:372:PRO:HD2	2.42	0.45
1:A:370:SER:O	1:A:371:ALA:O	2.35	0.45
1:A:156:ASN:O	1:A:160:MSE:HG3	2.17	0.44
1:B:105:MSE:HE3	1:B:115:PHE:HD2	1.81	0.44
1:A:342:TRP:HB3	1:B:243:ILE:HG22	1.98	0.44
1:A:162:HIS:CG	1:A:363:GLY:HA2	2.53	0.44
1:A:350:PRO:HD2	1:A:353:LEU:HD12	2.00	0.43
1:A:14:HIS:O	1:A:15:HIS:CD2	2.72	0.43
1:B:145:ALA:O	1:B:148:VAL:HG22	2.18	0.42
1:B:371:ALA:CB	1:B:372:PRO:CD	2.86	0.42
1:B:327:VAL:O	1:B:328:THR:CB	2.68	0.42
1:B:350:PRO:HD2	1:B:353:LEU:HD12	2.03	0.41
1:B:105:MSE:HG3	1:B:118:LEU:HD12	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	373/383 (97%)	355 (95%)	14 (4%)	4 (1%)	17	11
1	B	366/383 (96%)	351 (96%)	11 (3%)	4 (1%)	17	11
All	All	739/766 (96%)	706 (96%)	25 (3%)	8 (1%)	17	11

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	88	LYS
1	A	371	ALA
1	A	372	PRO
1	B	18	LEU
1	B	327	VAL
1	B	328	THR
1	B	371	ALA
1	A	134	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/302 (100%)	298 (99%)	3 (1%)	82	87
1	B	296/302 (98%)	291 (98%)	5 (2%)	68	74
All	All	597/604 (99%)	589 (99%)	8 (1%)	76	82

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	61	ARG
1	A	72	LEU
1	B	18	LEU
1	B	61	ARG
1	B	325	ASN
1	B	326	LEU
1	B	327	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	289	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	289	HIS
1	B	325	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	401	-	4,4,4	0.21	0	6,6,6	0.07	0
2	SO4	A	402	-	4,4,4	0.24	0	6,6,6	0.15	0
2	SO4	A	403	-	4,4,4	0.19	0	6,6,6	0.06	0
2	SO4	A	404	-	4,4,4	0.21	0	6,6,6	0.06	0
2	SO4	A	405	-	4,4,4	0.21	0	6,6,6	0.10	0
2	SO4	B	406	-	4,4,4	0.21	0	6,6,6	0.12	0
2	SO4	B	407	-	4,4,4	0.21	0	6,6,6	0.09	0
2	SO4	B	408	-	4,4,4	0.24	0	6,6,6	0.19	0
2	SO4	B	409	-	4,4,4	0.22	0	6,6,6	0.08	0
2	SO4	B	410	-	4,4,4	0.23	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	401	-	-	0/0/0/0	0/0/0/0
2	SO4	A	402	-	-	0/0/0/0	0/0/0/0
2	SO4	A	403	-	-	0/0/0/0	0/0/0/0
2	SO4	A	404	-	-	0/0/0/0	0/0/0/0
2	SO4	A	405	-	-	0/0/0/0	0/0/0/0
2	SO4	B	406	-	-	0/0/0/0	0/0/0/0
2	SO4	B	407	-	-	0/0/0/0	0/0/0/0
2	SO4	B	408	-	-	0/0/0/0	0/0/0/0
2	SO4	B	409	-	-	0/0/0/0	0/0/0/0
2	SO4	B	410	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	402	SO4	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	369/383 (96%)	0.77	45 (12%) 5 7	23, 36, 54, 63	0
1	B	363/383 (94%)	0.66	33 (9%) 11 16	24, 36, 54, 64	0
All	All	732/766 (95%)	0.71	78 (10%) 8 11	23, 36, 54, 64	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	326	LEU	7.9
1	B	17	THR	7.1
1	B	326	LEU	6.0
1	A	16	ALA	5.7
1	B	327	VAL	5.2
1	A	127	LYS	4.9
1	B	133	ALA	4.6
1	A	17	THR	4.4
1	A	131	THR	4.2
1	A	133	ALA	4.2
1	B	195	TYR	4.1
1	B	131	THR	4.0
1	A	327	VAL	4.0
1	A	383	ALA	3.9
1	A	15	HIS	3.9
1	B	371	ALA	3.9
1	A	87	LEU	3.8
1	B	18	LEU	3.8
1	A	91	ARG	3.7
1	A	137	VAL	3.7
1	B	174	GLU	3.6
1	A	92	GLU	3.4
1	B	91	ARG	3.4
1	B	370	SER	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	171	ALA	3.4
1	A	161	ARG	3.2
1	A	371	ALA	3.1
1	A	134	ALA	3.1
1	A	123	ALA	3.0
1	B	328	THR	3.0
1	B	128	LEU	3.0
1	A	12	LYS	2.9
1	A	159	ILE	2.8
1	B	177	ASP	2.8
1	A	369	ALA	2.8
1	B	209	GLY	2.8
1	A	177	ASP	2.8
1	B	129	ASP	2.8
1	B	310	ASN	2.8
1	A	310	ASN	2.7
1	B	381	ALA	2.7
1	A	89	SER	2.7
1	A	372	PRO	2.7
1	A	90	GLY	2.7
1	A	13	PRO	2.7
1	A	128	LEU	2.5
1	A	311	HIS	2.5
1	B	84	ASN	2.5
1	A	205	LEU	2.5
1	A	140	ALA	2.5
1	A	132	LYS	2.4
1	B	125	VAL	2.4
1	B	173	ALA	2.4
1	B	175	GLY	2.4
1	A	14	HIS	2.4
1	A	175	GLY	2.4
1	A	337	PRO	2.3
1	A	49	GLU	2.3
1	A	130	PRO	2.2
1	B	238	ILE	2.2
1	B	325	ASN	2.2
1	B	292	LYS	2.2
1	A	121	LEU	2.2
1	A	195	TYR	2.2
1	B	212	GLU	2.1
1	B	135	ALA	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	52	TRP	2.1
1	A	18	LEU	2.1
1	A	82	ALA	2.1
1	B	112	VAL	2.1
1	B	30	LEU	2.1
1	B	99	THR	2.0
1	A	156	ASN	2.0
1	B	138	ALA	2.0
1	B	136	GLU	2.0
1	A	45	VAL	2.0
1	A	174	GLU	2.0
1	A	370	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	409	5/5	0.73	0.35	35.30	78,79,79,79	0
2	SO4	A	401	5/5	0.89	0.27	4.53	43,44,45,45	0
2	SO4	A	404	5/5	0.86	0.27	2.79	57,58,59,59	0
2	SO4	B	406	5/5	0.64	0.28	1.14	70,70,71,72	0
2	SO4	A	403	5/5	0.85	0.23	0.85	57,59,60,60	0
2	SO4	B	407	5/5	0.84	0.20	0.40	37,37,39,39	0
2	SO4	A	402	5/5	0.98	0.09	-1.62	25,25,29,29	0
2	SO4	B	408	5/5	0.99	0.07	-1.62	25,26,28,28	0
2	SO4	B	410	5/5	0.89	0.41	-	95,95,95,95	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	405	5/5	0.88	0.29	-	55,56,56,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.