



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 27, 2016 – 05:38 AM BST

PDB ID : 5A3G
Title : Structure of herpesvirus nuclear egress complex subunit M50
Authors : Leigh, K.E.; Boeszoermenyi, A.; Mansueto, M.S.; Sharma, M.; Filman, D.J.;
Coen, D.M.; Wagner, G.; Hogle, J.M.; Arthanari, H.
Deposited on : 2015-06-01

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 7%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	171	<div> <div style="width: 79%; background-color: green;"></div> <div style="width: 16%; background-color: yellow;"></div> <div style="width: 5%; background-color: cyan;"></div> <div style="width: 0%; background-color: grey;"></div> <div style="width: 0%; background-color: red;"></div> </div> <div>79% 16% . .</div>

2 Ensemble composition and analysis

This entry contains 15 models. Model 12 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:7-A:171 (165)	0.93	12

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 5, 6, 7, 8, 11, 12, 13
2	1, 10, 14, 15
Single-model clusters	4; 9

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2711 atoms, of which 1356 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called M50.

Mol	Chain	Residues	Atoms							Trace
1	A	171	Total	C	H	N	O	S		0
			2711	851	1356	234	258	12		

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	EXPRESSION TAG	UNP H2A365
A	2	GLY	-	EXPRESSION TAG	UNP H2A365
A	3	HIS	-	EXPRESSION TAG	UNP H2A365

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: M50



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 12. Colouring as in section 4.1 above.

- Molecule 1: M50



5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING WITH TORSION ANGLE DYNAMICS SIMULATION FOLLOWED BY RDC AND WATERBOX REFINEMENT*.

Of the 50 calculated structures, 15 were deposited, based on the following criterion: *LOWEST ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	
CYANA	structure solution	3.0
NMRDRAW	structure solution	11
TALOSPLUS	structure solution	1.2009.0605.17
CCPNMR ANALYSIS	structure solution	2.4
NMRPIPE	structure solution	11
HMSIST	structure solution	2.11
CARA	structure solution	1.8.4

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	5a3g_cs.str
Number of chemical shift lists	1
Total number of shifts	300
Number of shifts mapped to atoms	300
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	7%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.74±0.01	0±0/1331 (0.0±0.0%)	1.31±0.03	14±2/1801 (0.8±0.1%)
All	All	0.74	0/19965 (0.0%)	1.31	208/27015 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	9.3±2.0
All	All	0	140

There are no bond-length outliers.

5 of 33 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	101	ARG	NE-CZ-NH1	12.11	126.36	120.30	7	11
1	A	123	ARG	NE-CZ-NH1	10.40	125.50	120.30	5	14
1	A	23	ARG	NE-CZ-NH1	9.38	124.99	120.30	11	14
1	A	19	ARG	NE-CZ-NH1	9.33	124.97	120.30	6	13
1	A	138	ARG	NE-CZ-NH1	9.06	124.83	120.30	7	13

There are no chirality outliers.

5 of 46 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	102	VAL	Peptide	12
1	A	89	TYR	Peptide,Sidechain	12
1	A	101	ARG	Peptide	9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	82	CYS	Peptide	7
1	A	146	TYR	Sidechain,Peptide	7

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1311	1313	1313	2±2
All	All	19665	19695	19695	31

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

5 of 19 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:76:VAL:HG22	1:A:115:VAL:HG22	0.86	1.48	10	3
1:A:113:ALA:C	1:A:114:LEU:HD23	0.62	2.15	13	6
1:A:29:LEU:HD11	1:A:48:ALA:HB1	0.58	1.76	15	2
1:A:139:LYS:HE2	1:A:146:TYR:CD1	0.55	2.36	15	3
1:A:101:ARG:CD	1:A:102:VAL:H	0.53	2.17	5	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/171 (96%)	145±2 (88±2%)	15±2 (9±1%)	4±2 (3±1%)	11	48
All	All	2460/2565 (96%)	2176 (88%)	222 (9%)	62 (3%)	11	48

5 of 21 unique Ramachandran outliers are listed below. They are sorted by the frequency of

occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	81	GLY	15
1	A	130	GLY	8
1	A	38	CYS	7
1	A	125	LYS	4
1	A	126	PRO	4

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	148/152 (97%)	131±3 (88±2%)	17±3 (12±2%)	11	54
All	All	2220/2280 (97%)	1963 (88%)	257 (12%)	11	54

5 of 83 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	139	LYS	13
1	A	114	LEU	12
1	A	36	LEU	12
1	A	101	ARG	11
1	A	116	VAL	9

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 7% for the well-defined parts and 7% for the entire structure.

7.1 Chemical shift list 1

File name: 5a3g_cs.str

Chemical shift list name: *assigned_chem_shift_list*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	300
Number of shifts mapped to atoms	300
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 7%, i.e. 148 atoms were assigned a chemical shift out of a possible 2050. 0 out of 32 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	0/815 (0%)	0/325 (0%)	0/330 (0%)	0/160 (0%)
Sidechain	148/1105 (13%)	74/643 (12%)	74/402 (18%)	0/60 (0%)
Aromatic	0/130 (0%)	0/68 (0%)	0/61 (0%)	0/1 (0%)
Overall	148/2050 (7%)	74/1036 (7%)	74/793 (9%)	0/221 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list). RCI is only applicable to proteins.