Full wwPDB X-ray Structure Validation Report (i)

Jan 31, 2016 – 06:17 PM GMT

PDB ID	:	1A4A
Title	:	AZURIN MUTANT WITH MET 121 REPLACED BY HIS, PH 6.5 CRYSTAL FORM, DATA COLLECTED AT 16 DEGREES CELSIUS
Authors	:	Messerschmidt, A.; Prade, L.
Deposited on	:	1998-01-28
Resolution	:	1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	129	81% 16% • •
1	B	129	82% 16% •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3057 atoms, of which 880 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AZURIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	129	1199	611	221	167	191	9	0	0	0
1	B	129	1199	611	221	167	191	9	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	ASP	ASN	CONFLICT	UNP P00280
A	42	SER	VAL	SEE REMARK 999	UNP P00280
A	57	GLU	GLN	CONFLICT	UNP P00280
A	121	HIS	MET	ENGINEERED	UNP P00280
B	16	ASP	ASN	CONFLICT	UNP P00280
B	42	SER	VAL	SEE REMARK 999	UNP P00280
B	57	GLU	GLN	CONFLICT	UNP P00280
B	121	HIS	MET	ENGINEERED	UNP P00280

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cu		
2	B	1	1	1	0	0
2	A	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
3	A	110	330	220	110	0	0
3	B	109	327	218	109	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: AZURIN

Chain A:



- Molecule 1: AZURIN

Chain B:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α, β, γ	75.82Å 74.42Å 98.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.89	Depositor
% Data completeness (in resolution range)	95.0 (8.00-1.89)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	0.09	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.189 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3057	wwPDB-VP
Average B, all atoms ($Å^2$)	18.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.76	0/997	1.36	10/1341 (0.7%)
1	B	0.74	0/997	1.34	8/1341 (0.6%)
All	All	0.75	0/1994	1.35	18/2682 (0.7%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	2	GLN	CA-CB-CG	9.20	133.64	113.40
1	A	118	TRP	CD1-CG-CD2	8.20	112.86	106.30
1	A	39	MET	CA-CB-CG	-7.79	100.05	113.30
1	B	118	TRP	CD1-CG-CD2	7.76	112.51	106.30
1	A	48	TRP	CD1-CG-CD2	7.68	112.44	106.30
1	B	48	TRP	CD1-CG-CD2	7.64	112.41	106.30
1	B	39	MET	CG-SD-CE	-7.52	88.17	100.20
1	A	118	TRP	CE2-CD2-CG	-7.28	101.48	107.30
1	A	48	TRP	CE2-CD2-CG	-7.26	101.49	107.30
1	B	48	TRP	CE2-CD2-CG	-7.25	101.50	107.30
1	B	118	TRP	CE2-CD2-CG	-6.63	101.99	107.30
1	A	127	LEU	CA-CB-CG	5.47	127.88	115.30
1	B	129	ASN	N-CA-C	5.28	125.26	111.00
1	B	108	TYR	CB-CG-CD2	-5.24	117.86	121.00
1	B	41	LYS	CA-CB-CG	5.22	124.89	113.40
1	A	110	TYR	CB-CG-CD2	-5.10	117.94	121.00
1	A	118	TRP	CG-CD1-NE1	-5.09	105.01	110.10
1	A	129	ASN	CA-CB-CG	5.08	124.57	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	978	221	950	11	0
1	B	978	221	950	9	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	110	220	0	5	0
3	B	109	218	0	4	0
All	All	2177	880	1900	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:MET:HB2	3:B:202:HOH:O	2.03	0.59
1:B:64:MET:HG3	1:B:115:PRO:HG3	1.88	0.54
1:A:40:ALA:HB1	3:A:140:HOH:O	2.07	0.54
1:B:24:LYS:HB2	3:B:232:HOH:O	2.09	0.53
1:A:4:GLU:HG2	1:A:30:THR:HB	1.91	0.53
1:A:10:ASN:HB2	3:A:142:HOH:O	2.11	0.50
1:A:64:MET:HG3	1:A:115:PRO:HG3	1.93	0.49
1:B:65:ASN:HB3	3:B:179:HOH:O	2.15	0.47
1:B:22:VAL:HB	1:B:127:LEU:HD12	1.96	0.46
1:A:1:ALA:N	3:A:204:HOH:O	2.48	0.46
1:A:1:ALA:O	1:A:2:GLN:HG3	2.15	0.45
1:A:112:CYS:HB3	1:A:121:HIS:HB3	1.98	0.45
1:B:18:LYS:HE2	3:B:136:HOH:O	2.16	0.45
1:B:49:VAL:O	1:B:110:TYR:HA	2.17	0.44
1:B:19:GLU:HA	1:B:124:THR:O	2.20	0.42
1:A:64:MET:HB2	3:A:161:HOH:O	2.18	0.41
1:B:103:THR:HA	1:B:104:PRO:HD3	1.93	0.41
1:A:49:VAL:O	1:A:110:TYR:HA	2.20	0.41
1:A:9:SER:HB2	3:A:174:HOH:O	2.21	0.40
1:A:52:LYS:HE3	1:A:52:LYS:HB2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	127/129 (98%)	121 (95%)	5 (4%)	1 (1%)	24 11
1	B	127/129 (98%)	122 (96%)	5 (4%)	0	100 100
All	All	254/258 (98%)	243 (96%)	10 (4%)	1 (0%)	39 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	103/103 (100%)	96 (93%)	7 (7%)	20 9
1	B	103/103 (100%)	98 (95%)	5 (5%)	31 18
All	All	206/206 (100%)	194 (94%)	12 (6%)	25 13

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	41	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	100	SER
1	A	106	GLU
1	A	115	PRO
1	A	122	LYS
1	A	126	LYS
1	B	14	GLN
1	B	98	ASP
1	B	126	LYS
1	B	127	LEU
1	B	129	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates (i)

EDS was not executed - this section will therefore be empty.

6.4 Ligands (i)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers (i)

EDS was not executed - this section will therefore be empty.