



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 06:49 PM BST

PDB ID : 2A4J  
Title : Solution structure of the C-terminal domain (T94-Y172) of the human centrin 2 in complex with a 17 residues peptide (P1-XPC) from xeroderma pigmentosum group C protein  
Authors : Yang, A.; Miron, S.; Mouawad, L.; Duchambon, P.; Blouquit, Y.; Craescu, C.T.  
Deposited on : 2005-06-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 70%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	79	
2	B	17	

## 2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 14 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:106-A:114, A:120-A:133, A:141-A:149, A:157-A:167, B:2-B:12 (54)	0.18	14

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 2 single-model clusters were found.

Cluster number	Models
1	3, 4, 8, 9, 10, 11, 14, 15, 17
2	6, 16, 19
3	7, 12
4	2, 18
5	5, 13
Single-model clusters	1; 20

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1608 atoms, of which 814 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Centrin 2.

Mol	Chain	Residues	Atoms						Trace
1	A	79	Total	C	H	N	O	S	0
			1283	399	638	104	139	3	

- Molecule 2 is a protein called 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells.


Mol	Chain	Residues	Atoms					Trace
2	B	17	Total	C	H	N	O	0
			325	97	176	31	21	

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Centrin 2



- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.2 Score per residue for model 2

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.3 Score per residue for model 3

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.4 Score per residue for model 4

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.5 Score per residue for model 5

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.6 Score per residue for model 6

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.7 Score per residue for model 7

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.8 Score per residue for model 8

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.9 Score per residue for model 9

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.10 Score per residue for model 10

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.11 Score per residue for model 11

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.12 Score per residue for model 12

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.13 Score per residue for model 13

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.14 Score per residue for model 14 (medoid)

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.15 Score per residue for model 15

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.16 Score per residue for model 16

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.17 Score per residue for model 17

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.18 Score per residue for model 18

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.19 Score per residue for model 19

- Molecule 1: Centrin 2




- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells




#### 4.2.20 Score per residue for model 20

- Molecule 1: Centrin 2



- Molecule 2: 17-mer peptide P1-XPC from DNA-repair protein complementing XP-C cells



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Insight II	refinement	2000

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5992
Number of chemical shift lists	2
Total number of shifts	947
Number of shifts mapped to atoms	947
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	70%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality

### 6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.13±0.01	0±0/366 (0.0±0.0%)	1.56±0.04	7±1/485 (1.4±0.2%)
2	B	1.26±0.02	0±0/94 (0.0±0.0%)	1.69±0.04	1±0/126 (1.2±0.4%)
All	All	1.16	0/9200 (0.0%)	1.59	167/12220 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.9±0.4
All	All	0	58

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	113	PHE	CB-CG-CD2	-11.82	112.53	120.80	13	20
1	A	113	PHE	CB-CG-CD1	8.44	126.71	120.80	13	4
1	A	157	VAL	CA-CB-CG1	8.32	123.37	110.90	5	13
2	B	12	ARG	NE-CZ-NH1	8.19	124.39	120.30	4	20
1	A	128	ARG	NE-CZ-NH1	7.90	124.25	120.30	18	20
1	A	164	ARG	NE-CZ-NH1	7.87	124.23	120.30	9	20
2	B	9	LEU	CB-CG-CD2	7.11	123.08	111.00	6	8
1	A	128	ARG	N-CA-CB	-6.90	98.17	110.60	19	2
1	A	113	PHE	CA-CB-CG	6.90	130.46	113.90	5	20
1	A	129	VAL	CA-CB-CG2	6.59	120.78	110.90	20	5
1	A	142	LEU	CB-CA-C	6.13	121.84	110.20	14	7
1	A	165	ILE	CA-CB-CG1	6.12	122.63	111.00	8	9
1	A	162	PHE	CB-CG-CD1	6.11	125.08	120.80	12	8
1	A	114	ASP	N-CA-CB	-6.11	99.61	110.60	6	2

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	126	LEU	CA-CB-CG	5.95	129.00	115.30	4	4
1	A	126	LEU	CB-CG-CD2	5.95	121.12	111.00	8	1
1	A	113	PHE	N-CA-CB	5.62	120.72	110.60	16	2
1	A	144	GLU	N-CA-CB	-5.20	101.25	110.60	11	1
2	B	12	ARG	N-CA-CB	-5.19	101.26	110.60	5	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	113	PHE	Sidechain	20
1	A	162	PHE	Sidechain	20
1	A	110	PHE	Sidechain	17
1	A	123	PHE	Sidechain	1

## 6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	362	390	390	7±2
2	B	92	112	112	3±2
All	All	9080	10040	10040	165

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:127:LYS:O	1:A:131:LYS:HB2	1.38	1.15	10	20
1:A:160:GLN:O	1:A:164:ARG:HB3	1.21	1.35	6	20
1:A:127:LYS:O	1:A:131:LYS:CB	1.12	1.98	17	20
1:A:160:GLN:O	1:A:164:ARG:CB	1.04	2.04	1	20
1:A:144:GLU:O	1:A:148:GLU:HB2	0.92	1.64	5	4
1:A:124:LYS:O	1:A:128:ARG:HB2	0.88	1.69	10	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:160:GLN:O	1:A:164:ARG:HB2	0.77	1.80	1	1
1:A:108:LYS:O	1:A:111:LYS:HB2	0.75	1.82	2	5
1:A:109:ALA:HB2	2:B:12:ARG:HG2	0.75	1.59	2	2
1:A:120:LYS:HG3	1:A:157:VAL:O	0.73	1.83	9	2
1:A:109:ALA:HB2	2:B:12:ARG:CB	0.71	2.15	11	4
1:A:109:ALA:HB2	2:B:12:ARG:HB2	0.65	1.66	11	1
1:A:109:ALA:HB2	2:B:12:ARG:HB3	0.62	1.70	1	5
1:A:129:VAL:HG23	2:B:9:LEU:HD11	0.61	1.73	3	7
1:A:120:LYS:CG	1:A:157:VAL:O	0.59	2.51	9	2
1:A:124:LYS:O	1:A:128:ARG:HG2	0.59	1.97	7	1
2:B:9:LEU:HA	2:B:11:ILE:HG22	0.56	1.77	12	20
1:A:124:LYS:O	1:A:128:ARG:HG3	0.56	2.00	14	2
1:A:120:LYS:CB	1:A:157:VAL:O	0.54	2.56	18	2
1:A:109:ALA:HB1	1:A:162:PHE:CZ	0.53	2.38	2	1
1:A:124:LYS:O	1:A:128:ARG:CG	0.51	2.59	18	2
1:A:113:PHE:HB3	1:A:121:ILE:HG21	0.51	1.81	8	2
1:A:109:ALA:CB	2:B:12:ARG:HB3	0.51	2.35	18	2
1:A:112:LEU:HB3	1:A:129:VAL:HG21	0.49	1.84	20	1
1:A:109:ALA:HB1	1:A:162:PHE:CE2	0.49	2.43	2	2
1:A:112:LEU:HD13	2:B:12:ARG:CB	0.48	2.38	2	1
2:B:10:LEU:O	2:B:12:ARG:HG2	0.45	2.12	7	1
2:B:9:LEU:HD12	2:B:11:ILE:CG2	0.44	2.42	12	2
1:A:129:VAL:HG22	2:B:9:LEU:CD2	0.44	2.43	20	1
1:A:129:VAL:HG13	2:B:9:LEU:HD11	0.43	1.90	20	1
1:A:127:LYS:O	1:A:131:LYS:HB3	0.43	2.07	8	1
1:A:109:ALA:HB2	2:B:12:ARG:CG	0.42	2.44	15	1
1:A:129:VAL:CG2	2:B:9:LEU:HD11	0.42	2.42	3	4
1:A:163:LEU:O	1:A:167:LYS:HB2	0.41	2.16	18	1
1:A:141:GLU:O	1:A:145:MET:SD	0.40	2.78	18	1
1:A:129:VAL:CG1	2:B:9:LEU:HD11	0.40	2.47	20	1

## 6.3 Torsion angles ⓘ

### 6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	43/79 (54%)	38±1 (89±2%)	5±1 (11±2%)	0±0 (0±1%)	59	88
2	B	11/17 (65%)	9±0 (82±4%)	1±1 (7±5%)	1±0 (11±4%)	1	8
All	All	1080/1920 (56%)	943 (87%)	111 (10%)	26 (2%)	12	49

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	11	ILE	20
2	B	12	ARG	5
1	A	114	ASP	1

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	40/72 (56%)	35±2 (88±4%)	5±2 (12±4%)	10	52
2	B	9/15 (60%)	8±1 (86±12%)	1±1 (14±12%)	8	47
All	All	980/1740 (56%)	855 (87%)	125 (13%)	10	51

All 24 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	142	LEU	20
1	A	162	PHE	19
2	B	9	LEU	9
1	A	157	VAL	9
1	A	148	GLU	9
2	B	5	LEU	8
1	A	126	LEU	7
2	B	12	ARG	6
1	A	163	LEU	6
1	A	161	GLU	6
1	A	166	MET	5
1	A	165	ILE	4
2	B	10	LEU	3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	113	PHE	3
1	A	129	VAL	2
1	A	146	ILE	1
1	A	108	LYS	1
1	A	114	ASP	1
1	A	106	ILE	1
1	A	133	LEU	1
1	A	125	ASN	1
1	A	159	GLU	1
1	A	121	ILE	1
1	A	141	GLU	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 70% for the well-defined parts and 67% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 5992

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	895
Number of shifts mapped to atoms	895
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	77	$1.33 \pm 0.33$	Should be applied
$^{13}\text{C}_\beta$	73	$1.55 \pm 0.22$	Should be applied
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	77	$-0.23 \pm 0.54$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 65%, i.e. 490 atoms were assigned a chemical shift out of a possible 758. 8 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	172/270 (64%)	86/108 (80%)	43/108 (40%)	43/54 (80%)
Sidechain	298/440 (68%)	189/258 (73%)	107/161 (66%)	2/21 (10%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	20/48 (42%)	20/26 (77%)	0/21 (0%)	0/1 (0%)
Overall	490/758 (65%)	295/392 (75%)	150/290 (52%)	45/76 (59%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 64%, i.e. 807 atoms were assigned a chemical shift out of a possible 1267. 10 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	309/480 (64%)	155/192 (81%)	77/192 (40%)	77/96 (80%)
Sidechain	474/731 (65%)	305/427 (71%)	165/266 (62%)	4/38 (11%)
Aromatic	24/56 (43%)	24/30 (80%)	0/25 (0%)	0/1 (0%)
Overall	807/1267 (64%)	484/649 (75%)	242/483 (50%)	81/135 (60%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	117	GLU	CG	25.50	42.24 – 29.94	-8.6
1	A	159	GLU	CG	26.60	42.24 – 29.94	-7.7
1	A	111	LYS	CD	22.80	34.86 – 23.06	-5.2
1	A	100	LYS	CD	22.80	34.86 – 23.06	-5.2

#### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



## 7.2 Chemical shift list 2

File name: BMRB entry 5992

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	52
Number of shifts mapped to atoms	52
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

### 7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 5%, i.e. 39 atoms were assigned a chemical shift out of a possible 758. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	19/270 (7%)	19/108 (18%)	0/108 (0%)	0/54 (0%)
Sidechain	14/440 (3%)	14/258 (5%)	0/161 (0%)	0/21 (0%)
Aromatic	6/48 (12%)	6/26 (23%)	0/21 (0%)	0/1 (0%)

*Continued on next page...*

Continued from previous page...

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Overall	39/758 (5%)	39/392 (10%)	0/290 (0%)	0/76 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 4%, i.e. 45 atoms were assigned a chemical shift out of a possible 1267. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	23/480 (5%)	23/192 (12%)	0/192 (0%)	0/96 (0%)
Sidechain	16/731 (2%)	16/427 (4%)	0/266 (0%)	0/38 (0%)
Aromatic	6/56 (11%)	6/30 (20%)	0/25 (0%)	0/1 (0%)
Overall	45/1267 (4%)	45/649 (7%)	0/483 (0%)	0/135 (0%)

## 7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

## 7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:

