



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:15 AM GMT

PDB ID : 3A4G
Title : Structure of cytochrome P450 vdh from Pseudonocardia autotrophica (trigonal crystal form)
Authors : Yasutake, Y.; Fujii, Y.; Cheon, W.K.; Arisawa, A.; Tamura, T.
Deposited on : 2009-07-07
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	411	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Vitamin D hydroxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	395	3062	1927	536	583	16	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	404	LEU	-	EXPRESSION TAG	UNP C4B644
A	405	GLU	-	EXPRESSION TAG	UNP C4B644
A	406	HIS	-	EXPRESSION TAG	UNP C4B644
A	407	HIS	-	EXPRESSION TAG	UNP C4B644
A	408	HIS	-	EXPRESSION TAG	UNP C4B644
A	409	HIS	-	EXPRESSION TAG	UNP C4B644
A	410	HIS	-	EXPRESSION TAG	UNP C4B644
A	411	HIS	-	EXPRESSION TAG	UNP C4B644

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 2-(2-METHOXYETHOXY)ETHANOL (three-letter code: PG0) (formula: $C_5H_{12}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	5	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	338	Total	O	0	0
			338	338		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vitamin D hydroxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	61.67Å 61.67Å 98.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.44 – 1.75 29.44 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.44-1.75) 99.3 (29.44-1.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.185 , 0.236 0.187 , 0.182	Depositor DCC
R_{free} test set	2120 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	22.1	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 52.2	EDS
Estimated twinning fraction	0.010 for -h,-k,l 0.038 for h,-h-k,-l 0.025 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 41973 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3451	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, PG0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/3128	0.80	2/4257 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	309	MET	CG-SD-CE	6.19	110.10	100.20
1	A	278	ARG	NE-CZ-NH1	5.30	122.95	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3062	0	3030	34	0
2	A	43	0	30	7	0
3	A	8	0	12	1	0
4	A	338	0	0	5	0
All	All	3451	0	3072	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:SER:HB2	1:A:176:PRO:HD2	1.08	1.06
1:A:175:SER:HB2	1:A:176:PRO:CD	1.91	1.00
1:A:175:SER:CB	1:A:176:PRO:HD2	2.01	0.90
1:A:214:SER:HA	1:A:219:ASP:O	1.79	0.81
2:A:412:HEM:HHC	2:A:412:HEM:HBB2	1.68	0.76
1:A:241:THR:HA	2:A:412:HEM:HBB1	1.75	0.69
1:A:40:LEU:HB3	1:A:41:PRO:HD2	1.82	0.61
1:A:40:LEU:HD12	1:A:45:VAL:CG2	2.30	0.60
1:A:40:LEU:HD12	1:A:45:VAL:HG21	1.83	0.59
1:A:36:ARG:HD2	1:A:37:LYS:O	2.03	0.58
1:A:37:LYS:HE2	1:A:44:PRO:HB2	1.86	0.57
1:A:211:LEU:O	1:A:220:ARG:NH2	2.38	0.55
2:A:412:HEM:HMC2	2:A:412:HEM:HBC2	1.88	0.55
1:A:254:THR:HG21	1:A:377:LEU:HD21	1.92	0.52
1:A:84:THR:O	1:A:86:MET:N	2.41	0.51
1:A:173:ASP:OD1	1:A:180:LYS:HE2	2.10	0.51
1:A:188:HIS:O	1:A:192:SER:HB2	2.12	0.49
3:A:420:PG0:H42	4:A:622:HOH:O	2.12	0.48
1:A:161:ARG:NH1	4:A:780:HOH:O	2.41	0.48
1:A:131:THR:HG23	1:A:399:GLY:C	2.35	0.46
2:A:412:HEM:CMC	2:A:412:HEM:HBC2	2.44	0.46
1:A:118:ILE:HD12	1:A:355:LEU:HD13	1.98	0.46
1:A:241:THR:OG1	2:A:412:HEM:HBB2	2.16	0.46
1:A:40:LEU:CD1	1:A:45:VAL:HG21	2.46	0.45
1:A:170:VAL:O	1:A:180:LYS:HG2	2.16	0.45
1:A:241:THR:HA	2:A:412:HEM:CBB	2.45	0.44
1:A:334:ALA:HA	4:A:791:HOH:O	2.17	0.43
1:A:363:ARG:HD3	4:A:835:HOH:O	2.17	0.43
1:A:215:ASP:OD1	1:A:215:ASP:C	2.57	0.43
1:A:157:PRO:HD2	1:A:190:TYR:OH	2.18	0.42
1:A:104:ARG:HD2	1:A:213:VAL:HG13	2.02	0.42
1:A:245:ILE:O	1:A:249:VAL:HG23	2.19	0.42
1:A:40:LEU:HD12	1:A:45:VAL:HG22	2.01	0.42
1:A:368:ARG:HG3	1:A:398:MET:HE3	2.02	0.42
1:A:349:GLY:HA3	2:A:412:HEM:C3C	2.55	0.41
1:A:215:ASP:HB2	4:A:669:HOH:O	2.19	0.41
1:A:40:LEU:CD1	1:A:45:VAL:CG2	2.96	0.41
1:A:179:ASP:N	1:A:179:ASP:OD2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	393/411 (96%)	376 (96%)	15 (4%)	2 (0%)	34 14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	334	ALA
1	A	175	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	329/343 (96%)	315 (96%)	14 (4%)	35 12

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ARG
1	A	74	ASP
1	A	87	MET
1	A	104	ARG
1	A	135	VAL
1	A	173	ASP
1	A	178	ASP
1	A	179	ASP
1	A	181	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	192	SER
1	A	194	LEU
1	A	195	LEU
1	A	223	GLN
1	A	355	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	ASN
1	A	223	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	412	1,4	30,50,50	2.35	8 (26%)	24,82,82	2.56	11 (45%)
3	PG0	A	420	-	7,7,7	0.69	0	6,6,6	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	412	1,4	-	0/10/54/54	0/0/8/8
3	PG0	A	420	-	-	0/5/5/5	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	412	HEM	C3B-C4B	-8.65	1.44	1.51
2	A	412	HEM	C3D-C4D	-5.18	1.44	1.51
2	A	412	HEM	C2C-C1C	-3.15	1.46	1.52
2	A	412	HEM	C2D-C1D	-2.36	1.44	1.51
2	A	412	HEM	C3C-CAC	2.01	1.55	1.51
2	A	412	HEM	FE-ND	2.03	2.08	1.97
2	A	412	HEM	C3B-CAB	2.78	1.56	1.51
2	A	412	HEM	FE-NC	3.48	2.09	1.95

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	412	HEM	CAA-C2A-C1A	-4.30	122.33	127.01
2	A	412	HEM	CMA-C3A-C4A	-2.35	124.48	128.36
2	A	412	HEM	C3B-CAB-CBB	-2.12	121.20	124.46
2	A	412	HEM	C3B-C4B-CHC	2.13	126.17	123.16
2	A	412	HEM	C2C-C1C-CHC	2.26	127.13	123.68
2	A	412	HEM	C2D-C3D-C4D	2.65	105.99	101.50
2	A	412	HEM	CMD-C2D-C3D	3.14	128.24	114.35
2	A	412	HEM	CAD-C3D-C4D	4.14	127.06	112.47
2	A	412	HEM	CMB-C2B-C3B	4.61	128.05	116.53
2	A	412	HEM	CMC-C2C-C3C	4.72	128.31	116.53
2	A	412	HEM	CAD-C3D-C2D	4.77	126.93	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	412	HEM	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	420	PG0	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/411 (96%)	0.60	49 (12%) 5 7	12, 24, 59, 82	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	170	VAL	9.3
1	A	167	TRP	9.2
1	A	176	PRO	8.3
1	A	77	ALA	7.9
1	A	174	ASP	6.8
1	A	80	PRO	6.7
1	A	81	ALA	6.1
1	A	78	ASP	5.8
1	A	175	SER	5.7
1	A	182	ALA	5.7
1	A	179	ASP	5.4
1	A	83	PRO	5.2
1	A	178	ASP	5.2
1	A	9	GLU	4.8
1	A	334	ALA	4.7
1	A	74	ASP	4.7
1	A	162	ASP	4.5
1	A	82	THR	4.4
1	A	181	ASN	4.2
1	A	79	MET	4.1
1	A	73	GLU	3.8
1	A	177	ALA	3.8
1	A	213	VAL	3.6
1	A	180	LYS	3.5
1	A	173	ASP	3.4
1	A	185	GLY	3.4
1	A	84	THR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	184	MET	3.1
1	A	216	GLU	3.0
1	A	350	ALA	2.8
1	A	166	ALA	2.8
1	A	244	LEU	2.7
1	A	143	PHE	2.6
1	A	236	ALA	2.5
1	A	333	ASP	2.5
1	A	212	ALA	2.5
1	A	172	VAL	2.4
1	A	197	ARG	2.4
1	A	193	ASP	2.3
1	A	217	ASP	2.3
1	A	403	ALA	2.2
1	A	171	LEU	2.2
1	A	348	LEU	2.2
1	A	215	ASP	2.2
1	A	67	TRP	2.1
1	A	117	ARG	2.1
1	A	233	LEU	2.0
1	A	186	LYS	2.0
1	A	335	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	PG0	A	420	8/8	0.79	0.18	1.65	31,35,37,38	0
2	HEM	A	412	43/43	0.98	0.15	0.84	11,14,20,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.