



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2017 – 11:58 PM EST

PDB ID : 5A4U  
Title : AtGSTF2 from Arabidopsis thaliana in complex with indole-3-aldehyde  
Authors : Ahmad, L.; Rylott, E.; Bruce, N.C.; Edwards, R.; Grogan, G.  
Deposited on : 2015-06-15  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	<div> <div>91%</div> <div>7% .</div> </div>
1	B	212	<div> <div>88%</div> <div>8% ...</div> </div>
1	C	212	<div> <div>90%</div> <div>8% .</div> </div>
1	D	212	<div> <div>90%</div> <div>7% .</div> </div>
1	E	212	<div> <div>%</div> <div>90%</div> <div>6% .</div> </div>
1	F	212	<div> <div>87%</div> <div>8% . .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	I3A	C	1213	-	-	-	X
2	I3A	C	1214	-	-	-	X
2	I3A	E	1213	-	-	-	X
2	I3A	F	1213	-	-	-	X
2	I3A	F	1214	-	-	-	X

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 10438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTATHIONE S-TRANSFERASE F2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1667	1072	280	313	2			
1	B	210	Total	C	N	O	S	0	1	0
			1668	1073	280	313	2			
1	C	211	Total	C	N	O	S	0	0	0
			1661	1069	282	308	2			
1	D	211	Total	C	N	O	S	0	1	0
			1652	1062	280	308	2			
1	E	211	Total	C	N	O	S	0	0	0
			1648	1062	274	310	2			
1	F	206	Total	C	N	O	S	0	2	0
			1630	1050	275	303	2			

- Molecule 2 is 1H-INDOLE-3-CARBALDEHYDE (three-letter code: I3A) (formula: C<sub>9</sub>H<sub>7</sub>NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			11	9	1	1		
2	B	1	Total	C	N	O	0	0
			11	9	1	1		
2	C	1	Total	C	N	O	0	0
			11	9	1	1		
2	C	1	Total	C	N	O	0	0
			11	9	1	1		
2	D	1	Total	C	N	O	0	0
			11	9	1	1		
2	E	1	Total	C	N	O	0	0
			11	9	1	1		
2	F	1	Total	C	N	O	0	0
			11	9	1	1		
2	F	1	Total	C	N	O	0	0
			11	9	1	1		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	E	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	97	Total	O	0	0
			97	97		
4	B	70	Total	O	0	0
			70	70		
4	C	64	Total	O	0	0
			64	64		
4	D	57	Total	O	0	0
			57	57		
4	E	54	Total	O	0	0
			54	54		
4	F	62	Total	O	0	0
			62	62		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GLUTATHIONE S-TRANSFERASE F2

Chain A: 




- Molecule 1: GLUTATHIONE S-TRANSFERASE F2

Chain B: 



- Molecule 1: GLUTATHIONE S-TRANSFERASE F2

Chain C: 




- Molecule 1: GLUTATHIONE S-TRANSFERASE F2

Chain D: 




- Molecule 1: GLUTATHIONE S-TRANSFERASE F2

Chain E: 



- Molecule 1: GLUTATHIONE S-TRANSFERASE F2

Chain F: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.86Å 94.41Å 152.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.38 – 2.00 80.25 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (80.38-2.00) 100.0 (80.25-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.198 , 0.234 0.206 , 0.241	Depositor DCC
$R_{free}$ test set	4311 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.9	Xtriage
Anisotropy	0.567	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10438	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: I3A, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.85	1/1704 (0.1%)	1.02	9/2311 (0.4%)
1	B	0.84	0/1709	1.00	9/2318 (0.4%)
1	C	0.82	1/1698 (0.1%)	1.02	8/2303 (0.3%)
1	D	0.83	1/1693 (0.1%)	0.98	7/2302 (0.3%)
1	E	0.80	0/1685	1.05	10/2290 (0.4%)
1	F	0.85	0/1670	1.04	12/2267 (0.5%)
All	All	0.83	3/10159 (0.0%)	1.02	55/13791 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	205	ARG	CD-NE	-5.53	1.37	1.46
1	C	139	GLU	CG-CD	5.41	1.60	1.51
1	D	205	ARG	CD-NE	-5.23	1.37	1.46

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	205	ARG	NE-CZ-NH2	-14.49	113.06	120.30
1	B	205	ARG	NE-CZ-NH2	-14.45	113.08	120.30
1	F	205	ARG	NE-CZ-NH2	-14.34	113.13	120.30
1	E	205	ARG	NE-CZ-NH2	-14.21	113.19	120.30
1	A	205	ARG	NE-CZ-NH2	-14.08	113.26	120.30
1	E	17	ARG	NE-CZ-NH2	-13.17	113.71	120.30
1	C	205	ARG	NE-CZ-NH2	-12.97	113.82	120.30
1	A	205	ARG	NE-CZ-NH1	12.72	126.66	120.30
1	B	205	ARG	NE-CZ-NH1	12.60	126.60	120.30
1	C	17	ARG	NE-CZ-NH2	-11.82	114.39	120.30
1	C	205	ARG	NE-CZ-NH1	11.61	126.10	120.30
1	E	205	ARG	NE-CZ-NH1	11.57	126.08	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	17	ARG	NE-CZ-NH1	11.02	125.81	120.30
1	D	205	ARG	NE-CZ-NH1	10.89	125.75	120.30
1	F	205	ARG	NE-CZ-NH1	10.33	125.46	120.30
1	A	17	ARG	NE-CZ-NH2	-9.38	115.61	120.30
1	C	17	ARG	NE-CZ-NH1	9.35	124.97	120.30
1	A	17	ARG	NE-CZ-NH1	9.34	124.97	120.30
1	D	17	ARG	NE-CZ-NH2	-8.96	115.82	120.30
1	F	17	ARG	NE-CZ-NH2	-8.45	116.08	120.30
1	B	17	ARG	NE-CZ-NH2	-7.53	116.53	120.30
1	F	17	ARG	NE-CZ-NH1	6.46	123.53	120.30
1	F	118	LEU	CB-CA-C	-6.42	97.99	110.20
1	B	127	ILE	N-CA-C	-6.42	93.66	111.00
1	A	17	ARG	CD-NE-CZ	6.42	132.59	123.60
1	B	128	TYR	N-CA-CB	6.33	121.99	110.60
1	E	124	PHE	CB-CA-C	6.29	122.99	110.40
1	F	120	PHE	CB-CA-C	6.27	122.93	110.40
1	D	35	GLU	CG-CD-OE2	-6.16	105.97	118.30
1	B	78	ARG	CG-CD-NE	5.94	124.28	111.80
1	C	128	TYR	CA-C-N	5.92	128.04	116.20
1	A	17	ARG	CG-CD-NE	5.89	124.17	111.80
1	D	17	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	E	3	GLY	N-CA-C	-5.81	98.56	113.10
1	C	17	ARG	CD-NE-CZ	5.80	131.72	123.60
1	A	69	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	82	GLN	CB-CA-C	-5.63	99.13	110.40
1	F	78	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	F	36	LEU	CA-CB-CG	5.56	128.09	115.30
1	E	55	VAL	N-CA-CB	-5.49	99.43	111.50
1	B	36	LEU	CA-CB-CG	5.48	127.90	115.30
1	A	140	GLU	OE1-CD-OE2	-5.46	116.75	123.30
1	E	190	THR	N-CA-CB	-5.42	99.99	110.30
1	C	55	VAL	N-CA-CB	-5.42	99.57	111.50
1	D	212	GLN	CA-C-O	5.40	131.44	120.10
1	D	49	ARG	NE-CZ-NH1	5.39	123.00	120.30
1	B	49	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	F	78	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	E	140	GLU	OE1-CD-OE2	-5.24	117.01	123.30
1	E	17	ARG	CD-NE-CZ	5.23	130.93	123.60
1	F	205	ARG	CG-CD-NE	-5.16	100.97	111.80
1	B	128	TYR	N-CA-C	-5.09	97.25	111.00
1	C	204	LYS	CB-CA-C	5.09	120.57	110.40
1	F	191	GLU	CB-CA-C	-5.05	100.29	110.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	205	ARG	CD-NE-CZ	5.04	130.66	123.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1667	0	1625	10	0
1	B	1668	0	1627	14	0
1	C	1661	0	1626	13	0
1	D	1652	0	1589	14	0
1	E	1648	0	1582	12	0
1	F	1630	0	1559	13	0
2	A	11	0	7	0	0
2	B	11	0	7	0	0
2	C	22	0	14	1	0
2	D	11	0	7	0	0
2	E	11	0	7	0	0
2	F	22	0	14	1	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
3	C	4	0	3	0	0
3	E	4	0	3	0	0
3	F	4	0	3	0	0
4	A	97	0	0	1	0
4	B	70	0	0	4	0
4	C	64	0	0	3	0
4	D	57	0	0	2	0
4	E	54	0	0	2	0
4	F	62	0	0	1	0
All	All	10438	0	9679	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:GLN:HG3	4:B:2045:HOH:O	1.61	0.98
1:A:77:HIS:HE1	1:A:97:TYR:OH	1.74	0.70
1:E:77:HIS:HE1	1:E:97:TYR:OH	1.74	0.70
1:F:77:HIS:HE1	1:F:97:TYR:OH	1.75	0.68
1:D:77:HIS:HE1	1:D:97:TYR:OH	1.75	0.68
1:E:81:ASN:HD22	1:E:81:ASN:C	1.96	0.67
1:B:77:HIS:HE1	1:B:97:TYR:OH	1.77	0.67
1:C:88:GLN:HG2	1:C:96:GLN:HB3	1.77	0.67
1:D:33[A]:HIS:NE2	1:D:35:GLU:OE1	2.28	0.66
1:C:117:LYS:CE	4:C:2059:HOH:O	2.43	0.66
1:E:88:GLN:HG2	1:E:96:GLN:HB3	1.78	0.65
1:D:81:ASN:C	1:D:81:ASN:HD22	1.99	0.65
1:C:77:HIS:HE1	1:C:97:TYR:OH	1.79	0.65
1:B:88:GLN:HG2	1:B:96:GLN:HB3	1.79	0.65
1:F:88:GLN:HG2	1:F:96:GLN:HB3	1.79	0.64
1:D:88:GLN:HG2	1:D:96:GLN:HB3	1.79	0.64
1:E:17:ARG:NH2	4:E:2007:HOH:O	2.16	0.64
1:C:9:HIS:CD2	1:C:11:ALA:H	2.15	0.63
1:C:9:HIS:HD2	1:C:11:ALA:H	1.48	0.61
1:A:88:GLN:HG2	1:A:96:GLN:HB3	1.83	0.60
1:B:36:LEU:HB3	1:C:127:ILE:HG12	1.84	0.59
1:A:124:PHE:CD2	1:E:128:TYR:HE1	2.23	0.56
1:C:88:GLN:NE2	4:C:2041:HOH:O	2.34	0.56
1:A:124:PHE:CD2	1:E:128:TYR:CE1	2.94	0.55
1:D:120:PHE:HZ	1:F:124:PHE:CD2	2.27	0.52
1:B:165:THR:HG23	4:B:2034:HOH:O	2.09	0.51
1:D:131:THR:HG22	1:F:41:HIS:NE2	2.25	0.51
1:E:121:GLU:O	1:E:132:THR:HB	2.10	0.51
1:B:124:PHE:O	1:B:127:ILE:O	2.29	0.51
1:C:121:GLU:O	1:C:132:THR:HB	2.11	0.51
1:B:165:THR:HG22	4:B:2037:HOH:O	2.11	0.50
1:B:121:GLU:O	1:B:132:THR:HB	2.12	0.50
1:E:108:ASP:OD1	1:F:109:HIS:HE1	1.95	0.50
1:D:121:GLU:O	1:D:132:THR:HB	2.11	0.49
1:A:121:GLU:O	1:A:132:THR:HB	2.13	0.49
1:A:108:ASP:OD1	1:B:109:HIS:HE1	1.96	0.49
1:B:127:ILE:O	1:B:128:TYR:CD2	2.66	0.48
1:C:17:ARG:NH2	4:C:2010:HOH:O	2.26	0.47
1:C:98:ALA:HB2	2:C:1214:ALA:H5	1.96	0.47
1:D:81:ASN:C	1:D:81:ASN:ND2	2.68	0.47
1:F:119:ALA:HB2	1:F:180:LEU:CD2	2.45	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:160:TYR:OH	1:C:201:GLU:CD	2.53	0.47
1:E:127:ILE:C	1:E:128:TYR:O	2.53	0.46
1:E:24:GLU:OE2	1:E:205:ARG:HD3	2.15	0.46
1:A:77:HIS:CE1	1:A:97:TYR:OH	2.63	0.46
1:F:77:HIS:CE1	1:F:97:TYR:OH	2.64	0.46
1:D:24:GLU:OE2	1:D:205:ARG:HD3	2.15	0.46
1:E:77:HIS:CE1	1:E:97:TYR:OH	2.63	0.46
1:F:24:GLU:OE2	1:F:205:ARG:HD3	2.17	0.45
1:F:98:ALA:HB2	2:F:1214:I3A:H5	1.98	0.45
1:C:24:GLU:OE2	1:C:205:ARG:HD3	2.17	0.45
1:D:88:GLN:HB2	4:D:2028:HOH:O	2.17	0.45
1:F:88:GLN:HB2	4:F:2039:HOH:O	2.18	0.44
1:E:88:GLN:HB2	4:E:2029:HOH:O	2.18	0.43
1:A:117:LYS:HE2	4:A:2069:HOH:O	2.17	0.43
1:A:24:GLU:OE2	1:A:205:ARG:HD3	2.18	0.43
1:B:37:LYS:HA	1:C:127:ILE:HD11	2.01	0.43
1:F:88:GLN:HA	1:F:88:GLN:HE21	1.83	0.43
1:D:77:HIS:CE1	1:D:97:TYR:OH	2.65	0.43
1:B:165:THR:CG2	4:B:2034:HOH:O	2.65	0.42
1:B:24:GLU:OE2	1:B:205:ARG:HD3	2.19	0.42
1:D:165:THR:HG22	4:D:2029:HOH:O	2.20	0.42
1:B:126:SER:O	1:B:128:TYR:O	2.37	0.42
1:D:126:SER:OG	1:F:37:LYS:HB3	2.20	0.42
1:D:131:THR:CG2	1:F:41:HIS:NE2	2.83	0.42
1:B:44:GLU:OE1	1:B:48:SER:HB3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/212 (99%)	207 (99%)	2 (1%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	209/212 (99%)	205 (98%)	3 (1%)	1 (0%)	34	26
1	C	209/212 (99%)	206 (99%)	3 (1%)	0	100	100
1	D	210/212 (99%)	208 (99%)	2 (1%)	0	100	100
1	E	209/212 (99%)	206 (99%)	2 (1%)	1 (0%)	34	26
1	F	204/212 (96%)	203 (100%)	0	1 (0%)	34	26
All	All	1250/1272 (98%)	1235 (99%)	12 (1%)	3 (0%)	52	48

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	128	TYR
1	B	128	TYR
1	F	67	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/182 (94%)	166 (97%)	5 (3%)	50	49
1	B	173/182 (95%)	167 (96%)	6 (4%)	43	40
1	C	170/182 (93%)	166 (98%)	4 (2%)	57	58
1	D	167/182 (92%)	162 (97%)	5 (3%)	48	47
1	E	165/182 (91%)	159 (96%)	6 (4%)	42	39
1	F	163/182 (90%)	158 (97%)	5 (3%)	47	46
All	All	1009/1092 (92%)	978 (97%)	31 (3%)	47	46

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	SER
1	A	17	ARG
1	A	88	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	132	THR
1	A	187	LYS
1	B	12	SER
1	B	36	LEU
1	B	82	GLN
1	B	88	GLN
1	B	127	ILE
1	B	132	THR
1	C	12	SER
1	C	55	VAL
1	C	88	GLN
1	C	132	THR
1	D	12	SER
1	D	36	LEU
1	D	81	ASN
1	D	88	GLN
1	D	132	THR
1	E	12	SER
1	E	32	VAL
1	E	55	VAL
1	E	81	ASN
1	E	88	GLN
1	E	132	THR
1	F	12	SER
1	F	36	LEU
1	F	37	LYS
1	F	82	GLN
1	F	88	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	77	HIS
1	A	88	GLN
1	A	122	GLN
1	A	212	GLN
1	B	77	HIS
1	B	88	GLN
1	B	109	HIS
1	B	122	GLN
1	C	9	HIS

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	C	77	HIS
1	C	81	ASN
1	C	122	GLN
1	D	77	HIS
1	D	81	ASN
1	D	88	GLN
1	D	109	HIS
1	D	122	GLN
1	E	26	ASN
1	E	77	HIS
1	E	81	ASN
1	E	82	GLN
1	E	88	GLN
1	E	122	GLN
1	F	77	HIS
1	F	88	GLN
1	F	109	HIS
1	F	122	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	I3A	A	1213	-	11,12,12	2.36	6 (54%)	11,16,16	1.03	0
3	ACT	A	1214	-	0,3,3	0.00	-	0,3,3	0.00	-
2	I3A	B	1213	-	11,12,12	2.16	5 (45%)	11,16,16	1.09	2 (18%)
3	ACT	B	1214	-	0,3,3	0.00	-	0,3,3	0.00	-
2	I3A	C	1213	-	11,12,12	2.18	4 (36%)	11,16,16	1.35	2 (18%)
2	I3A	C	1214	-	11,12,12	2.01	5 (45%)	11,16,16	1.21	1 (9%)
3	ACT	C	1215	-	0,3,3	0.00	-	0,3,3	0.00	-
2	I3A	D	1213	-	11,12,12	2.46	6 (54%)	11,16,16	1.46	3 (27%)
2	I3A	E	1213	-	11,12,12	1.99	4 (36%)	11,16,16	1.07	2 (18%)
3	ACT	E	1214	-	0,3,3	0.00	-	0,3,3	0.00	-
2	I3A	F	1213	-	11,12,12	2.25	6 (54%)	11,16,16	1.14	1 (9%)
2	I3A	F	1214	-	11,12,12	1.86	4 (36%)	11,16,16	1.23	1 (9%)
3	ACT	F	1215	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	I3A	A	1213	-	-	0/0/2/2	0/2/2/2
3	ACT	A	1214	-	-	0/0/0/0	0/0/0/0
2	I3A	B	1213	-	-	0/0/2/2	0/2/2/2
3	ACT	B	1214	-	-	0/0/0/0	0/0/0/0
2	I3A	C	1213	-	-	0/0/2/2	0/2/2/2
2	I3A	C	1214	-	-	0/0/2/2	0/2/2/2
3	ACT	C	1215	-	-	0/0/0/0	0/0/0/0
2	I3A	D	1213	-	-	0/0/2/2	0/2/2/2
2	I3A	E	1213	-	-	0/0/2/2	0/2/2/2
3	ACT	E	1214	-	-	0/0/0/0	0/0/0/0
2	I3A	F	1213	-	-	0/0/2/2	0/2/2/2
2	I3A	F	1214	-	-	0/0/2/2	0/2/2/2
3	ACT	F	1215	-	-	0/0/0/0	0/0/0/0

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1213	I3A	C7-C8	-4.47	1.33	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1213	I3A	C7-C8	-3.86	1.34	1.41
2	F	1213	I3A	C7-C8	-3.82	1.34	1.41
2	B	1213	I3A	C3-C3'	-3.69	1.35	1.46
2	A	1213	I3A	C3-C3'	-3.69	1.35	1.46
2	C	1213	I3A	C4-C9	-3.58	1.34	1.42
2	C	1213	I3A	C7-C8	-3.58	1.35	1.41
2	E	1213	I3A	C4-C9	-3.58	1.35	1.42
2	C	1214	I3A	C3-C3'	-3.53	1.35	1.46
2	D	1213	I3A	C3-C3'	-3.48	1.36	1.46
2	A	1213	I3A	C4-C9	-3.37	1.35	1.42
2	F	1214	I3A	C3-C3'	-3.30	1.36	1.46
2	F	1213	I3A	C3-C3'	-3.18	1.37	1.46
2	B	1213	I3A	C7-C8	-3.10	1.36	1.41
2	B	1213	I3A	C4-C9	-3.03	1.36	1.42
2	E	1213	I3A	C7-C8	-2.95	1.36	1.41
2	C	1213	I3A	C3-C3'	-2.93	1.37	1.46
2	C	1213	I3A	C3-C9	-2.84	1.34	1.41
2	C	1214	I3A	C4-C9	-2.83	1.36	1.42
2	E	1213	I3A	C3-C3'	-2.81	1.38	1.46
2	D	1213	I3A	C9-C8	-2.76	1.35	1.42
2	B	1213	I3A	C3-C9	-2.65	1.35	1.41
2	A	1213	I3A	C3-C9	-2.61	1.35	1.41
2	A	1213	I3A	C9-C8	-2.53	1.35	1.42
2	F	1213	I3A	C3-C9	-2.52	1.35	1.41
2	F	1213	I3A	C9-C8	-2.43	1.36	1.42
2	D	1213	I3A	C3-C9	-2.33	1.35	1.41
2	F	1213	I3A	C4-C9	-2.33	1.37	1.42
2	D	1213	I3A	C4-C9	-2.27	1.37	1.42
2	F	1214	I3A	C4-C9	-2.20	1.37	1.42
2	E	1213	I3A	C9-C8	-2.17	1.36	1.42
2	A	1213	I3A	C2-N	-2.16	1.32	1.36
2	C	1214	I3A	C9-C8	-2.14	1.36	1.42
2	F	1213	I3A	C8-N	-2.08	1.32	1.38
2	C	1214	I3A	C7-C8	-2.05	1.38	1.41
2	D	1213	I3A	C8-N	-2.02	1.32	1.38
2	B	1213	I3A	C9-C8	-2.00	1.37	1.42
2	F	1214	I3A	C6-C5	2.22	1.43	1.38
2	C	1214	I3A	C6-C7	2.43	1.42	1.36
2	F	1214	I3A	C6-C7	2.52	1.42	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1214	I3A	C5-C6-C7	-3.17	115.87	120.45
2	F	1214	I3A	C5-C6-C7	-3.11	115.96	120.45
2	E	1213	I3A	C4-C9-C3	-2.21	132.43	135.73
2	F	1213	I3A	C5-C4-C9	-2.15	117.89	120.88
2	B	1213	I3A	C4-C9-C3	-2.15	132.52	135.73
2	D	1213	I3A	C4-C9-C3	-2.14	132.54	135.73
2	D	1213	I3A	C5-C4-C9	-2.11	117.96	120.88
2	C	1213	I3A	C5-C6-C7	-2.03	117.52	120.45
2	E	1213	I3A	C9-C3-C3'	2.04	130.82	126.44
2	C	1213	I3A	C9-C3-C3'	2.25	131.26	126.44
2	B	1213	I3A	C9-C3-C3'	2.40	131.59	126.44
2	D	1213	I3A	C9-C3-C3'	2.60	132.01	126.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1214	I3A	1	0
2	F	1214	I3A	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/212 (99%)	-0.37	1 (0%) 91 92	19, 29, 42, 57	0
1	B	210/212 (99%)	-0.33	1 (0%) 91 92	19, 30, 47, 61	0
1	C	211/212 (99%)	-0.18	1 (0%) 91 92	20, 35, 54, 65	0
1	D	211/212 (99%)	-0.26	0 100 100	20, 34, 49, 60	0
1	E	211/212 (99%)	-0.13	2 (0%) 85 86	21, 36, 55, 74	0
1	F	206/212 (97%)	-0.35	1 (0%) 91 92	19, 30, 48, 65	0
All	All	1260/1272 (99%)	-0.27	6 (0%) 91 92	19, 32, 50, 74	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	136	VAL	2.7
1	E	2	ALA	2.6
1	C	2	ALA	2.3
1	A	131	THR	2.2
1	E	128	TYR	2.2
1	B	127	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	I3A	F	1214	11/11	0.74	0.26	7.83	42,48,54,54	0
2	I3A	C	1214	11/11	0.83	0.28	7.68	41,48,53,54	0
2	I3A	E	1213	11/11	0.91	0.15	3.38	31,33,36,39	0
2	I3A	C	1213	11/11	0.92	0.13	2.69	31,38,44,46	0
2	I3A	F	1213	11/11	0.94	0.11	2.17	28,30,34,35	0
2	I3A	D	1213	11/11	0.94	0.12	1.21	29,31,33,33	0
2	I3A	B	1213	11/11	0.94	0.11	0.67	25,27,31,33	0
3	ACT	E	1214	4/4	0.96	0.12	0.46	20,22,23,25	0
2	I3A	A	1213	11/11	0.95	0.10	0.33	28,29,31,33	0
3	ACT	C	1215	4/4	0.96	0.11	0.22	25,25,27,27	0
3	ACT	B	1214	4/4	0.95	0.10	-0.19	24,26,27,28	0
3	ACT	A	1214	4/4	0.99	0.09	-0.40	23,24,24,25	0
3	ACT	F	1215	4/4	0.97	0.08	-1.49	23,26,28,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.