



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:40 PM GMT

PDB ID : 4A57
Title : CRYSTAL STRUCTURE OF TOXOPLASMA GONDII NUCLEOSIDE
TRIPHOSPHATE DIPHOSPHOHYDROLASE 3 (NTPDASE3)
Authors : Krug, U.; Zebisch, M.; Straeter, N.
Deposited on : 2011-10-24
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	611	<div> <div>3%</div> <div>92%</div> <div>5%</div> <div>.</div> </div>
1	B	611	<div> <div>6%</div> <div>93%</div> <div>.</div> <div>.</div> </div>
1	C	611	<div> <div>2%</div> <div>94%</div> <div>.</div> <div>.</div> </div>
1	D	611	<div> <div>4%</div> <div>94%</div> <div>.</div> <div>.</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 19870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEOSIDE-TRIPHOSPHATASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	594	Total	C	N	O	S	0	6	0
			4640	2916	823	874	27			
1	B	594	Total	C	N	O	S	0	3	0
			4617	2905	813	872	27			
1	C	591	Total	C	N	O	S	0	4	0
			4611	2898	817	869	27			
1	D	595	Total	C	N	O	S	0	2	0
			4624	2905	817	875	27			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	EXPRESSION TAG	UNP Q27893
A	629	GLU	-	EXPRESSION TAG	UNP Q27893
A	630	HIS	-	EXPRESSION TAG	UNP Q27893
A	631	HIS	-	EXPRESSION TAG	UNP Q27893
A	632	HIS	-	EXPRESSION TAG	UNP Q27893
A	633	HIS	-	EXPRESSION TAG	UNP Q27893
A	634	HIS	-	EXPRESSION TAG	UNP Q27893
A	635	HIS	-	EXPRESSION TAG	UNP Q27893
B	25	MET	-	EXPRESSION TAG	UNP Q27893
B	629	GLU	-	EXPRESSION TAG	UNP Q27893
B	630	HIS	-	EXPRESSION TAG	UNP Q27893
B	631	HIS	-	EXPRESSION TAG	UNP Q27893
B	632	HIS	-	EXPRESSION TAG	UNP Q27893
B	633	HIS	-	EXPRESSION TAG	UNP Q27893
B	634	HIS	-	EXPRESSION TAG	UNP Q27893
B	635	HIS	-	EXPRESSION TAG	UNP Q27893
C	25	MET	-	EXPRESSION TAG	UNP Q27893
C	629	GLU	-	EXPRESSION TAG	UNP Q27893
C	630	HIS	-	EXPRESSION TAG	UNP Q27893
C	631	HIS	-	EXPRESSION TAG	UNP Q27893
C	632	HIS	-	EXPRESSION TAG	UNP Q27893

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	633	HIS	-	EXPRESSION TAG	UNP Q27893
C	634	HIS	-	EXPRESSION TAG	UNP Q27893
C	635	HIS	-	EXPRESSION TAG	UNP Q27893
D	25	MET	-	EXPRESSION TAG	UNP Q27893
D	629	GLU	-	EXPRESSION TAG	UNP Q27893
D	630	HIS	-	EXPRESSION TAG	UNP Q27893
D	631	HIS	-	EXPRESSION TAG	UNP Q27893
D	632	HIS	-	EXPRESSION TAG	UNP Q27893
D	633	HIS	-	EXPRESSION TAG	UNP Q27893
D	634	HIS	-	EXPRESSION TAG	UNP Q27893
D	635	HIS	-	EXPRESSION TAG	UNP Q27893

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Cl 3 3	0	0
2	C	1	Total Cl 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	451	Total O 452 452	0	1
3	B	306	Total O 307 307	0	1
3	C	366	Total O 366 366	0	0
3	D	248	Total O 249 249	0	1


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NUCLEOSIDE-TRIPHOSPHATASE 1




• Molecule 1: NUCLEOSIDE-TRIPHOSPHATASE 1



• Molecule 1: NUCLEOSIDE-TRIPHOSPHATASE 1



• Molecule 1: NUCLEOSIDE-TRIPHOSPHATASE 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.96Å 165.94Å 97.51Å 90.00° 97.03° 90.00°	Depositor
Resolution (Å)	39.43 – 2.00 39.43 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (39.43-2.00) 97.2 (39.43-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.00Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.174 , 0.210 0.179 , 0.220	Depositor DCC
R_{free} test set	1844 reflections (1.02%)	DCC
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.170	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 183317 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19870	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/4748	0.63	0/6424
1	B	0.49	0/4716	0.63	0/6384
1	C	0.50	0/4711	0.63	0/6374
1	D	0.49	0/4719	0.61	0/6386
All	All	0.50	0/18894	0.63	0/25568

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4640	0	4632	17	0
1	B	4617	0	4600	15	0
1	C	4611	0	4590	10	0
1	D	4624	0	4599	11	0
2	A	3	0	0	2	0
2	C	1	0	0	1	0
3	A	452	0	0	0	0
3	B	307	0	0	1	0
3	C	366	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	249	0	0	1	0
All	All	19870	0	18421	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:419:ASP:HB3	1:B:422:LYS:HD3	1.55	0.89
1:A:300[A]:ARG:HG3	1:B:464:ILE:HD12	1.68	0.76
1:A:487:MET:HG3	1:A:571:MET:HE1	1.72	0.70
1:A:476:GLN:HG3	2:A:1630:CL:CL	2.30	0.69
1:B:289:LEU:HD11	1:B:295[A]:LEU:HD21	1.75	0.67
1:A:113:ARG:HB2	1:A:190:GLY:O	2.00	0.62
1:B:185:LEU:HB3	1:B:228:THR:HG23	1.88	0.55
1:A:45:GLY:HA3	1:D:139:PHE:O	2.08	0.54
1:D:459:PRO:HD2	3:D:2182:HOH:O	2.09	0.53
1:B:45:GLY:HA3	1:C:139:PHE:O	2.11	0.50
1:D:243:THR:HG23	1:D:602:VAL:HB	1.95	0.48
1:A:185:LEU:HB3	1:A:228:THR:HG23	1.96	0.48
1:A:411:MET:HB3	1:A:411:MET:HE2	1.73	0.48
1:C:337:LEU:HD22	1:C:436:ILE:HD13	1.96	0.48
1:B:487:MET:HG3	1:B:571:MET:HE1	1.96	0.47
1:D:185:LEU:HB3	1:D:228:THR:HG23	1.95	0.47
1:C:123:TYR:OH	1:C:144:GLN:HG2	2.15	0.46
1:A:64:GLN:HB2	1:A:179:LEU:HD12	1.97	0.46
1:B:376:GLY:HA3	1:B:532:ILE:HD12	1.96	0.46
1:C:54:ASN:HB3	1:C:58:ARG:HE	1.81	0.46
1:C:191:VAL:O	1:C:199:ARG:HD3	2.16	0.46
1:C:476:GLN:HG3	2:C:1630:CL:CL	2.53	0.45
1:A:515:VAL:HG11	1:A:569:ARG:HG3	1.99	0.45
1:D:398:VAL:HG13	1:D:420:PRO:HB2	1.99	0.45
1:B:139:PHE:O	1:C:45:GLY:HA3	2.16	0.45
1:A:311:PRO:HB2	2:A:1631:CL:CL	2.54	0.45
1:A:392:LEU:HB3	1:A:418:ILE:HD13	1.99	0.45
1:B:390:LYS:HE2	1:B:424:LEU:HD21	1.99	0.44
1:D:187:SER:HG	1:D:203:PHE:HZ	1.64	0.44
1:D:370:VAL:HG22	1:D:380:VAL:HG22	2.00	0.43
1:A:123:TYR:HB3	1:A:141:TYR:CD2	2.53	0.43
1:A:474:VAL:HG11	1:A:567:LEU:CD2	2.49	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:411:MET:HB3	1:B:411:MET:HE2	1.78	0.43
1:B:393:ALA:HA	1:B:418:ILE:HG21	2.01	0.43
1:B:487:MET:HG3	1:B:571:MET:CE	2.49	0.43
1:A:139:PHE:O	1:D:45:GLY:HA3	2.19	0.43
1:D:337:LEU:HD22	1:D:436:ILE:HD13	2.00	0.43
1:D:333:ALA:HA	1:D:559:LYS:HE2	2.01	0.42
1:B:137:LEU:HD23	1:C:42:ILE:HD11	2.02	0.42
1:C:185:LEU:HB3	1:C:228:THR:HG23	2.02	0.42
1:C:243:THR:HG23	1:C:602:VAL:HB	2.01	0.41
1:D:182:PRO:HB3	1:D:227:TRP:CZ3	2.55	0.41
1:A:474:VAL:HG11	1:A:567:LEU:HD23	2.03	0.41
1:A:376:GLY:HA2	1:A:537:PRO:HG3	2.02	0.41
1:B:71:ALA:O	1:B:187:SER:HA	2.20	0.41
1:A:337:LEU:HD22	1:A:436:ILE:HD13	2.03	0.41
1:B:526:CYS:HB3	3:B:2175:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	598/611 (98%)	588 (98%)	8 (1%)	2 (0%)	46	41
1	B	595/611 (97%)	587 (99%)	8 (1%)	0	100	100
1	C	593/611 (97%)	584 (98%)	9 (2%)	0	100	100
1	D	595/611 (97%)	588 (99%)	7 (1%)	0	100	100
All	All	2381/2444 (97%)	2347 (99%)	32 (1%)	2 (0%)	56	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	128	TRP
1	A	297	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	511/521 (98%)	508 (99%)	3 (1%)	90	93
1	B	508/521 (98%)	501 (99%)	7 (1%)	74	77
1	C	507/521 (97%)	505 (100%)	2 (0%)	93	95
1	D	508/521 (98%)	506 (100%)	2 (0%)	93	95
All	All	2034/2084 (98%)	2020 (99%)	14 (1%)	90	91

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	366	SER
1	A	607	HIS
1	A	609	VAL
1	B	179[A]	LEU
1	B	179[B]	LEU
1	B	193	ASP
1	B	366	SER
1	B	404	SER
1	B	422	LYS
1	B	607	HIS
1	C	591	ASP
1	C	607	HIS
1	D	523	ARG
1	D	607	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	348	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	284	GLN
1	C	548	GLN
1	D	348	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	594/611 (97%)	-0.17	21 (3%)	48 49	13, 24, 55, 92	0
1	B	594/611 (97%)	0.02	34 (5%)	27 29	16, 30, 64, 126	0
1	C	591/611 (96%)	-0.23	10 (1%)	73 73	14, 25, 50, 88	0
1	D	595/611 (97%)	-0.12	23 (3%)	43 45	13, 31, 60, 81	0
All	All	2374/2444 (97%)	-0.13	88 (3%)	45 47	13, 28, 59, 126	0

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	132	PRO	5.9
1	C	590	PRO	5.4
1	A	592	GLY	5.2
1	B	137	LEU	5.1
1	A	590	PRO	4.6
1	A	258	CYS	4.5
1	A	591	ASP	4.4
1	B	126	LYS	4.3
1	B	591	ASP	4.3
1	B	590	PRO	4.2
1	A	260	ILE	4.1
1	B	125	GLY	4.1
1	B	262	GLU	3.9
1	B	265	VAL	3.8
1	B	37	ASP	3.8
1	B	259	MET	3.8
1	B	418	ILE	3.7
1	B	258	CYS	3.7
1	B	592	GLY	3.7
1	B	260	ILE	3.7
1	A	127	ASP	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	398	VAL	3.6
1	B	127	ASP	3.6
1	B	398	VAL	3.5
1	A	130	SER	3.5
1	B	133	VAL	3.4
1	A	131	ARG	3.4
1	B	131	ARG	3.3
1	B	419	ASP	3.3
1	B	535	ASP	3.2
1	D	265	VAL	3.0
1	D	374	PRO	3.0
1	B	36	ALA	3.0
1	C	591	ASP	3.0
1	A	266	LYS	3.0
1	D	418	ILE	2.9
1	A	256	ALA	2.9
1	C	593	LYS	2.9
1	A	259	MET	2.8
1	C	40	LYS	2.8
1	D	195	HIS	2.8
1	C	132	PRO	2.8
1	B	589	GLY	2.7
1	B	266	LYS	2.7
1	D	130	SER	2.7
1	B	124	ALA	2.7
1	B	397	SER	2.7
1	D	520	GLU	2.7
1	D	403	ILE	2.6
1	D	505	ARG	2.6
1	A	589	GLY	2.6
1	D	258	CYS	2.5
1	D	260	ILE	2.5
1	D	264	GLY	2.5
1	B	263	TYR	2.4
1	C	41	ARG	2.4
1	D	401	PRO	2.4
1	A	254	ASP	2.4
1	D	535	ASP	2.4
1	D	268	CYS	2.4
1	D	426	GLU	2.3
1	A	268	CYS	2.3
1	A	132	PRO	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	544	ALA	2.3
1	B	128	TRP	2.3
1	B	38	THR	2.3
1	A	588	ALA	2.3
1	A	37	ASP	2.3
1	A	265	VAL	2.3
1	B	268	CYS	2.2
1	A	593	LYS	2.2
1	C	44	VAL	2.2
1	D	294	VAL	2.2
1	D	397	SER	2.2
1	C	131	ARG	2.2
1	B	534	THR	2.2
1	C	265	VAL	2.1
1	A	137	LEU	2.1
1	B	417	SER	2.1
1	C	42	ILE	2.1
1	D	543	ASN	2.1
1	B	123	TYR	2.1
1	B	588	ALA	2.0
1	D	419	ASP	2.0
1	B	421	THR	2.0
1	D	132	PRO	2.0
1	A	124	ALA	2.0
1	D	524	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CL	C	1630	1/1	0.99	0.05	-2.63	34,34,34,34	0
2	CL	A	1630	1/1	0.99	0.05	-2.72	32,32,32,32	0
2	CL	A	1632	1/1	0.84	0.16	-	36,36,36,36	1
2	CL	A	1631	1/1	0.98	0.07	-	29,29,29,29	1

6.5 Other polymers [i](#)

There are no such residues in this entry.