



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:19 PM GMT

PDB ID : 1A6E  
Title : THERMOSOME-MG-ADP-ALF3 COMPLEX  
Authors : Ditzel, L.; Loewe, J.; Stock, D.; Stetter, K.-O.; Huber, H.; Huber, R.; Steinbacher, S.  
Deposited on : 1998-02-24  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	545	
2	B	543	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9390 atoms, of which 1742 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THERMOSOME (ALPHA SUBUNIT).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	503	Total	C	H	N	O	S	884	0	0
			4668	2356	884	662	752	14			

- Molecule 2 is a protein called THERMOSOME (BETA SUBUNIT).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	502	Total	C	H	N	O	S	858	0	0
			4656	2370	858	651	758	19			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total 27	C 10	N 5	O 10	P 2	0	0
4	A	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 5 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula:  $\text{AlF}_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	Al	F	0	0
			4	1	3		
5	A	1	Total	Al	F	0	0
			4	1	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	O 1	0	0
6	B	1	Total 1	O 1	0	0



I511	A424	E346	N272
M512	Q425	R347	N273
I513	K426	V348	L274
I514	I427	E349	R275
R515	R430	Q350	R276
I516	Q431	V351	N277
	Q432	K352	V278
V519	L433	V353	
I520		G354	I281
A521	E436	E355	
THR		D356	V284
LYS	D440	Y357	
SER	A441	K358	N287
SER	I442	T359	
SER	E443	F360	I290
SER	E444	V361	T291
SER	I445	T362	Q292
SER	P446	G363	K293
ASN			
PRO	A450	N366	D296
PRO		P367	D297
LYS	L455	K368	N298
SER	D456	A369	A299
GLY	P457	V370	Q300
SER	I458	S371	H301
SER	D459	I372	T302
SER	I460	L373	L303
GLU	L461	V374	S304
SER	L462	R375	R305
SER		G376	
GLU	M472		A310
ASP	K473	I389	V311
	T474	T390	R312
	Y475	D391	R313
	G476		V314
	I477	H394	K315
		V395	K316
	I484	V396	S317
	M487	A399	L322
	V488	L400	
	K489	E401	A325
	M490	D402	T326
	G491		G327
	V492	Y405	A328
		A406	S329
	P495	A407	I330
	I496	G408	V331
	R497	G409	S332
	V498		T333
	G499	T412	I334
	R500		
	Q501	E415	I337
	A502	I416	S338
	I503		R339
		L420	S340
	T507	R421	D341
		S422	L342
	A510	Y423	G343

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	167.80 Å   167.80 Å   202.30 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	8.00 – 3.20	Depositor
% Data completeness (in resolution range)	92.5 (8.00-3.20)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.181 , 0.285	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	9390	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP, AF3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/3812	0.74	2/5139 (0.0%)
2	B	0.52	0/3834	0.74	1/5166 (0.0%)
All	All	0.50	0/7646	0.74	3/10305 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	244	LEU	CA-CB-CG	5.21	127.28	115.30
1	A	57	ASP	N-CA-C	5.18	125.00	111.00
1	A	160	SER	N-CA-C	5.15	124.91	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3784	884	3935	191	0
2	B	3798	858	3892	195	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	27	0	12	1	0
4	B	27	0	12	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	4	0	0	0	0
5	B	4	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
All	All	7648	1742	7851	374	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 374 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ILE:HD11	1:A:321:LEU:HD11	1.22	1.11
2:B:172:LEU:HD22	2:B:389:ILE:HD11	1.46	0.96
1:A:281:LYS:HD2	1:A:305:GLU:HG3	1.48	0.95
2:B:50:LEU:HB2	2:B:58:VAL:HG13	1.48	0.93
2:B:64:VAL:HG22	2:B:95:THR:HG21	1.49	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	501/545 (92%)	450 (90%)	41 (8%)	10 (2%)	9	48
2	B	500/543 (92%)	463 (93%)	36 (7%)	1 (0%)	52	88
All	All	1001/1088 (92%)	913 (91%)	77 (8%)	11 (1%)	17	62

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	ASP
1	A	145	SER
1	A	377	THR
1	A	378	ASP
1	A	160	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/442 (93%)	375 (91%)	36 (9%)	12	45
2	B	410/446 (92%)	376 (92%)	34 (8%)	14	49
All	All	821/888 (92%)	751 (92%)	70 (8%)	13	47

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	467	ASP
2	B	67	LEU
2	B	462	LEU
1	A	478	LEU
1	A	516	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	424	ASN
1	A	480	ASN
2	B	394	HIS
1	A	431	GLN
1	A	451	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	ADP	A	898	3,5	22,29,29	1.20	3 (13%)	27,45,45	2.66	4 (14%)
5	AF3	A	899	3,4,6	0,3,3	0.00	-	0,3,3	0.00	-
4	ADP	B	998	3,5	22,29,29	1.16	3 (13%)	27,45,45	2.67	3 (11%)
5	AF3	B	999	3,4,6	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	898	3,5	-	0/12/32/32	0/3/3/3
5	AF3	A	899	3,4,6	-	0/0/0/0	0/0/0/0
4	ADP	B	998	3,5	-	0/12/32/32	0/3/3/3
5	AF3	B	999	3,4,6	-	0/0/0/0	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	898	ADP	C5-N7	-2.43	1.31	1.39
4	B	998	ADP	C5-N7	-2.22	1.31	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	898	ADP	O4'-C1'	2.42	1.44	1.41
4	B	998	ADP	O4'-C1'	2.66	1.44	1.41
4	B	998	ADP	PB-O2B	2.80	1.64	1.54

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	998	ADP	N3-C2-N1	-12.38	119.41	128.89
4	A	898	ADP	N3-C2-N1	-11.90	119.79	128.89
4	A	898	ADP	C2'-C1'-N9	-3.79	108.50	114.29
4	B	998	ADP	C2'-C1'-N9	-3.70	108.63	114.29
4	A	898	ADP	C4-C5-N7	-3.05	106.67	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	898	ADP	1	0
4	B	998	ADP	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.