



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:22 AM GMT

PDB ID : 2A8H
Title : Crystal structure of catalytic domain of TACE with Thiomorpholine Sulfonamide Hydroxamate inhibitor
Authors : Levin, J.I.; Chen, J.M.; Laakso, L.M.; Du, M.; Schmid, J.; Xu, W.; Cummons, T.; Xu, J.; Jin, G.; Barone, D.; Skotnicki, J.S.
Deposited on : 2005-07-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	 4% 80% 14% • •
1	B	271	 8% 72% 20% • 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4295 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADAM 17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			2054	1290	347	404	13			
1	B	257	Total	C	N	O	S	0	0	0
			2038	1282	344	399	13			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	266	ALA	SER	ENGINEERED	UNP P78536
A	452	GLN	ASN	ENGINEERED	UNP P78536
A	478	GLY	-	EXPRESSION TAG	UNP P78536
A	479	SER	-	EXPRESSION TAG	UNP P78536
A	480	HIS	-	EXPRESSION TAG	UNP P78536
A	481	HIS	-	EXPRESSION TAG	UNP P78536
A	482	HIS	-	EXPRESSION TAG	UNP P78536
A	483	HIS	-	EXPRESSION TAG	UNP P78536
A	484	HIS	-	EXPRESSION TAG	UNP P78536
A	485	HIS	-	EXPRESSION TAG	UNP P78536
B	266	ALA	SER	ENGINEERED	UNP P78536
B	452	GLN	ASN	ENGINEERED	UNP P78536
B	478	GLY	-	EXPRESSION TAG	UNP P78536
B	479	SER	-	EXPRESSION TAG	UNP P78536
B	480	HIS	-	EXPRESSION TAG	UNP P78536
B	481	HIS	-	EXPRESSION TAG	UNP P78536
B	482	HIS	-	EXPRESSION TAG	UNP P78536
B	483	HIS	-	EXPRESSION TAG	UNP P78536
B	484	HIS	-	EXPRESSION TAG	UNP P78536
B	485	HIS	-	EXPRESSION TAG	UNP P78536

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 4-({4-[(4-AMINO BUT-2-YNYL)OXY]PHENYL}SULFONYL)-N-HYDROXY-2,2-DIMETHYLTHIOMORPHOLINE-3-CARBOXAMIDE (three-letter code: 4NH) (formula: C₁₇H₂₃N₃O₅S₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			27	17	3	5	2		
3	B	1	Total	C	N	O	S	0	0
			27	17	3	5	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	91	Total	O	0	0
			91	91		
4	B	56	Total	O	0	0
			56	56		

- Molecule 1: ADAM 17



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.37Å 59.21Å 198.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 19.80 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-2.30) 99.1 (19.80-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11.88 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.281 0.216 , 0.279	Depositor DCC
R_{free} test set	1313 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 25906 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4295	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 4NH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2101	0.67	0/2837
1	B	0.54	0/2085	0.63	0/2814
All	All	0.62	0/4186	0.65	0/5651

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	372	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2054	0	1960	30	0
1	B	2038	0	1951	52	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	27	0	22	1	0
3	B	27	0	22	3	0
4	A	91	0	0	2	0
4	B	56	0	0	3	0
All	All	4295	0	3955	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:370:TYR:HA	1:B:371:SER:CB	1.52	1.37
1:B:370:TYR:CA	1:B:371:SER:HB2	1.68	1.21
1:B:371:SER:N	1:B:372:PRO:HD2	1.40	1.17
1:B:371:SER:OG	1:B:376:LYS:O	1.70	1.08
1:B:373:VAL:HG22	1:B:374:GLY:H	1.10	1.06
1:A:369:TYR:CE2	1:A:378:ILE:HG12	1.95	1.02
1:B:371:SER:H	1:B:372:PRO:HD2	1.19	1.01
1:A:378:ILE:HG13	1:A:379:TYR:N	1.77	0.99
1:A:283:ARG:HH11	1:A:283:ARG:HB2	1.31	0.95
1:B:278:GLN:HE22	1:B:474:SER:N	1.64	0.94
1:B:373:VAL:CG2	1:B:374:GLY:N	2.30	0.94
1:B:371:SER:N	1:B:372:PRO:CD	2.30	0.92
1:B:278:GLN:HE22	1:B:474:SER:H	0.94	0.92
1:B:373:VAL:HG22	1:B:374:GLY:N	1.75	0.92
1:B:278:GLN:NE2	1:B:474:SER:H	1.74	0.83
1:B:370:TYR:CA	1:B:371:SER:CB	2.37	0.76
1:A:415:HIS:HE1	3:A:158:4NH:O4	1.66	0.76
1:B:307:GLU:HG2	4:B:66:HOH:O	1.85	0.75
1:B:370:TYR:HA	1:B:371:SER:HB2	0.76	0.74
1:B:269:ASN:HD21	1:B:452:GLN:HE22	1.37	0.72
1:A:297:HIS:HD2	1:A:299:ASN:H	1.39	0.71
1:A:269:ASN:HD21	1:A:452:GLN:HE22	1.43	0.65
1:B:312:TRP:H	1:B:341:GLN:NE2	1.97	0.63
1:A:279:ILE:HG21	1:A:282:ILE:HG13	1.79	0.63
1:A:306:ASN:O	1:A:309:LYS:HG2	1.98	0.62
1:A:290:GLU:HB3	1:B:429:GLN:HG2	1.80	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:ARG:HH11	1:A:283:ARG:CB	2.09	0.62
1:A:297:HIS:CD2	1:A:299:ASN:H	2.17	0.61
1:B:428:ASP:N	1:B:428:ASP:OD1	2.27	0.61
1:A:378:ILE:HD11	1:A:380:LEU:CD2	2.32	0.60
1:B:415:HIS:CE1	3:B:159:4NH:O4	2.54	0.60
1:A:378:ILE:HD11	1:A:380:LEU:HD23	1.84	0.59
1:B:306:ASN:HD21	1:B:308:GLU:HG3	1.67	0.59
1:B:375:LYS:O	1:B:376:LYS:CB	2.51	0.58
1:B:361:HIS:HA	4:B:75:HOH:O	2.04	0.58
1:B:361:HIS:HB3	1:B:380:LEU:HD12	1.85	0.58
1:B:304:TYR:CD1	1:B:305:PRO:HA	2.39	0.57
1:B:439:ALA:HB1	3:B:159:4NH:HN21	1.70	0.56
1:A:234:ARG:O	1:A:238:TYR:HD1	1.88	0.56
1:B:409:HIS:CD2	3:B:159:4NH:O5	2.59	0.56
1:B:297:HIS:CD2	1:B:299:ASN:H	2.24	0.56
1:A:369:TYR:HE2	1:A:378:ILE:HG12	1.63	0.55
1:A:260:ASP:OD1	1:A:263:ARG:NH2	2.38	0.54
1:B:226:LYS:HB3	1:B:280:GLU:HB2	1.89	0.54
1:B:375:LYS:O	1:B:376:LYS:HB3	2.09	0.53
1:A:369:TYR:CZ	1:A:378:ILE:HG21	2.44	0.53
1:B:357:ARG:CZ	1:B:360:SER:HB2	2.39	0.53
1:A:436:TYR:CD1	1:A:437:PRO:HD2	2.45	0.52
1:B:370:TYR:HA	1:B:371:SER:HB3	1.75	0.50
1:B:306:ASN:HD22	1:B:306:ASN:C	2.14	0.50
1:A:289:GLN:HG2	1:A:297:HIS:CG	2.47	0.50
1:A:292:LYS:HB3	1:A:293:PRO:HD2	1.94	0.49
1:B:219:ASP:C	1:B:220:PRO:O	2.51	0.49
1:B:269:ASN:HD21	1:B:452:GLN:NE2	2.09	0.49
1:B:230:VAL:HG22	1:B:283:ARG:HB2	1.94	0.48
1:B:370:TYR:C	1:B:372:PRO:HD2	2.25	0.48
1:B:224:THR:HG21	1:B:278:GLN:HE21	1.77	0.48
1:B:255:ILE:HG23	1:B:277:ILE:HD12	1.95	0.47
1:B:220:PRO:C	1:B:222:LYS:H	2.17	0.47
1:B:312:TRP:H	1:B:341:GLN:HE21	1.61	0.46
1:A:348:LEU:HD21	1:A:389:ASN:HB2	1.98	0.46
1:B:330:SER:HB3	1:B:379:TYR:CZ	2.51	0.46
1:B:356:PRO:HD3	4:B:146:HOH:O	2.16	0.45
1:A:217:ASP:HB3	4:A:9:HOH:O	2.15	0.45
1:A:397:LYS:O	1:A:401:LEU:HG	2.17	0.45
1:B:389:ASN:O	1:B:392:LYS:HG3	2.17	0.44
1:B:373:VAL:HG23	1:B:374:GLY:N	2.27	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:416:ASP:HA	1:A:417:PRO:HD3	1.86	0.44
1:A:279:ILE:HG21	1:A:282:ILE:CG1	2.47	0.44
1:B:343:PHE:HB2	1:B:347:THR:HB	2.00	0.43
1:B:370:TYR:CG	1:B:371:SER:HB3	2.52	0.43
1:B:304:TYR:CG	1:B:305:PRO:HA	2.53	0.43
1:A:406:GLU:OE1	1:A:406:GLU:HA	2.18	0.43
1:B:371:SER:O	1:B:372:PRO:C	2.58	0.42
1:B:401:LEU:HD11	1:B:442:GLY:CA	2.49	0.42
1:A:306:ASN:HB2	1:A:309:LYS:HE2	2.02	0.42
1:B:352:TYR:HB3	1:B:362:GLY:HA2	2.02	0.42
1:A:369:TYR:O	1:A:378:ILE:HG23	2.20	0.41
1:A:308:GLU:HB2	4:A:21:HOH:O	2.21	0.41
1:B:458:ILE:HG22	1:B:462:ILE:HD12	2.01	0.41
1:A:378:ILE:HD11	1:A:380:LEU:HD21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/271 (95%)	247 (96%)	8 (3%)	2 (1%)	24	27
1	B	255/271 (94%)	240 (94%)	10 (4%)	5 (2%)	9	7
All	All	512/542 (94%)	487 (95%)	18 (4%)	7 (1%)	14	13

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	371	SER
1	B	372	PRO
1	B	376	LYS
1	A	312	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	365	CYS
1	B	365	CYS
1	B	220	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/232 (96%)	211 (95%)	11 (5%)	30	41
1	B	220/232 (95%)	207 (94%)	13 (6%)	24	32
All	All	442/464 (95%)	418 (95%)	24 (5%)	27	36

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	223	ASN
1	A	277	ILE
1	A	283	ARG
1	A	302	LYS
1	A	306	ASN
1	A	308	GLU
1	A	327	GLU
1	A	341	GLN
1	A	378	ILE
1	A	407	LEU
1	A	448	LYS
1	B	223	ASN
1	B	306	ASN
1	B	308	GLU
1	B	309	LYS
1	B	322	SER
1	B	392	LYS
1	B	407	LEU
1	B	414	GLU
1	B	420	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	422	GLU
1	B	426	ASN
1	B	428	ASP
1	B	460	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	ASN
1	A	297	HIS
1	A	341	GLN
1	A	452	GLN
1	B	223	ASN
1	B	278	GLN
1	B	297	HIS
1	B	306	ASN
1	B	320	GLN
1	B	341	GLN
1	B	426	ASN
1	B	429	GLN
1	B	452	GLN
1	B	471	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	4NH	A	158	2	26,28,28	2.17	3 (11%)	32,40,40	1.36	6 (18%)
3	4NH	B	159	2	26,28,28	2.17	3 (11%)	32,40,40	1.36	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4NH	A	158	2	-	0/21/41/41	0/1/2/2
3	4NH	B	159	2	-	0/21/41/41	0/1/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	159	4NH	C4-N1	2.84	1.52	1.48
3	A	158	4NH	C4-N1	2.89	1.52	1.48
3	B	159	4NH	C3-N1	4.22	1.52	1.48
3	A	158	4NH	C3-N1	4.29	1.52	1.48
3	A	158	4NH	S2-N1	9.25	1.76	1.63
3	B	159	4NH	S2-N1	9.28	1.76	1.63

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	158	4NH	C6-S2-N1	-3.14	102.24	107.38
3	B	159	4NH	C6-S2-N1	-3.13	102.27	107.38
3	B	159	4NH	O5-N3-C16	-2.88	115.80	119.86
3	A	158	4NH	O5-N3-C16	-2.87	115.81	119.86
3	B	159	4NH	O1-S2-C6	-2.49	104.77	108.00
3	A	158	4NH	O1-S2-C6	-2.48	104.79	108.00
3	A	158	4NH	C14-C15-N2	-2.40	109.56	113.08
3	B	159	4NH	C14-C15-N2	-2.39	109.59	113.08
3	B	159	4NH	O1-S2-N1	2.05	111.32	106.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	158	4NH	O1-S2-N1	2.05	111.33	106.97
3	A	158	4NH	C3-N1-S2	2.70	124.32	119.43
3	B	159	4NH	C3-N1-S2	2.71	124.32	119.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	158	4NH	1	0
3	B	159	4NH	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/271 (95%)	0.08	10 (3%) 43 52	12, 25, 43, 56	1 (0%)
1	B	257/271 (94%)	0.56	23 (8%) 12 17	16, 36, 59, 72	0
All	All	516/542 (95%)	0.32	33 (6%) 23 31	12, 30, 53, 72	1 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	373	VAL	10.3
1	B	372	PRO	5.3
1	B	359	ASN	4.3
1	B	369	TYR	4.3
1	B	370	TYR	4.3
1	A	293	PRO	4.2
1	B	361	HIS	4.1
1	B	375	LYS	3.7
1	B	294	GLY	3.4
1	A	291	VAL	3.3
1	B	360	SER	3.2
1	B	371	SER	3.0
1	B	270	ALA	2.8
1	B	391	GLY	2.8
1	A	428	ASP	2.8
1	B	292	LYS	2.8
1	B	295	GLU	2.8
1	A	308	GLU	2.7
1	B	293	PRO	2.6
1	B	390	TYR	2.5
1	B	221	MET	2.4
1	B	357	ARG	2.4
1	A	241	ARG	2.2
1	B	327	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	302	LYS	2.2
1	B	364	VAL	2.2
1	B	374	GLY	2.2
1	A	292	LYS	2.1
1	B	458	ILE	2.1
1	B	358	ALA	2.1
1	A	217	ASP	2.0
1	A	238	TYR	2.0
1	A	294	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	4NH	A	158	27/27	0.90	0.16	1.18	23,30,33,34	0
3	4NH	B	159	27/27	0.89	0.17	0.75	29,38,39,40	0
2	ZN	A	486	1/1	0.99	0.06	-	18,18,18,18	0
2	ZN	B	486	1/1	0.97	0.06	-	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.