



wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 02:09 PM BST

PDB ID : 5A9K
EMDB ID: : EMD-3087
Title : Structural basis for DNA strand separation by a hexameric replicative helicase
Authors : Chaban, Y.; Stead, J.A.; Ryzhenkova, K.; Whelan, F.; Lamber, K.; Antson, F.; Sanders, C.M.; Orlova, E.V.
Deposited on : 2015-07-21
Resolution : 19.00 Å(reported)
Based on PDB ID : 2V9P

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk27241


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 19.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726
Sidechain outliers	111093	686

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	305	69% 19% • 11%
1	B	305	71% 14% • 12%
1	C	305	70% 16% • 12%
1	D	305	73% 14% • 10%
1	E	305	68% 20% • 11%
1	F	305	64% 22% •• 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	1580	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 13059 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called REPLICATION PROTEIN E1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	272	Total	C	N	O	S	0	0
			2183	1404	375	394	10		
1	B	268	Total	C	N	O	S	0	0
			2149	1386	370	383	10		
1	C	267	Total	C	N	O	S	0	0
			2142	1382	369	381	10		
1	D	274	Total	C	N	O	S	0	0
			2191	1409	377	395	10		
1	E	270	Total	C	N	O	S	0	0
			2170	1398	373	389	10		
1	F	269	Total	C	N	O	S	0	0
			2158	1392	372	384	10		

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			AltConf
2	A	1	Total	O	P	0
			10	8	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			AltConf
2	A	1	Total	O	P	0
			10	8	2	
2	B	1	Total	O	P	0
			10	8	2	
2	B	1	Total	O	P	0
			10	8	2	
2	C	1	Total	O	P	0
			10	8	2	
2	C	1	Total	O	P	0
			10	8	2	
2	D	1	Total	O	P	0
			5	4	1	
2	E	1	Total	O	P	0
			5	4	1	
2	F	1	Total	O	P	0
			5	4	1	

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
3	B	1	Total	Mg	0
			1	1	
3	A	1	Total	Mg	0
			1	1	
3	C	1	Total	Mg	0
			1	1	


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		AltConf
4	A	3	Total	O	0
			3	3	
4	B	3	Total	O	0
			3	3	
4	C	3	Total	O	0
			3	3	
4	D	4	Total	O	0
			4	4	
4	E	2	Total	O	0
			2	2	
4	F	3	Total	O	0
			3	3	


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: REPLICATION PROTEIN E1



• Molecule 1: REPLICATION PROTEIN E1




• Molecule 1: REPLICATION PROTEIN E1




• Molecule 1: REPLICATION PROTEIN E1






• Molecule 1: REPLICATION PROTEIN E1

Chain E: 68% 20% 11%



• Molecule 1: REPLICATION PROTEIN E1

Chain F: 64% 22% 12%



4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	Not provided	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	FRAMES	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	20	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	62000	Depositor
Image detector	GATAN ULTRASCAN 4000 (4K X 4K)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >2	RMSZ	# Z >2
1	A	0.48	0/2240	0.59	0/3037
1	B	0.50	1/2206 (0.0%)	0.62	2/2990 (0.1%)
1	C	0.49	0/2199	0.61	0/2980
1	D	0.47	0/2248	0.64	4/3047 (0.1%)
1	E	0.48	0/2227	0.59	0/3017
1	F	0.49	0/2215	0.62	0/3001
All	All	0.48	1/13335 (0.0%)	0.61	6/18072 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	327	GLU	CD-OE1	5.96	1.32	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	486	ARG	NE-CZ-NH2	-7.44	116.58	120.30
1	B	486	ARG	NE-CZ-NH1	6.21	123.40	120.30
1	D	531	ARG	NE-CZ-NH2	-6.18	117.21	120.30
1	D	570	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	D	531	ARG	NE-CZ-NH1	5.74	123.17	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2183	0	2132	40	0
1	B	2149	0	2110	28	0
1	C	2142	0	2103	28	0
1	D	2191	0	2140	32	0
1	E	2170	0	2132	43	0
1	F	2158	0	2124	35	0
2	A	10	0	0	2	0
2	B	10	0	0	1	0
2	C	10	0	0	0	0
2	D	5	0	0	1	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
4	C	3	0	0	0	0
4	D	4	0	0	0	0
4	E	2	0	0	0	0
4	F	3	0	0	0	0
All	All	13059	0	12741	183	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 183 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:304:GLU:HG2	1:D:305:SER:H	1.18	1.06
1:E:453:SER:HB2	1:F:512:GLN:HE22	1.32	0.94
1:A:441:MET:HE1	1:A:559:ILE:H	1.32	0.92
1:A:304:GLU:HG2	1:A:305:SER:H	1.41	0.83
2:A:1580:PO4:O1	2:A:1581:PO4:O2	1.97	0.82

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/305 (87%)	244 (92%)	20 (8%)	2 (1%)	24	69
1	B	262/305 (86%)	251 (96%)	10 (4%)	1 (0%)	39	80
1	C	261/305 (86%)	252 (97%)	9 (3%)	0	100	100
1	D	268/305 (88%)	256 (96%)	8 (3%)	4 (2%)	13	57
1	E	264/305 (87%)	246 (93%)	18 (7%)	0	100	100
1	F	263/305 (86%)	239 (91%)	17 (6%)	7 (3%)	6	45
All	All	1584/1830 (87%)	1488 (94%)	82 (5%)	14 (1%)	26	67

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	506	LYS
1	D	553	GLY
1	F	479	ASP
1	F	506	LYS
1	B	576	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/259 (88%)	210 (92%)	19 (8%)	14	49
1	B	225/259 (87%)	208 (92%)	17 (8%)	16	53
1	C	224/259 (86%)	207 (92%)	17 (8%)	16	53

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	229/259 (88%)	215 (94%)	14 (6%)	23	60
1	E	228/259 (88%)	212 (93%)	16 (7%)	19	56
1	F	226/259 (87%)	200 (88%)	26 (12%)	7	32
All	All	1361/1554 (88%)	1252 (92%)	109 (8%)	20	50

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	543	ARG
1	D	527	GLN
1	F	508	LYS
1	C	548	CYS
1	D	387	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	459	ASN
1	D	507	HIS
1	F	427	ASN
1	B	527	GLN
1	C	427	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	1580	3	4,4,4	0.76	0	6,6,6	0.25	0
2	PO4	A	1581	3	4,4,4	1.06	0	6,6,6	0.25	0
2	PO4	B	1579	3	4,4,4	0.84	0	6,6,6	0.28	0
2	PO4	B	1580	3	4,4,4	0.92	0	6,6,6	0.27	0
2	PO4	C	1579	3	4,4,4	1.59	1 (25%)	6,6,6	0.44	0
2	PO4	C	1580	3	4,4,4	1.52	1 (25%)	6,6,6	0.49	0
2	PO4	D	1580	-	4,4,4	0.68	0	6,6,6	0.24	0
2	PO4	E	1579	-	4,4,4	0.54	0	6,6,6	0.23	0
2	PO4	F	1580	-	4,4,4	0.71	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	1580	3	-	0/0/0/0	0/0/0/0
2	PO4	A	1581	3	-	0/0/0/0	0/0/0/0
2	PO4	B	1579	3	-	0/0/0/0	0/0/0/0
2	PO4	B	1580	3	-	0/0/0/0	0/0/0/0
2	PO4	C	1579	3	-	0/0/0/0	0/0/0/0
2	PO4	C	1580	3	-	0/0/0/0	0/0/0/0
2	PO4	D	1580	-	-	0/0/0/0	0/0/0/0
2	PO4	E	1579	-	-	0/0/0/0	0/0/0/0
2	PO4	F	1580	-	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1579	PO4	P-O3	-2.53	1.45	1.53
2	C	1580	PO4	P-O2	-2.51	1.46	1.53

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1580	PO4	2	0
2	A	1581	PO4	1	0
2	B	1579	PO4	1	0
2	B	1580	PO4	1	0
2	D	1580	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.