



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:25 AM GMT

PDB ID : 2ADV
Title : Crystal Structures Of Glutaryl 7-Aminocephalosporanic Acid Acylase: mutational study of activation mechanism
Authors : Kim, J.K.; Yang, I.S.; Shin, H.J.; Cho, K.J.; Ryu, E.K.; Kim, S.H.; Park, S.S.; Kim, K.H.
Deposited on : 2005-07-21
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.




Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1764 (2.26-2.22)
Ramachandran outliers	100387	1724 (2.26-2.22)
Sidechain outliers	100360	1724 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	166	 90% 7% ..
2	B	28	 93% 7%
3	C	500	 88% 10% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5746 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaryl 7- Aminocephalosporanic Acid Acylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	161	Total	C	N	O	S	0	0	0
			1255	798	220	236	1			

- Molecule 2 is a protein called Glutaryl 7- Aminocephalosporanic Acid Acylase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	28	Total	C	N	O	0	0	0
			208	131	38	39			

- Molecule 3 is a protein called Glutaryl 7- Aminocephalosporanic Acid Acylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	494	Total	C	N	O	S	0	0	0
			3909	2472	689	737	11			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	103	Total	O	0	0
			103	103		
4	B	11	Total	O	0	0
			11	11		
4	C	260	Total	O	0	0
			260	260		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Glutaryl 7- Aminocephalosporanic Acid Acylase

Chain A:  90% 7% ..





• Molecule 2: Glutaryl 7- Aminocephalosporanic Acid Acylase

Chain B:  93% 7%



• Molecule 3: Glutaryl 7- Aminocephalosporanic Acid Acylase

Chain C:  88% 10% ..



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.78 Å 73.78 Å 384.65 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.22 – 2.24	Depositor
% Data completeness (in resolution range)	94.6 (48.22-2.24)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.183 , 0.210	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5746	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/1297	0.83	5/1777 (0.3%)
2	B	0.49	0/214	0.67	0/295
3	C	0.55	0/4016	0.82	16/5478 (0.3%)
All	All	0.56	0/5527	0.82	21/7550 (0.3%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121	ASP	CB-CG-OD2	8.49	125.94	118.30
3	C	420	ASP	CB-CG-OD2	7.98	125.48	118.30
1	A	143	ARG	NE-CZ-NH2	-7.71	116.45	120.30
3	C	287	ASP	CB-CG-OD2	7.13	124.72	118.30
3	C	203	ASP	CB-CG-OD2	6.77	124.39	118.30
3	C	133	ASP	CB-CG-OD2	6.51	124.16	118.30
3	C	497	ASP	CB-CG-OD2	6.38	124.04	118.30
3	C	214	ASP	CB-CG-OD2	6.32	123.99	118.30
3	C	164	ASP	CB-CG-OD2	6.17	123.85	118.30
1	A	23	ASP	CB-CG-OD2	6.03	123.73	118.30
3	C	110	ASP	CB-CG-OD2	5.96	123.67	118.30
1	A	143	ARG	NE-CZ-NH1	5.94	123.27	120.30
3	C	227	ASP	CB-CG-OD2	5.76	123.48	118.30
3	C	444	ASP	CB-CG-OD2	5.62	123.36	118.30
1	A	52	ASP	CB-CG-OD2	5.55	123.30	118.30
3	C	442	ASP	CB-CG-OD2	5.47	123.22	118.30
3	C	42	ASP	CB-CG-OD2	5.46	123.21	118.30
3	C	390	ASP	CB-CG-OD2	5.38	123.14	118.30
3	C	306	ASP	CB-CG-OD2	5.28	123.05	118.30
3	C	373	ASP	CB-CG-OD2	5.23	123.00	118.30
3	C	311	ASP	CB-CG-OD2	5.10	122.89	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1255	0	1167	2	0
2	B	208	0	203	1	0
3	C	3909	0	3754	16	0
4	A	103	0	0	0	0
4	B	11	0	0	0	0
4	C	260	0	0	4	0
All	All	5746	0	5124	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:31:PHE:CD2	4:C:783:HOH:O	2.48	0.65
3:C:50:GLN:HE22	3:C:57:ARG:HH11	1.56	0.51
3:C:300:HIS:HE1	4:C:762:HOH:O	1.94	0.51
2:B:27:THR:O	2:B:27:THR:HG23	2.10	0.51
3:C:31:PHE:HD2	4:C:783:HOH:O	1.88	0.50
1:A:35:ALA:HB3	1:A:36:PRO:HD3	1.95	0.48
3:C:190:TYR:OH	3:C:247:PRO:HB3	2.14	0.48
3:C:29:ASP:OD1	3:C:29:ASP:N	2.48	0.47
3:C:197:PRO:HB3	3:C:220:TRP:CD2	2.49	0.47
3:C:229:LEU:O	3:C:231:ARG:HG3	2.16	0.45
3:C:300:HIS:HD2	4:C:550:HOH:O	1.98	0.45
3:C:378:THR:HB	3:C:379:PRO:HA	1.98	0.45
1:A:56:ARG:HA	1:A:69:TRP:CH2	2.53	0.43
3:C:426:ALA:O	3:C:427:ALA:HB3	2.19	0.42
3:C:485:HIS:HD2	3:C:488:ASP:OD2	2.03	0.41
3:C:245:ASP:N	3:C:245:ASP:OD1	2.53	0.41
3:C:54:PRO:HD2	3:C:150:LEU:HD22	2.03	0.40
3:C:309:LEU:N	3:C:310:PRO:CD	2.83	0.40
3:C:95:ARG:NH2	3:C:124:VAL:HG11	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/166 (96%)	150 (94%)	7 (4%)	2 (1%)	15	9
2	B	26/28 (93%)	25 (96%)	1 (4%)	0	100	100
3	C	492/500 (98%)	479 (97%)	12 (2%)	1 (0%)	52	60
All	All	677/694 (98%)	654 (97%)	20 (3%)	3 (0%)	39	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	163	PRO
3	C	177	PHE
1	A	7	GLN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/128 (96%)	120 (98%)	3 (2%)	57	67
2	B	22/22 (100%)	21 (96%)	1 (4%)	34	37
3	C	409/415 (99%)	395 (97%)	14 (3%)	44	53
All	All	554/565 (98%)	536 (97%)	18 (3%)	46	56

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	116	GLN
1	A	143	ARG
2	B	21	ASN
3	C	29	ASP
3	C	30	TYR
3	C	50	GLN
3	C	83	LEU
3	C	100	ARG
3	C	105	ARG
3	C	122	SER
3	C	199	ARG
3	C	227	ASP
3	C	286	ASP
3	C	323	GLU
3	C	460	MET
3	C	475	TYR
3	C	479	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
3	C	50	GLN
3	C	300	HIS
3	C	480	GLN
3	C	485	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.