



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:44 PM GMT

PDB ID : 4AD6
Title : Synthesis and SAR of guanine based analogues for HPPK inhibitors
Authors : Chhabra, S.; Peat, T.S.; Swarbrick, J.
Deposited on : 2011-12-22
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	<div> <div></div> <div>69% 29% ...</div> </div>
1	B	161	<div> <div>6%</div> <div>80% 18% ...</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2804 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 7,8-DIHYDRO-6-HYDROXYMETHYLPTERIN-PYROPHOSPHOKINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	160	Total	C	N	O	S	0	3	0
			1306	828	224	247	7			
1	B	160	Total	C	N	O	S	0	3	0
			1301	824	222	247	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP C8MLE4
A	-1	SER	-	EXPRESSION TAG	UNP C8MLE4
A	0	HIS	-	EXPRESSION TAG	UNP C8MLE4
B	-2	GLY	-	EXPRESSION TAG	UNP C8MLE4
B	-1	SER	-	EXPRESSION TAG	UNP C8MLE4
B	0	HIS	-	EXPRESSION TAG	UNP C8MLE4

- Molecule 2 is 7-(2-HYDROXYETHYL)-8-MERCAPTOGUANINE (three-letter code: GSY) (formula: C₇H₉N₅O₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			15	7	5	2	1		
2	B	1	Total	C	N	O	S	0	0
			15	7	5	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	89	Total	O	0	0
			89	89		
3	B	78	Total	O	0	0
			78	78		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 7,8-DIHYDRO-6-HYDROXYMETHYLPTERIN-PYROPHOSPHOKINASE



- Molecule 1: 7,8-DIHYDRO-6-HYDROXYMETHYLPTERIN-PYROPHOSPHOKINASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	36.59Å 75.71Å 51.36Å 90.00° 99.71° 90.00°	Depositor
Resolution (Å)	30.00 – 1.85 29.50 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.0 (30.00-1.85) 98.0 (29.50-1.85)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.208 , 0.264 0.208 , 0.262	Depositor DCC
R_{free} test set	1194 reflections (5.44%)	DCC
Wilson B-factor (Å ²)	13.8	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	2 of 23143 reflections (0.009%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2804	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.10 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.3647e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	1/1328 (0.1%)	1.04	7/1799 (0.4%)
1	B	0.90	2/1326 (0.2%)	0.99	0/1797
All	All	0.91	3/2654 (0.1%)	1.01	7/3596 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	101	TYR	CE1-CZ	6.19	1.46	1.38
1	A	89	TRP	CD2-CE2	5.45	1.47	1.41
1	B	89	TRP	CD2-CE2	5.06	1.47	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	ARG	NE-CZ-NH2	-8.78	115.91	120.30
1	A	92	ARG	NE-CZ-NH1	7.29	123.95	120.30
1	A	142	LEU	CA-CB-CG	6.98	131.35	115.30
1	A	15	ARG	NE-CZ-NH2	-6.62	116.99	120.30
1	A	152	ASP	CB-CG-OD1	5.32	123.08	118.30
1	A	117	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	A	151	ASP	CB-CG-OD1	5.09	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1306	0	1336	39	0
1	B	1301	0	1326	37	0
2	A	15	0	9	1	0
2	B	15	0	9	0	0
3	A	89	0	0	7	0
3	B	78	0	0	3	0
All	All	2804	0	2680	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43[B]:THR:HG23	1:A:154:VAL:HG22	1.43	1.00
1:B:25:ILE:HG22	1:B:29:TYR:HE2	1.30	0.97
1:B:25:ILE:HG22	1:B:29:TYR:CE2	2.05	0.92
1:A:115:HIS:HD2	1:A:117:ARG:H	1.09	0.91
1:A:43[B]:THR:HG21	1:A:123:PHE:HB3	1.54	0.89
1:A:115:HIS:CD2	1:A:117:ARG:H	1.91	0.89
1:B:2:ILE:HD11	1:B:102:GLY:N	1.89	0.87
1:B:2:ILE:HD11	1:B:102:GLY:CA	2.09	0.82
1:A:32:ILE:O	3:A:2027:HOH:O	1.98	0.82
1:A:43[B]:THR:HG22	3:A:2038:HOH:O	1.82	0.77
1:A:43[B]:THR:CG2	1:A:154:VAL:HG22	2.18	0.74
1:B:115:HIS:CD2	1:B:117:ARG:H	2.05	0.73
2:A:1159:GSY:S10	1:B:110:LYS:HG2	2.29	0.73
1:B:2:ILE:HG22	1:B:63:THR:O	1.90	0.71
1:A:115:HIS:HD2	1:A:117:ARG:N	1.89	0.68
1:A:83[B]:ARG:O	1:A:92:ARG:HD3	1.94	0.67
1:A:29:TYR:HE1	1:A:76:LYS:HZ2	1.40	0.67
1:B:2:ILE:HG23	1:B:63:THR:HG23	1.77	0.67
1:A:88:ARG:HG2	3:A:2057:HOH:O	1.95	0.65
1:B:154:VAL:O	3:B:2074:HOH:O	2.15	0.65
1:B:25:ILE:O	1:B:29:TYR:CD2	2.50	0.65
1:B:25:ILE:O	1:B:29:TYR:HD2	1.80	0.65
1:B:32:ILE:O	3:B:2022:HOH:O	2.15	0.63
1:B:2:ILE:HD11	1:B:102:GLY:HA3	1.80	0.63
1:B:2:ILE:HG23	1:B:63:THR:CG2	2.28	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2:ILE:CD1	1:B:102:GLY:CA	2.77	0.61
1:A:43[B]:THR:HG21	1:A:123:PHE:CB	2.29	0.60
1:A:81:LEU:HD12	1:A:94:LEU:HD22	1.82	0.60
1:A:83[A]:ARG:O	1:A:92:ARG:HD3	2.02	0.58
1:B:2:ILE:CG2	1:B:63:THR:O	2.53	0.57
1:B:2:ILE:CD1	1:B:102:GLY:HA3	2.35	0.56
1:B:115:HIS:NE2	1:B:117:ARG:HB2	2.20	0.56
1:A:79:GLU:HB2	1:B:75:LEU:HD11	1.87	0.56
1:A:105:MET:CG	1:A:138:PRO:HB2	2.35	0.56
1:B:2:ILE:CG2	1:B:63:THR:HG23	2.35	0.56
1:A:71:LEU:O	1:A:75:LEU:HG	2.04	0.56
1:A:6:LEU:HB2	1:A:59:VAL:HG12	1.89	0.55
1:B:25:ILE:CG2	1:B:29:TYR:CE2	2.87	0.55
1:B:54:PHE:CE2	3:B:2009:HOH:O	2.53	0.54
1:B:26:LEU:HD23	1:B:73[B]:CYS:SG	2.48	0.54
1:A:37:ILE:HG22	1:A:38:SER:O	2.09	0.52
1:B:115:HIS:ND1	1:B:116:PRO:HD2	2.26	0.51
1:B:25:ILE:CG2	1:B:29:TYR:HE2	2.15	0.50
1:B:115:HIS:CD2	1:B:117:ARG:HB2	2.47	0.50
1:A:75:LEU:HD23	3:A:2051:HOH:O	2.11	0.50
1:A:85:ARG:HG3	1:A:92:ARG:HD2	1.94	0.50
1:A:85:ARG:HG2	3:A:2055:HOH:O	2.12	0.48
1:B:88:ARG:HG3	1:B:89:TRP:H	1.79	0.48
1:A:71:LEU:HG	1:A:75:LEU:HD11	1.95	0.47
1:A:114:PRO:HG2	1:A:137:GLU:OE2	2.14	0.47
1:A:23:ILE:HD13	1:A:37:ILE:HD13	1.97	0.47
1:A:72:GLU:HG3	1:A:76:LYS:HE2	1.97	0.46
1:B:86:LYS:HB2	1:B:89:TRP:HB2	1.96	0.46
1:B:57:LEU:HD12	1:B:57:LEU:C	2.36	0.46
1:B:86:LYS:HD2	1:B:89:TRP:CE2	2.51	0.46
1:B:66:THR:OG1	1:B:69:GLN:HG3	2.16	0.45
1:A:100:LEU:HD23	1:A:114:PRO:HG3	1.97	0.45
1:A:81:LEU:CD1	1:A:94:LEU:HD22	2.45	0.45
1:B:115:HIS:CD2	1:B:117:ARG:N	2.82	0.45
1:A:2:ILE:HD13	1:A:101:TYR:OH	2.17	0.44
1:A:11:ASN:O	1:A:52:PRO:HB2	2.18	0.44
1:A:68:LEU:HD21	1:A:108:LEU:HD12	1.99	0.44
1:A:146:ASP:OD2	3:A:2084:HOH:O	2.21	0.43
1:B:1:MET:SD	1:B:64:THR:HG23	2.57	0.43
1:B:68:LEU:HD22	1:B:110:LYS:HE3	2.00	0.43
1:A:43[A]:THR:HG22	1:A:154:VAL:HG22	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:ASN:HA	1:A:20:ASN:HD22	1.70	0.42
1:A:5:TYR:CE2	1:A:132:ALA:HB2	2.55	0.41
1:A:142:LEU:HB3	1:A:147:LEU:HG	2.03	0.41
1:B:100:LEU:HD23	1:B:114:PRO:HG3	2.02	0.41
1:A:126:ILE:N	1:A:127:PRO:HD2	2.35	0.41
1:B:8:LEU:N	1:B:8:LEU:HD12	2.36	0.41
1:A:100:LEU:HD22	3:A:2070:HOH:O	2.19	0.41
1:A:57:LEU:HD12	1:A:57:LEU:C	2.41	0.41
1:B:7:GLY:C	1:B:8:LEU:HD12	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/161 (100%)	159 (99%)	2 (1%)	0	100	100
1	B	161/161 (100%)	157 (98%)	3 (2%)	1 (1%)	30	13
All	All	322/322 (100%)	316 (98%)	5 (2%)	1 (0%)	46	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	89	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/147 (102%)	147 (98%)	3 (2%)	63	47
1	B	150/147 (102%)	145 (97%)	5 (3%)	45	25
All	All	300/294 (102%)	292 (97%)	8 (3%)	52	34

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	ASN
1	A	142	LEU
1	A	156	ARG
1	B	2	ILE
1	B	75	LEU
1	B	92	ARG
1	B	103	GLU
1	B	156	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	20	ASN
1	A	62	GLN
1	A	115	HIS
1	B	3	GLN
1	B	62	GLN
1	B	115	HIS
1	B	134	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GSY	A	1159	-	12,16,16	2.02	3 (25%)	9,23,23	2.56	5 (55%)
2	GSY	B	1159	-	12,16,16	2.51	4 (33%)	9,23,23	2.41	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GSY	A	1159	-	-	0/3/3/3	0/2/2/2
2	GSY	B	1159	-	-	0/3/3/3	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1159	GSY	C8-N7	-6.05	1.30	1.36
2	A	1159	GSY	C8-N7	-3.64	1.32	1.36
2	B	1159	GSY	C2-N1	2.11	1.39	1.35
2	B	1159	GSY	C13-C14	2.31	1.57	1.50
2	A	1159	GSY	C6-N1	3.43	1.39	1.33
2	A	1159	GSY	C2-N1	3.98	1.42	1.35
2	B	1159	GSY	C6-N1	4.64	1.41	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1159	GSY	N3-C2-N1	-4.57	120.48	127.44
2	B	1159	GSY	C5-C6-N1	-4.19	117.87	123.59
2	A	1159	GSY	C5-C6-N1	-3.91	118.24	123.59
2	A	1159	GSY	N3-C2-N1	-3.56	122.03	127.44
2	A	1159	GSY	C14-C13-N7	-2.35	105.89	110.75
2	B	1159	GSY	C6-N1-C2	2.10	118.85	115.94
2	A	1159	GSY	N2-C2-N1	2.43	121.22	117.20
2	A	1159	GSY	C13-N7-C8	4.13	128.49	125.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1159	GSY	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	160/161 (99%)	0.28	2 (1%) 79 79	5, 11, 23, 29	4 (2%)
1	B	160/161 (99%)	0.59	9 (5%) 28 26	6, 14, 35, 48	0
All	All	320/322 (99%)	0.43	11 (3%) 49 46	5, 13, 27, 48	4 (1%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	89	TRP	7.3
1	B	87	GLU	5.5
1	B	84	ILE	5.5
1	B	85[A]	ARG	5.5
1	B	86	LYS	4.5
1	B	91	PRO	4.3
1	B	105	MET	3.2
1	A	84	ILE	2.9
1	B	116	PRO	2.9
1	B	88	ARG	2.5
1	A	-1	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GSY	B	1159	15/15	0.93	0.11	-0.46	8,9,10,11	0
2	GSY	A	1159	15/15	0.94	0.10	-1.03	9,10,14,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.