



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:20 AM GMT

PDB ID : 3AE9
Title : Crystal structure of porcine heart mitochondrial complex II bound with N-(3-Pentafluorophenoxy-phenyl)-2-trifluoromethyl-benzamide
Authors : Harada, S.; Sasaki, T.; Shindo, M.; Kido, Y.; Inaoka, D.K.; Omori, J.; Osanai, A.; Sakamoto, K.; Mao, J.; Matsuoka, S.; Inoue, M.; Honma, T.; Tanaka, A.; Kita, K.
Deposited on : 2010-02-04
Resolution : 3.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1198 (3.40-3.24)
Clashscore	102246	1280 (3.40-3.24)
Ramachandran outliers	100387	1260 (3.40-3.24)
Sidechain outliers	100360	1259 (3.40-3.24)
RSRZ outliers	91569	1203 (3.40-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	622	<div> <div>7%</div> <div>82% 16% ..</div> </div>
2	B	252	<div> <div>2%</div> <div>83% 12% 5%</div> </div>
3	C	140	<div> <div>4%</div> <div>91% 7% ..</div> </div>
4	D	103	<div> <div>4%</div> <div>97% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	FD8	C	1201	-	-	-	X
12	EPH	D	1306	-	-	-	X

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 8677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	613	Total	C	N	O	S	0	0	0
			4729	2954	848	895	32			

- Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	239	Total	C	N	O	S	0	0	0
			1922	1214	326	360	22			


- Molecule 3 is a protein called Succinate dehydrogenase cytochrome b560 subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	138	Total	C	N	O	S	0	0	0
			1064	695	179	183	7			

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	102	Total	C	N	O	S	0	0	0
			765	499	128	133	5			

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 6 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 7 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 8 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 9 is MALONATE ION (three-letter code: MLI) (formula: $\text{C}_3\text{H}_2\text{O}_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	O		0	0
			7	3	4			

- Molecule 10 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 11 is N-[3-(PENTAFLUOROPHENOXY)PHENYL]-2-(TRIFLUOROMETHYL) BENZAMIDE (three-letter code: FD8) (formula: $C_{20}H_9F_8NO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	C	1	Total	C	F	N	O	0	0
			31	20	8	1	2		

- Molecule 12 is L-ALPHA-PHOSPHATIDYL-BETA-OLEOYL-GAMMA-PALMITOYL-PHOSPHATIDYLETHANOLAMINE (three-letter code: EPH) (formula: $C_{39}H_{68}NO_8P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	D	1	Total	C	N	O	P	0	0
			44	34	1	8	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial




- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial



- Molecule 3: Succinate dehydrogenase cytochrome b560 subunit, mitochondrial



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.61Å 84.25Å 295.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.76 – 3.31 37.76 – 3.31	Depositor EDS
% Data completeness (in resolution range)	97.8 (37.76-3.31) 97.8 (37.76-3.31)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 3.32Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.257 , 0.307 0.256 , 0.305	Depositor DCC
R_{free} test set	1367 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	97.3	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 55.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 27276 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8677	wwPDB-VP
Average B, all atoms (Å ²)	124.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MLI, F3S, FES, EPH, HEM, FD8, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/4828	0.47	0/6531
2	B	0.31	0/1964	0.46	0/2648
3	C	0.32	0/1091	0.43	0/1483
4	D	0.29	0/784	0.42	0/1066
All	All	0.31	0/8667	0.46	0/11728

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4729	0	4618	57	0
2	B	1922	0	1901	15	0
3	C	1064	0	1104	8	0
4	D	765	0	773	0	0
5	A	53	0	31	8	0
6	B	4	0	0	0	0
7	B	8	0	0	0	0
8	B	7	0	0	1	0
9	A	7	0	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	C	43	0	30	2	0
11	C	31	0	9	2	0
12	D	44	0	53	0	0
All	All	8677	0	8521	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:HIS:NE2	5:A:700:FAD:HM82	1.36	1.41
1:A:57:HIS:CE1	5:A:700:FAD:HM82	2.00	0.96
1:A:61:ALA:HB3	1:A:155:GLY:HA3	1.49	0.91
1:A:49:THR:HA	5:A:700:FAD:H8A	1.59	0.83
1:A:76:ASN:HD21	1:A:78:ARG:HB2	1.55	0.71
1:A:415:LEU:HG	5:A:700:FAD:C2	2.22	0.69
1:A:59:VAL:HG13	1:A:159:LEU:HD23	1.78	0.66
1:A:518:THR:HA	1:A:521:VAL:HG22	1.81	0.62
1:A:113:ASN:HD22	2:B:138:GLY:H	1.47	0.61
1:A:57:HIS:NE2	5:A:700:FAD:HM81	2.10	0.61
1:A:566:LYS:HB2	1:A:567:PRO:HD2	1.83	0.59
1:A:258:ILE:HD11	1:A:265:ILE:HG23	1.85	0.59
11:C:1201:FD8:O1	11:C:1201:FD8:H14	2.03	0.58
2:B:51:ILE:HD11	2:B:59:LEU:HD22	1.85	0.57
1:A:111:LEU:HD11	1:A:419:VAL:HG21	1.86	0.57
1:A:61:ALA:HA	5:A:700:FAD:N5	2.21	0.56
1:A:60:ALA:HB1	5:A:700:FAD:HM72	1.87	0.55
2:B:102:LEU:HD22	2:B:166:THR:HG21	1.89	0.54
1:A:56:SER:O	1:A:59:VAL:HG22	2.08	0.54
1:A:182:LEU:HD13	1:A:237:MET:HB3	1.90	0.54
1:A:134:GLN:HG3	1:A:410:LEU:HD21	1.89	0.54
3:C:52:LEU:HB3	10:C:1305:HEM:HAC	1.89	0.53
1:A:415:LEU:HA	1:A:418:LEU:HD22	1.93	0.50
1:A:67:ALA:HB2	1:A:108:VAL:HG21	1.93	0.50
1:A:52:PHE:HB3	1:A:55:ARG:HG2	1.94	0.49
1:A:219:ARG:HD2	1:A:228:HIS:HA	1.93	0.49
1:A:516:TRP:HB3	2:B:60:THR:HG21	1.94	0.48
2:B:181:PRO:HG3	8:B:304:F3S:S3	2.53	0.48
2:B:231:PRO:O	2:B:235:ILE:HD12	2.14	0.48
1:A:59:VAL:HA	1:A:156:HIS:HA	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:188:TYR:O	2:B:192:ILE:HD13	2.13	0.48
1:A:233:ASP:HB2	5:A:700:FAD:N1A	2.28	0.48
1:A:504:LEU:HD11	1:A:530:MET:HG3	1.96	0.48
1:A:322:VAL:HG12	1:A:323:TYR:H	1.79	0.47
1:A:366:TYR:HE1	1:A:414:SER:HG	1.62	0.47
1:A:116:MET:HA	1:A:161:THR:HG21	1.97	0.47
2:B:79:GLY:HA3	3:C:34:ARG:HB2	1.96	0.47
1:A:579:HIS:HD2	1:A:581:ARG:H	1.63	0.46
1:A:64:GLY:HA2	1:A:154:THR:HG21	1.98	0.46
1:A:100:MET:HA	1:A:420:VAL:HG11	1.97	0.46
1:A:152:ASP:HB2	1:A:339:PRO:HD2	1.99	0.45
1:A:118:PHE:HA	1:A:150:VAL:HG22	1.99	0.45
1:A:215:GLY:HA2	1:A:398:GLU:HB3	1.98	0.45
1:A:150:VAL:O	1:A:150:VAL:HG12	2.16	0.45
2:B:181:PRO:HA	2:B:235:ILE:HD11	1.98	0.45
1:A:568:ILE:HG13	1:A:569:GLN:N	2.32	0.45
1:A:299:ASP:O	1:A:303:ARG:HB2	2.17	0.45
3:C:52:LEU:HD21	3:C:98:LEU:HA	1.99	0.45
1:A:10:SER:C	1:A:12:GLN:H	2.20	0.45
1:A:322:VAL:HG12	1:A:323:TYR:N	2.32	0.44
2:B:219:MET:CE	2:B:232:GLY:HA3	2.48	0.44
1:A:150:VAL:HB	1:A:154:THR:HA	1.98	0.44
1:A:163:TYR:O	1:A:167:LEU:HD22	2.18	0.44
3:C:91:LYS:HE3	3:C:141:ALA:O	2.18	0.44
1:A:221:TYR:CG	1:A:364:VAL:HG21	2.53	0.44
1:A:463:THR:HG23	1:A:464:ILE:HG13	1.99	0.43
1:A:289:ALA:HA	1:A:290:PRO:HD3	1.86	0.43
1:A:104:ALA:HA	1:A:416:LEU:HD11	1.99	0.43
1:A:104:ALA:HB3	1:A:105:PRO:HD3	2.00	0.43
1:A:258:ILE:HG22	1:A:260:GLY:H	1.83	0.42
3:C:102:THR:HG23	10:C:1305:HEM:HAB	2.02	0.42
1:A:254:HIS:HA	1:A:255:PRO:HD3	1.84	0.42
1:A:25:VAL:HG21	1:A:192:VAL:HG11	2.01	0.42
3:C:119:ILE:N	3:C:120:PRO:HD2	2.35	0.42
1:A:199:ASP:HA	1:A:515:VAL:HG22	2.02	0.42
1:A:342:SER:HA	1:A:353:VAL:HG11	2.02	0.42
2:B:139:LYS:HG3	2:B:140:GLN:HG2	2.02	0.41
11:C:1201:FD8:O1	11:C:1201:FD8:C14	2.69	0.41
2:B:16:ILE:HG12	2:B:99:ILE:HB	2.03	0.41
1:A:81:PHE:HA	1:A:101:THR:HG21	2.03	0.41
1:A:521:VAL:HA	1:A:524:LEU:HD12	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:233:LYS:HA	3:C:117:LEU:HD13	2.02	0.41
2:B:116:SER:HB2	3:C:15:PHE:HD1	1.86	0.41
2:B:62:ARG:HH12	2:B:115:LEU:HD12	1.86	0.41
1:A:461:ASN:HA	1:A:508:LYS:HD2	2.03	0.40
1:A:57:HIS:O	1:A:57:HIS:CG	2.75	0.40
1:A:72:MET:HG2	1:A:128:GLN:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	611/622 (98%)	571 (94%)	38 (6%)	2 (0%)	46	81
2	B	237/252 (94%)	222 (94%)	15 (6%)	0	100	100
3	C	136/140 (97%)	129 (95%)	7 (5%)	0	100	100
4	D	100/103 (97%)	97 (97%)	3 (3%)	0	100	100
All	All	1084/1117 (97%)	1019 (94%)	63 (6%)	2 (0%)	52	85

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	290	PRO
1	A	260	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	499/506 (99%)	482 (97%)	17 (3%)	44	77
2	B	214/220 (97%)	203 (95%)	11 (5%)	29	67
3	C	117/118 (99%)	115 (98%)	2 (2%)	68	86
4	D	76/76 (100%)	74 (97%)	2 (3%)	54	82
All	All	906/920 (98%)	874 (96%)	32 (4%)	43	76

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	THR
1	A	65	ILE
1	A	74	GLU
1	A	121	THR
1	A	129	ARG
1	A	131	PHE
1	A	157	SER
1	A	165	ARG
1	A	167	LEU
1	A	313	ARG
1	A	368	MET
1	A	418	LEU
1	A	458	ARG
1	A	507	LEU
1	A	556	LYS
1	A	561	GLU
1	A	597	LEU
2	B	10	ARG
2	B	51	ILE
2	B	63	ARG
2	B	66	ARG
2	B	87	ARG
2	B	159	ILE
2	B	189	ARG
2	B	192	ILE
2	B	214	ARG
2	B	217	THR
2	B	230	ASN
3	C	52	LEU
3	C	122	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	78	LEU
4	D	108	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	76	ASN
1	A	113	ASN
1	A	128	GLN
1	A	246	GLN
1	A	408	ASN
1	A	461	ASN
1	A	474	GLN
1	A	571	GLN
1	A	579	HIS
2	B	121	GLN
2	B	174	ASN
2	B	230	ASN
3	C	17	ASN
3	C	104	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	FAD	A	700	1	48,58,58	1.24	6 (12%)	54,89,89	2.19	8 (14%)
9	MLI	A	701	-	0,6,6	0.00	-	0,7,7	0.00	-
6	FES	B	302	2	0,4,4	0.00	-	0,4,4	0.00	-
7	SF4	B	303	2	0,12,12	0.00	-	0,24,24	0.00	-
8	F3S	B	304	2	0,9,9	0.00	-	0,15,15	0.00	-
11	FD8	C	1201	-	33,33,33	3.62	7 (21%)	48,49,49	0.73	0
10	HEM	C	1305	3,4	30,50,50	2.17	9 (30%)	24,82,82	2.35	10 (41%)
12	EPH	D	1306	-	42,43,48	1.50	6 (14%)	43,48,53	1.45	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FAD	A	700	1	-	0/30/50/50	0/6/6/6
9	MLI	A	701	-	-	0/0/4/4	0/0/0/0
6	FES	B	302	2	-	0/0/4/4	0/1/1/1
7	SF4	B	303	2	-	0/0/48/48	0/6/5/5
8	F3S	B	304	2	-	0/0/24/24	0/0/3/3
11	FD8	C	1201	-	-	0/18/18/18	0/3/3/3
10	HEM	C	1305	3,4	-	0/10/54/54	0/0/8/8
12	EPH	D	1306	-	-	0/47/47/52	0/0/0/0

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	C	1305	HEM	C3B-C4B	-7.17	1.45	1.51
10	C	1305	HEM	C3D-C4D	-4.77	1.45	1.51
10	C	1305	HEM	C2C-C1C	-3.61	1.45	1.52
11	C	1201	FD8	C9-N	-2.79	1.36	1.41
10	C	1305	HEM	C2D-C1D	-2.02	1.45	1.51
10	C	1305	HEM	C3C-CAC	2.03	1.55	1.51
10	C	1305	HEM	C4C-NC	2.08	1.38	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	700	FAD	C5X-N5	2.10	1.38	1.35
10	C	1305	HEM	FE-NC	2.25	2.04	1.95
10	C	1305	HEM	C1C-NC	2.34	1.38	1.36
5	A	700	FAD	C2A-N1A	2.55	1.38	1.33
10	C	1305	HEM	FE-ND	2.63	2.11	1.97
5	A	700	FAD	C1'-N10	2.65	1.51	1.48
5	A	700	FAD	C4-N3	2.99	1.38	1.33
12	D	1306	EPH	C15-C16	3.15	1.52	1.29
5	A	700	FAD	C4X-N5	3.46	1.38	1.33
12	D	1306	EPH	C13-C12	3.67	1.52	1.31
12	D	1306	EPH	C25-C24	3.68	1.52	1.31
12	D	1306	EPH	C29-C28	3.68	1.52	1.31
5	A	700	FAD	C2A-N3A	3.75	1.38	1.32
12	D	1306	EPH	O2-C4	4.21	1.46	1.33
12	D	1306	EPH	O1-C3	4.23	1.47	1.34
11	C	1201	FD8	C15-C16	7.42	1.53	1.39
11	C	1201	FD8	C15-C20	7.58	1.53	1.39
11	C	1201	FD8	C19-C18	8.55	1.53	1.37
11	C	1201	FD8	C20-C19	8.55	1.53	1.37
11	C	1201	FD8	C18-C17	8.57	1.53	1.37
11	C	1201	FD8	C17-C16	8.60	1.53	1.37

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	700	FAD	N3A-C2A-N1A	-12.10	119.63	128.89
5	A	700	FAD	P-O3P-PA	-3.63	122.55	132.73
12	D	1306	EPH	C30-C29-C28	-3.54	110.89	127.06
10	C	1305	HEM	C3B-CAB-CBB	-3.35	119.31	124.46
10	C	1305	HEM	C3C-CAC-CBC	-2.92	119.98	124.46
5	A	700	FAD	C4X-C4-N3	-2.49	120.18	123.59
5	A	700	FAD	C4B-O4B-C1B	-2.26	107.24	109.72
12	D	1306	EPH	C14-C15-C16	-2.25	111.10	125.00
12	D	1306	EPH	C14-C13-C12	-2.22	111.25	125.00
12	D	1306	EPH	C11-C12-C13	-2.06	110.98	125.34
10	C	1305	HEM	C3B-C4B-NB	-2.06	107.69	111.63
12	D	1306	EPH	C26-C25-C24	-2.06	111.00	125.34
12	D	1306	EPH	C23-C24-C25	-2.02	111.31	125.34
10	C	1305	HEM	C2D-C3D-C4D	2.07	105.00	101.50
10	C	1305	HEM	C3B-C4B-CHC	2.50	126.69	123.16
12	D	1306	EPH	O2-C4-C18	2.60	119.82	111.90
10	C	1305	HEM	CMD-C2D-C3D	2.82	126.83	114.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	700	FAD	C5X-C9A-N10	2.94	119.86	117.62
5	A	700	FAD	O4B-C1B-N9A	3.21	114.81	108.10
5	A	700	FAD	C4X-N5-C5X	3.34	120.60	116.76
10	C	1305	HEM	CMC-C2C-C3C	3.83	126.09	116.53
10	C	1305	HEM	CMB-C2B-C3B	3.86	126.16	116.53
12	D	1306	EPH	O1-C3-C5	4.36	120.99	111.53
10	C	1305	HEM	CAD-C3D-C2D	4.55	126.30	113.22
10	C	1305	HEM	CAD-C3D-C4D	4.60	128.70	112.47
5	A	700	FAD	C4-N3-C2	5.46	119.97	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	700	FAD	8	0
8	B	304	F3S	1	0
11	C	1201	FD8	2	0
10	C	1305	HEM	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	613/622 (98%)	0.37	43 (7%) 19 19	89, 120, 184, 205	0
2	B	239/252 (94%)	0.29	6 (2%) 61 60	91, 113, 152, 184	0
3	C	138/140 (98%)	0.13	6 (4%) 39 37	94, 117, 183, 207	0
4	D	102/103 (99%)	0.06	4 (3%) 43 41	98, 120, 160, 176	0
All	All	1092/1117 (97%)	0.29	59 (5%) 29 28	89, 118, 178, 207	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	35	SER	8.8
1	A	327	HIS	5.0
1	A	326	LEU	4.7
3	C	81	CYS	4.6
1	A	357	PRO	4.6
1	A	277	ILE	4.5
1	A	322	VAL	4.3
1	A	356	GLU	4.2
1	A	321	HIS	4.0
3	C	143	MET	4.0
1	A	323	TYR	3.8
2	B	11	ILE	3.8
1	A	317	PRO	3.8
1	A	325	GLN	3.6
1	A	280	GLN	3.5
2	B	34	GLU	3.4
1	A	288	TYR	3.4
3	C	84	PRO	3.3
1	A	622	TYR	3.2
1	A	10	SER	3.1
1	A	279	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	21	GLY	3.1
1	A	353	VAL	3.1
2	B	70	CYS	3.0
1	A	225	THR	2.6
1	A	328	HIS	2.6
1	A	315	CYS	2.6
1	A	337	ARG	2.6
1	A	354	THR	2.6
3	C	80	LEU	2.6
1	A	367	ASN	2.5
1	A	276	LEU	2.5
1	A	230	SER	2.5
1	A	576	PHE	2.4
2	B	245	TYR	2.4
1	A	57	HIS	2.4
1	A	358	ILE	2.3
1	A	216	GLY	2.3
1	A	398	GLU	2.3
4	D	36	SER	2.3
1	A	332	GLU	2.3
3	C	20	LEU	2.3
1	A	359	PRO	2.2
1	A	215	GLY	2.2
1	A	11	THR	2.2
1	A	352	ASP	2.2
1	A	155	GLY	2.2
4	D	96	ALA	2.1
1	A	226	SER	2.1
1	A	365	HIS	2.1
1	A	320	ASP	2.1
1	A	329	LEU	2.1
1	A	214	THR	2.1
2	B	246	LYS	2.1
2	B	71	GLY	2.0
1	A	227	ALA	2.0
4	D	136	LEU	2.0
1	A	305	MET	2.0
1	A	369	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
12	EPH	D	1306	44/49	0.82	0.67	3.79	150,153,156,156	0
11	FD8	C	1201	31/31	0.85	0.34	3.53	136,137,138,138	0
10	HEM	C	1305	43/43	0.96	0.26	1.70	101,102,103,104	0
5	FAD	A	700	53/53	0.90	0.40	0.92	98,100,101,101	0
9	MLI	A	701	7/7	0.75	0.31	-0.06	222,222,222,222	0
8	F3S	B	304	7/7	0.99	0.21	-0.48	99,99,99,99	0
7	SF4	B	303	8/8	1.00	0.20	-1.00	85,85,86,86	0
6	FES	B	302	4/4	0.97	0.22	-1.06	100,100,100,101	0

6.5 Other polymers [i](#)

There are no such residues in this entry.