



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:27 AM GMT

PDB ID : 2AGL  
Title : Crystal structure of the phenylhydrazine adduct of aromatic amine dehydrogenase from *Alcaligenes faecalis*  
Authors : Masgrau, L.; Roujeinikova, A.; Johannissen, L.O.; Hothi, P.; Basran, J.; Ranaghan, K.E.; Mulholland, A.J.; Sutcliffe, M.J.; Scrutton, N.S.; Leys, D.  
Deposited on : 2005-07-27  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865

**i**





## X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



<b>Metric</b>	<b>Whole archive (#Entries)</b>	<b>Similar resolution (#Entries, resolution range(Å))</b>
R <sub>free</sub>	91344	1199 (1.40-1.40)
Clashscore	102246	1295 (1.40-1.40)
Ramachandran outliers	100387	1259 (1.40-1.40)
Sidechain outliers	100360	1258 (1.40-1.40)
RSRZ outliers	91569	1198 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	135	
1	H	135	
2	A	361	
2	B	361	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PHZ	D	1	-	-	-	X
3	PHZ	H	1	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8464 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aromatic amine dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	109	Total	C	N	O	S	0	1	0
			842	514	148	165	15			
1	H	119	Total	C	N	O	S	0	4	0
			933	572	163	183	15			

- Molecule 2 is a protein called Aromatic amine dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	360	Total	C	N	O	S	0	2	0
			2811	1773	491	533	14			
2	B	360	Total	C	N	O	S	0	2	0
			2788	1758	484	532	14			

- Molecule 3 is 1-PHENYLHYDRAZINE (three-letter code: PHZ) (formula: C<sub>6</sub>H<sub>8</sub>N<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	C	N	0	0
			8	6	2		
3	D	1	Total	C	N	0	0
			8	6	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	464	Total	O	0	0
			464	464		
4	B	393	Total	O	0	0
			393	393		
4	D	117	Total	O	0	0
			117	117		
4	H	99	Total	O	0	1
			100	100		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aromatic amine dehydrogenase

Chain D: 




- Molecule 1: Aromatic amine dehydrogenase

Chain H: 



- Molecule 2: Aromatic amine dehydrogenase

Chain A: 



- Molecule 2: Aromatic amine dehydrogenase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.69Å 88.67Å 80.52Å 90.00° 90.24° 90.00°	Depositor
Resolution (Å)	15.00 – 1.40 80.52 – 1.40	Depositor EDS
% Data completeness (in resolution range)	96.2 (15.00-1.40) 96.2 (80.52-1.40)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.49 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.1.9999	Depositor
R, $R_{free}$	0.170 , 0.196 0.172 , 0.197	Depositor DCC
$R_{free}$ test set	9440 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	12.2	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.7	EDS
Estimated twinning fraction	0.014 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	2 of 194937 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8464	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRQ, PHZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	D	0.56	0/849	0.82	0/1156
1	H	0.57	0/942	0.81	3/1285 (0.2%)
2	A	0.59	0/2883	0.81	2/3907 (0.1%)
2	B	0.57	0/2861	0.82	4/3881 (0.1%)
All	All	0.57	0/7535	0.81	9/10229 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	147	ARG	NE-CZ-NH1	7.05	123.83	120.30
2	A	316	ASP	CB-CG-OD2	6.17	123.86	118.30
2	B	378	ASP	CB-CG-OD2	5.90	123.61	118.30
2	A	308	ASP	CB-CG-OD2	5.57	123.31	118.30
2	B	299	ASP	CB-CG-OD2	5.41	123.17	118.30
2	B	347	ASP	CB-CG-OD2	5.32	123.09	118.30
1	H	147	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	H	118	ASP	CB-CG-OD2	5.20	122.98	118.30
2	B	385	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	842	0	738	10	0
1	H	933	0	816	13	0
2	A	2811	0	2752	22	0
2	B	2788	0	2700	18	0
3	D	8	0	6	0	0
3	H	8	0	6	0	0
4	A	464	0	0	3	0
4	B	393	0	0	3	0
4	D	117	0	0	3	0
4	H	100	0	0	6	0
All	All	8464	0	7018	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:118:ASP:OD2	4:H:212:HOH:O	1.92	0.86
1:D:179:LEU:C	4:D:251:HOH:O	2.20	0.79
2:A:124:ASN:HD21	2:A:178:GLY:H	1.31	0.77
2:A:104:ARG:HH11	2:A:106:HIS:HE1	1.30	0.77
1:H:105:SER:HA	1:H:164:ASN:HD21	1.50	0.76
2:B:104:ARG:HH11	2:B:106:HIS:HE1	1.34	0.74
4:H:232:HOH:O	2:B:118[B]:MET:SD	2.45	0.74
1:H:115:ASN:O	4:H:212:HOH:O	2.04	0.73
2:B:124:ASN:HD21	2:B:178:GLY:H	1.34	0.73
1:D:105:SER:HA	1:D:164:ASN:HD21	1.54	0.71
2:B:91:TYR:OH	2:B:428:HIS:HD2	1.77	0.67
2:A:91:TYR:OH	2:A:428:HIS:HD2	1.78	0.66
2:A:297:SER:OG	4:A:896:HOH:O	2.06	0.66
1:D:95:THR:HG22	4:D:247:HOH:O	1.96	0.65
2:B:197:GLN:NE2	2:B:227:SER:H	1.97	0.63
1:H:120:LYS:N	4:H:212:HOH:O	2.11	0.61
2:A:197:GLN:NE2	2:A:227:SER:H	1.99	0.61
1:D:167:SER:HB2	2:B:177:GLN:HE22	1.67	0.58
1:H:167:SER:HB2	2:A:177:GLN:HE22	1.68	0.57
2:A:91:TYR:OH	2:A:428:HIS:CD2	2.59	0.55
2:A:197:GLN:HE21	2:A:227:SER:H	1.53	0.54
2:A:335:ARG:H	2:A:386:GLN:HE22	1.53	0.54
2:B:106:HIS:HD2	2:B:418:ALA:O	1.90	0.54
2:B:197:GLN:HE21	2:B:227:SER:H	1.56	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:115:ASN:HB3	4:H:212:HOH:O	2.06	0.54
1:D:109:TRQ:HB2	1:D:160:TRP:NE1	2.22	0.53
1:D:71:GLU:N	4:D:278:HOH:O	2.42	0.53
1:H:61:SER:N	4:H:245:HOH:O	2.40	0.53
2:A:335:ARG:H	2:A:386:GLN:NE2	2.06	0.53
1:H:109:TRQ:HB2	1:H:160:TRP:NE1	2.24	0.53
2:B:335:ARG:H	2:B:386:GLN:HE22	1.56	0.52
1:H:108:SER:HB2	1:H:126[A]:TYR:O	2.10	0.52
1:D:155:HIS:HE1	4:B:451:HOH:O	1.92	0.51
2:B:335:ARG:H	2:B:386:GLN:NE2	2.10	0.49
2:A:143:HIS:HD2	4:A:693:HOH:O	1.95	0.49
1:H:159:ASN:HD22	2:A:177:GLN:NE2	2.11	0.47
2:A:106:HIS:HD2	2:A:418:ALA:O	1.98	0.47
2:A:133:GLY:O	2:A:428:HIS:HE1	1.98	0.47
2:A:73:ARG:HD3	2:A:74:GLU:O	2.15	0.46
2:B:275:ILE:HA	2:B:290:SER:HA	1.98	0.46
2:A:248:LEU:HD13	2:A:263:ARG:HD3	1.97	0.45
2:B:236:ARG:NH2	2:B:254:GLU:HG3	2.32	0.45
1:D:146:GLU:CD	2:B:354:LYS:HZ1	2.21	0.44
2:B:343:PHE:HB3	2:B:357:ALA:HB2	2.00	0.43
1:H:146:GLU:CD	2:A:354:LYS:HZ1	2.21	0.43
1:D:159:ASN:HD22	2:B:177:GLN:NE2	2.16	0.42
2:A:275:ILE:HA	2:A:290:SER:HA	2.01	0.42
1:H:155:HIS:HE1	4:A:447:HOH:O	2.00	0.42
2:A:128:GLN:HE22	2:A:186:ARG:C	2.22	0.42
2:A:131:ASN:H	2:A:187:GLN:HE22	1.67	0.42
2:A:396:GLY:HA2	2:A:422:SER:O	2.20	0.41
2:B:128:GLN:HE22	2:B:186:ARG:C	2.23	0.41
2:B:396:GLY:HA2	2:B:422:SER:O	2.21	0.41
2:B:143:HIS:HD2	4:B:597:HOH:O	2.04	0.41
1:H:109:TRQ:HB2	1:H:160:TRP:HE1	1.86	0.41
2:A:186:ARG:HH22	2:A:426:GLN:NE2	2.19	0.40
1:D:155:HIS:CE1	4:B:451:HOH:O	2.71	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	107/135 (79%)	104 (97%)	3 (3%)	0	100	100
1	H	120/135 (89%)	114 (95%)	6 (5%)	0	100	100
2	A	360/361 (100%)	348 (97%)	11 (3%)	1 (0%)	46	19
2	B	359/361 (99%)	345 (96%)	12 (3%)	2 (1%)	30	8
All	All	946/992 (95%)	911 (96%)	32 (3%)	3 (0%)	46	19

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	216	VAL
2	A	200	SER
2	B	200	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	96/112 (86%)	96 (100%)	0	100	100
1	H	104/112 (93%)	104 (100%)	0	100	100
2	A	305/305 (100%)	303 (99%)	2 (1%)	88	70
2	B	301/305 (99%)	299 (99%)	2 (1%)	88	70
All	All	806/834 (97%)	802 (100%)	4 (0%)	92	78

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	104	ARG
2	A	308	ASP
2	B	104	ARG
2	B	388	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (38) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	139	GLN
1	D	143	GLN
1	D	155	HIS
1	D	164	ASN
1	H	139	GLN
1	H	143	GLN
1	H	155	HIS
1	H	164	ASN
2	A	86	GLN
2	A	106	HIS
2	A	124	ASN
2	A	128	GLN
2	A	143	HIS
2	A	177	GLN
2	A	180	ASN
2	A	187	GLN
2	A	197	GLN
2	A	231	GLN
2	A	266	GLN
2	A	386	GLN
2	A	424	GLN
2	A	426	GLN
2	A	428	HIS
2	B	86	GLN
2	B	106	HIS
2	B	124	ASN
2	B	128	GLN
2	B	143	HIS
2	B	177	GLN
2	B	180	ASN
2	B	187	GLN
2	B	197	GLN
2	B	231	GLN
2	B	266	GLN
2	B	386	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	424	GLN
2	B	426	GLN
2	B	428	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	TRQ	D	109	1,3	13,16,18	2.49	4 (30%)	8,22,26	1.66	3 (37%)
1	TRQ	H	109	1,3	13,16,18	2.55	4 (30%)	8,22,26	1.90	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TRQ	D	109	1,3	-	0/3/16/21	0/2/2/2
1	TRQ	H	109	1,3	-	0/3/16/21	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	109	TRQ	CH2-CZ3	-3.59	1.41	1.49
1	D	109	TRQ	CH2-CZ3	-3.31	1.41	1.49
1	H	109	TRQ	O7-CZ2	3.15	1.27	1.22
1	D	109	TRQ	O7-CZ2	3.16	1.27	1.22

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	109	TRQ	CE3-CZ3	4.18	1.42	1.33
1	D	109	TRQ	CE3-CZ3	4.27	1.42	1.33
1	D	109	TRQ	CD2-CG	5.73	1.47	1.40
1	H	109	TRQ	CD2-CG	6.19	1.48	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	109	TRQ	CB-CG-CD1	-3.08	124.17	127.97
1	D	109	TRQ	CB-CG-CD1	-2.66	124.68	127.97
1	D	109	TRQ	CD1-NE1-CE2	2.18	109.96	104.34
1	H	109	TRQ	CD1-NE1-CE2	2.33	110.37	104.34
1	D	109	TRQ	CB-CG-CD2	2.36	129.96	124.40
1	H	109	TRQ	CB-CG-CD2	2.61	130.55	124.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	109	TRQ	1	0
1	H	109	TRQ	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PHZ	D	1	1	8,8,8	1.38	1 (12%)	8,9,9	0.51	0
3	PHZ	H	1	1	8,8,8	1.33	1 (12%)	8,9,9	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PHZ	D	1	1	-	0/2/2/2	0/1/1/1
3	PHZ	H	1	1	-	0/2/2/2	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	1	PHZ	CG-N2	2.80	1.47	1.41
3	D	1	PHZ	CG-N2	3.10	1.48	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	D	108/135 (80%)	-0.41	0	100 100	10, 15, 23, 36	0
1	H	118/135 (87%)	-0.17	5 (4%)	40 38	11, 16, 35, 43	0
2	A	360/361 (99%)	-0.58	2 (0%)	90 89	8, 13, 22, 44	0
2	B	360/361 (99%)	-0.28	1 (0%)	94 93	9, 17, 29, 46	0
All	All	946/992 (95%)	-0.40	8 (0%)	87 86	8, 15, 28, 46	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	303	ASP	3.5
1	H	66	LEU	3.4
1	H	117	HIS	2.6
1	H	67	ALA	2.4
2	A	73	ARG	2.1
1	H	68	ASN	2.1
1	H	62	LEU	2.1
2	A	431	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	TRQ	D	109	15/17	0.93	0.07	-	12,14,16,17	0
1	TRQ	H	109	15/17	0.95	0.06	-	11,12,15,16	0



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PHZ	H	1	8/8	0.87	0.11	8.59	18,24,26,26	0
3	PHZ	D	1	8/8	0.85	0.12	7.32	19,22,23,26	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.