



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:05 PM GMT

PDB ID : 5AGK  
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with  
(2S)-2-Amino-5-(2-(methylsulfinyl)acetimidamido) pentanoic acid  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2015-02-02  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	 3% 82% 14% •
1	B	422	 % 89% 8% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	RGE	A	801[A]	-	-	-	X
4	RGE	A	801[B]	-	-	-	X
5	ACT	A	860	-	-	-	X
5	ACT	B	860	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, BRAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	0	4	1
			3329	2131	567	609	22			
1	B	411	Total	C	N	O	S	0	3	0
			3354	2146	574	612	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is (2S)-2-AMINO-5-(2-(METHYLSULFINYL)ACETIMIDAMIDO)PENTANOIC ACID (three-letter code: RGE) (formula: C<sub>8</sub>H<sub>17</sub>N<sub>3</sub>O<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	1
			18	9	3	4	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	1
			18	9	3	4	2		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	201	Total	O	0	0
			201	201		
7	B	216	Total	O	0	0
			216	216		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



#### • Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.75Å 110.90Å 164.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.37 – 2.00 49.37 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.1 (49.37-2.00) 95.3 (49.37-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.31 (at 2.00Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.171 , 0.215 0.178 , 0.218	Depositor DCC
$R_{free}$ test set	3089 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	44.1	Xtriage
Anisotropy	0.466	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 62049 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7265	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, ACT, H4B, RGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3434	0.51	0/4659
1	B	0.36	0/3456	0.50	0/4685
All	All	0.36	0/6890	0.51	0/9344

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3329	0	3242	37	0
1	B	3354	0	3274	21	0
2	A	43	0	30	4	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	18	0	10	5	0
4	B	18	0	10	4	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	201	0	0	2	0
7	B	216	0	0	1	0
All	All	7265	0	6632	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:565:PRO:HB2	4:A:801[B]:RGE:H52C	1.58	0.85
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.64	0.79
2:B:750:HEM:HH2	2:B:750:HEM:HBB2	1.74	0.70
1:A:567:VAL:HG23	4:A:801[B]:RGE:H51C	1.76	0.66
2:A:750:HEM:HBB2	2:A:750:HEM:HH2	1.79	0.64
1:A:631:VAL:HG11	1:B:628:GLN:HG3	1.78	0.64
1:B:322:LEU:HD13	1:B:699:ARG:HH21	1.64	0.63
1:A:350:THR:N	1:A:353:GLN:OE1	2.31	0.63
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.79	0.62
1:A:436:HIS:ND1	7:A:2065:HOH:O	2.31	0.61
1:B:567:VAL:HG23	4:B:801[B]:RGE:H51C	1.84	0.59
1:B:620:LYS:HD3	1:B:622:SER:H	1.71	0.56
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.89	0.54
1:B:706:TYR:OH	2:B:750:HEM:O2D	2.18	0.54
1:B:388:ILE:O	1:B:392:SER:N	2.37	0.52
1:A:316:LEU:HD12	1:A:319:LYS:HE2	1.92	0.51
1:A:552:ASP:OD1	1:A:555:LYS:NZ	2.44	0.50
1:B:414:ARG:HD3	1:B:678:TRP:CD2	2.46	0.49
1:B:404:GLY:HA3	1:B:574:ILE:HD13	1.95	0.48
1:A:567:VAL:HG23	4:A:801[B]:RGE:C5	2.42	0.48
1:B:565:PRO:O	4:B:801[B]:RGE:H53C	2.15	0.47
1:A:510:TRP:CE2	1:A:521:PRO:HD3	2.50	0.47
1:A:668[B]:CYS:HB2	7:A:2175:HOH:O	2.15	0.46
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.50	0.46
1:B:525:GLN:HE21	1:B:528:GLY:HA2	1.81	0.46
1:A:620:LYS:HB2	1:A:620:LYS:NZ	2.31	0.46
1:A:517:PHE:HB2	1:A:560:LYS:HE3	1.97	0.46
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.97	0.46
1:A:321:THR:HG21	1:A:701:THR:HG22	1.98	0.45
1:A:481:ARG:HD3	1:A:498:ASN:HD21	1.80	0.45
1:B:335:ILE:HB	1:B:338:PRO:HG3	1.99	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.99	0.45
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.99	0.45
1:A:369:ILE:HG13	1:A:371:ARG:HG3	1.98	0.44
1:A:628:GLN:NE2	1:B:632:GLU:OE2	2.50	0.44
1:A:450:THR:HA	1:A:455:LEU:HD22	1.99	0.44
1:A:706:TYR:OH	2:A:750:HEM:O2D	2.25	0.43
1:B:355:PHE:CE1	1:B:385:ASN:HB2	2.53	0.43
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.53	0.43
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.99	0.43
1:A:360:GLU:O	1:A:364:GLN:HG3	2.19	0.42
1:B:440:ASN:ND2	7:B:2077:HOH:O	2.52	0.42
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.07	0.42
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.54	0.42
1:A:676:TRP:CZ2	1:A:680:VAL:HG21	2.55	0.41
1:A:496:PRO:HA	1:A:499:VAL:HG23	2.02	0.41
1:A:475:TRP:CZ2	1:A:531:PRO:HG3	2.56	0.41
1:A:632:GLU:OE2	1:B:628:GLN:NE2	2.52	0.41
1:A:423:LYS:O	1:A:457[B]:SER:OG	2.38	0.41
1:A:709:ASP:HA	1:A:710:PRO:HD3	1.94	0.41
1:A:481:ARG:NH1	1:A:498:ASN:OD1	2.54	0.41
1:B:565:PRO:HB2	4:B:801[B]:RGE:H52C	2.02	0.41
1:A:595:VAL:O	1:A:599:CYS:HB2	2.21	0.41
1:B:595:VAL:O	1:B:599:CYS:HB2	2.21	0.40
2:B:750:HEM:NB	4:B:801[B]:RGE:H2A	2.37	0.40
1:A:414:ARG:HD3	1:A:678:TRP:CD2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	408/422 (97%)	396 (97%)	12 (3%)	0	100 100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	410/422 (97%)	401 (98%)	9 (2%)	0	100	100
All	All	818/844 (97%)	797 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/377 (97%)	356 (97%)	11 (3%)	48	47
1	B	369/377 (98%)	360 (98%)	9 (2%)	57	58
All	All	736/754 (98%)	716 (97%)	20 (3%)	52	52

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	321	THR
1	A	523	LEU
1	A	527	ASN
1	A	535	GLN
1	A	547	ARG
1	A	550	LYS
1	A	608	GLU
1	A	612	LYS
1	A	645	LYS
1	A	663	GLU
1	A	715	VAL
1	B	315	THR
1	B	337	LEU
1	B	351	LYS
1	B	353	GLN
1	B	392	SER
1	B	540	LEU
1	B	547	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	550	LYS
1	B	620	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	535	GLN
1	B	642	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	750	1	30,50,50	2.20	8 (26%)	24,82,82	2.57	9 (37%)
3	H4B	A	760	-	13,18,18	0.75	0	11,26,26	2.30	5 (45%)
4	RGE	A	801[A]	-	9,14,14	2.85	3 (33%)	8,17,17	3.12	2 (25%)
4	RGE	A	801[B]	-	9,14,14	3.02	3 (33%)	8,17,17	2.61	2 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	ACT	A	860	-	1,3,3	1.34	0	0,3,3	0.00	-
2	HEM	B	750	1	30,50,50	2.04	8 (26%)	24,82,82	2.56	9 (37%)
3	H4B	B	760	-	13,18,18	0.77	0	11,26,26	2.34	5 (45%)
4	RGE	B	801[A]	-	9,14,14	2.68	3 (33%)	8,17,17	3.35	3 (37%)
4	RGE	B	801[B]	-	9,14,14	3.19	3 (33%)	8,17,17	3.11	3 (37%)
5	ACT	B	860	-	1,3,3	0.98	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	RGE	A	801[A]	-	-	0/8/15/15	0/0/0/0
4	RGE	A	801[B]	-	-	0/8/15/15	0/0/0/0
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/10/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	RGE	B	801[A]	-	-	0/8/15/15	0/0/0/0
4	RGE	B	801[B]	-	-	0/8/15/15	0/0/0/0
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	801[B]	RGE	C2-S3	-7.32	1.65	1.80
2	A	750	HEM	C3B-C4B	-6.61	1.46	1.51
4	A	801[B]	RGE	C2-S3	-6.55	1.67	1.80
4	A	801[A]	RGE	C2-S3	-6.34	1.67	1.80
2	B	750	HEM	C3B-C4B	-6.06	1.46	1.51
2	A	750	HEM	C3D-C4D	-5.71	1.44	1.51
4	B	801[A]	RGE	C2-S3	-5.70	1.68	1.80
2	B	750	HEM	C3D-C4D	-5.10	1.45	1.51
2	B	750	HEM	C2C-C1C	-3.85	1.45	1.52
2	A	750	HEM	C2C-C1C	-3.80	1.45	1.52
4	A	801[B]	RGE	C5-S3	-2.89	1.65	1.77
4	B	801[B]	RGE	C5-S3	-2.78	1.65	1.77
4	B	801[A]	RGE	C5-S3	-2.76	1.66	1.77
4	A	801[A]	RGE	C5-S3	-2.59	1.66	1.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C2B-C1B	-2.19	1.44	1.51
2	B	750	HEM	C2B-C1B	-2.08	1.45	1.51
2	B	750	HEM	C2D-C1D	-2.04	1.45	1.51
2	A	750	HEM	C3C-CAC	2.02	1.55	1.51
2	B	750	HEM	C3C-CAC	2.06	1.55	1.51
2	B	750	HEM	FE-ND	2.12	2.08	1.97
2	A	750	HEM	C3B-CAB	2.15	1.55	1.51
2	B	750	HEM	C1C-NC	2.16	1.38	1.36
2	A	750	HEM	FE-NB	2.43	2.10	1.97
2	A	750	HEM	FE-NC	3.38	2.09	1.95
4	B	801[A]	RGE	O4-S3	4.86	1.62	1.51
4	A	801[A]	RGE	O4-S3	5.08	1.63	1.51
4	B	801[B]	RGE	O4-S3	5.43	1.64	1.51
4	A	801[B]	RGE	O4-S3	5.50	1.64	1.51

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	CBA-CAA-C2A	-6.28	101.28	112.53
2	A	750	HEM	CBA-CAA-C2A	-6.25	101.33	112.53
3	A	760	H4B	N3-C2-N1	-3.23	120.24	125.53
3	B	760	H4B	N3-C2-N1	-3.12	120.42	125.53
2	B	750	HEM	CBD-CAD-C3D	-2.58	106.05	113.55
2	A	750	HEM	CBD-CAD-C3D	-2.53	106.20	113.55
4	B	801[A]	RGE	C1-C2-S3	-2.15	103.28	109.68
4	B	801[B]	RGE	O4-S3-C2	2.08	109.19	106.31
3	A	760	H4B	C4A-C8A-N8	2.10	120.91	118.43
3	B	760	H4B	N2-C2-N1	2.16	120.78	117.20
4	B	801[B]	RGE	C1-C2-S3	2.18	116.16	109.68
4	A	801[B]	RGE	O4-S3-C2	2.22	109.39	106.31
3	B	760	H4B	C2-N1-C8A	2.29	119.68	114.54
2	A	750	HEM	CMD-C2D-C3D	2.42	125.03	114.35
2	B	750	HEM	C2D-C3D-C4D	2.47	105.69	101.50
2	B	750	HEM	C3B-C4B-CHC	2.52	126.70	123.16
2	A	750	HEM	C2D-C3D-C4D	2.55	105.82	101.50
4	B	801[A]	RGE	O4-S3-C5	2.78	111.02	106.41
3	A	760	H4B	C4-N3-C2	2.89	119.95	115.94
4	A	801[A]	RGE	O4-S3-C5	2.96	111.31	106.41
2	B	750	HEM	CMD-C2D-C3D	2.98	127.53	114.35
2	A	750	HEM	CMB-C2B-C3B	3.00	124.01	116.53
3	A	760	H4B	C2-N1-C8A	3.14	121.59	114.54
2	A	750	HEM	C3B-C4B-CHC	3.15	127.60	123.16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	760	H4B	C4-N3-C2	3.82	121.25	115.94
2	B	750	HEM	CMB-C2B-C3B	3.83	126.08	116.53
2	B	750	HEM	CAD-C3D-C4D	4.27	127.52	112.47
2	A	750	HEM	CAD-C3D-C4D	4.34	127.78	112.47
3	A	760	H4B	C4-C4A-C8A	4.40	118.55	114.56
2	A	750	HEM	CAD-C3D-C2D	4.57	126.37	113.22
3	B	760	H4B	C4-C4A-C8A	4.60	118.73	114.56
2	B	750	HEM	CAD-C3D-C2D	4.72	126.78	113.22
2	B	750	HEM	CMC-C2C-C3C	4.84	128.62	116.53
2	A	750	HEM	CMC-C2C-C3C	5.41	130.02	116.53
4	A	801[B]	RGE	C5-S3-C2	6.79	107.71	97.84
4	B	801[B]	RGE	C5-S3-C2	8.14	109.68	97.84
4	A	801[A]	RGE	C5-S3-C2	8.15	109.69	97.84
4	B	801[A]	RGE	C5-S3-C2	8.69	110.48	97.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	4	0
4	A	801[A]	RGE	1	0
4	A	801[B]	RGE	4	0
2	B	750	HEM	3	0
4	B	801[B]	RGE	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	408/422 (96%)	0.24	11 (2%) 58 58	18, 43, 88, 113	0
1	B	411/422 (97%)	0.11	6 (1%) 76 77	20, 36, 69, 101	0
All	All	819/844 (97%)	0.18	17 (2%) 67 67	18, 39, 82, 113	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	4.8
1	A	355	PHE	4.2
1	A	299	ARG	4.2
1	A	715	VAL	4.1
1	A	551	PHE	3.8
1	A	322	LEU	3.3
1	B	299	ARG	3.1
1	A	388	ILE	2.8
1	A	554	PHE	2.8
1	A	369	ILE	2.5
1	A	300	PHE	2.5
1	B	355	PHE	2.4
1	A	321	THR	2.4
1	B	352	ASP	2.3
1	A	394	TYR	2.3
1	B	321	THR	2.2
1	B	351	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	ACT	A	860	4/4	0.95	0.27	13.31	47,53,53,53	0
5	ACT	B	860	4/4	0.95	0.22	3.51	43,44,45,46	0
4	RGE	A	801[B]	15/15	0.97	0.15	2.21	23,34,39,42	3
4	RGE	A	801[A]	15/15	0.97	0.15	2.21	13,34,39,42	3
6	ZN	B	900	1/1	0.99	0.12	1.57	33,33,33,33	0
2	HEM	B	750	43/43	0.98	0.16	1.43	15,26,33,44	0
2	HEM	A	750	43/43	0.98	0.13	0.61	14,26,39,44	0
4	RGE	B	801[A]	15/15	0.97	0.13	0.15	16,30,36,41	3
4	RGE	B	801[B]	15/15	0.97	0.13	0.15	23,32,41,45	3
3	H4B	B	760	17/17	0.95	0.13	-0.21	20,25,30,36	0
3	H4B	A	760	17/17	0.93	0.11	-0.87	22,27,30,33	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.