



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:07 PM GMT

PDB ID : 5AH2
Title : The sliding clamp of Mycobacterium smegmatis in complex with a natural product.
Authors : Lukat, P.; Kling, A.; Heinz, D.W.; Mueller, R.
Deposited on : 2015-02-04
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4587 (2.14-2.10)
Clashscore	102246	5132 (2.14-2.10)
Ramachandran outliers	100387	5080 (2.14-2.10)
Sidechain outliers	100360	5081 (2.14-2.10)
RSRZ outliers	91569	4597 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>5%</div> </div> </div>
1	B	401	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>6%</div> </div> </div>
1	C	401	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>
1	D	401	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>
2	E	11	<div> <div></div> <div> <div>55%</div> <div>27%</div> <div>18%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	11	
2	G	11	
2	H	11	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NA	A	1398	-	-	-	X
3	NA	C	1398	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE III SUBUNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	11	0
			2842	1804	484	550	4			
1	B	377	Total	C	N	O	S	0	10	0
			2791	1782	472	533	4			
1	C	382	Total	C	N	O	S	0	4	0
			2784	1770	469	541	4			
1	D	379	Total	C	N	O	S	0	9	0
			2766	1765	477	520	4			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP A0QND6
A	-2	GLY	-	EXPRESSION TAG	UNP A0QND6
A	-1	GLY	-	EXPRESSION TAG	UNP A0QND6
A	0	ARG	-	EXPRESSION TAG	UNP A0QND6
B	-3	GLY	-	EXPRESSION TAG	UNP A0QND6
B	-2	GLY	-	EXPRESSION TAG	UNP A0QND6
B	-1	GLY	-	EXPRESSION TAG	UNP A0QND6
B	0	ARG	-	EXPRESSION TAG	UNP A0QND6
C	-3	GLY	-	EXPRESSION TAG	UNP A0QND6
C	-2	GLY	-	EXPRESSION TAG	UNP A0QND6
C	-1	GLY	-	EXPRESSION TAG	UNP A0QND6
C	0	ARG	-	EXPRESSION TAG	UNP A0QND6
D	-3	GLY	-	EXPRESSION TAG	UNP A0QND6
D	-2	GLY	-	EXPRESSION TAG	UNP A0QND6
D	-1	GLY	-	EXPRESSION TAG	UNP A0QND6
D	0	ARG	-	EXPRESSION TAG	UNP A0QND6

- Molecule 2 is a protein called GRISELIMYCIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	11	Total	C	N	O	0	0	0
			79	57	10	12			
2	F	11	Total	C	N	O	0	0	0
			79	57	10	12			
2	G	11	Total	C	N	O	0	0	0
			79	57	10	12			
2	H	11	Total	C	N	O	0	0	0
			79	57	10	12			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	273	Total	O	0	0
			273	273		
4	B	190	Total	O	0	0
			190	190		
4	C	183	Total	O	0	0
			183	183		
4	D	124	Total	O	0	0
			124	124		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA POLYMERASE III SUBUNIT BETA




- Molecule 1: DNA POLYMERASE III SUBUNIT BETA



- Molecule 1: DNA POLYMERASE III SUBUNIT BETA



- Molecule 1: DNA POLYMERASE III SUBUNIT BETA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.24Å 125.87Å 94.90Å 90.00° 104.42° 90.00°	Depositor
Resolution (Å)	77.71 – 2.13 77.71 – 2.13	Depositor EDS
% Data completeness (in resolution range)	100.0 (77.71-2.13) 100.0 (77.71-2.13)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.12Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.182 , 0.231 0.192 , 0.238	Depositor DCC
R_{free} test set	5071 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	34.0	Xtriage
Anisotropy	0.158	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 102012 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12271	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NA, NZC, MVA, MLU, MP8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2906	0.56	0/3965
1	B	0.40	0/2863	0.56	0/3906
1	C	0.39	0/2838	0.54	0/3875
1	D	0.35	0/2835	0.51	0/3872
2	E	0.52	0/24	1.21	0/26
2	F	0.53	0/24	1.39	0/26
2	G	0.56	0/24	1.20	0/26
2	H	0.45	0/24	1.13	0/26
All	All	0.39	0/11538	0.55	0/15722

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	E	0	1
2	F	0	1
2	G	0	1
2	H	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	E	8	MVA	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	F	8	MVA	Peptide
2	G	8	MVA	Peptide
2	H	8	MVA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2842	0	2888	23	0
1	B	2791	0	2865	21	0
1	C	2784	0	2800	27	0
1	D	2766	0	2819	21	0
2	E	79	0	96	6	0
2	F	79	0	96	6	0
2	G	79	0	96	8	0
2	H	79	0	96	4	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	273	0	0	7	0
4	B	190	0	0	4	0
4	C	183	0	0	4	0
4	D	124	0	0	0	0
All	All	12271	0	11756	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:79[B]:VAL:HG11	1:B:105:VAL:HG11	1.71	0.72
1:B:348[A]:GLU:HG2	1:B:349:ARG:HG3	1.72	0.71
1:A:391:MET:HB2	2:E:5:LEU:HD23	1.75	0.69
1:A:92:PRO:HB2	1:A:94:LYS:HD2	1.75	0.69
1:B:181:ARG:HD3	2:F:10:MLU:HD22	1.76	0.68
1:D:391:MET:HB2	2:H:5:LEU:HD23	1.77	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:LEU:HD11	2:E:7:LEU:HD11	1.80	0.62
1:A:181:ARG:HD3	2:E:10:MLU:HD22	1.82	0.61
1:A:364[A]:ARG:NH1	1:A:384:THR:OG1	2.29	0.61
1:B:30:SER:OG	4:B:2016:HOH:O	2.15	0.60
1:A:159:LEU:HD12	1:A:161:MET:HG2	1.83	0.59
1:A:105:VAL:HG23	1:A:118:THR:HG22	1.85	0.58
1:D:14:ARG:NH1	1:D:70:GLU:OE1	2.37	0.57
1:D:349[B]:ARG:HB2	1:D:366:ALA:HB3	1.85	0.57
1:C:391:MET:HB2	2:G:5:LEU:HD23	1.86	0.57
1:B:262:LEU:HD11	2:F:7:LEU:HD11	1.87	0.56
1:D:181:ARG:HD3	2:H:10:MLU:HD22	1.86	0.56
1:B:280:GLU:OE2	1:B:284[A]:ARG:NH2	2.34	0.56
1:B:28:ALA:HA	1:B:31:LEU:HG	1.88	0.56
1:A:246[B]:ARG:NH1	4:A:2100:HOH:O	2.39	0.55
1:B:31:LEU:HD11	1:B:43[B]:VAL:HG23	1.87	0.55
1:B:36:THR:O	1:B:38:PRO:HD3	2.07	0.54
1:C:161:MET:O	4:C:2089:HOH:O	2.19	0.54
1:A:364[B]:ARG:NH1	4:A:2201:HOH:O	2.31	0.53
1:C:37:ILE:HG22	1:C:39:VAL:HG12	1.91	0.53
1:D:92:PRO:HB2	1:D:94:LYS:HG2	1.93	0.51
1:D:91:LEU:HD11	1:D:107:LEU:HD21	1.91	0.51
1:B:124:TYR:OH	4:B:2024:HOH:O	2.19	0.51
1:B:55:SER:HB3	1:B:64:GLU:HG3	1.93	0.51
1:C:113:ARG:NH1	4:C:2045:HOH:O	2.43	0.50
1:B:31:LEU:HD11	1:B:43[A]:VAL:HG13	1.94	0.50
1:A:86:ASP:OD1	4:A:2067:HOH:O	2.20	0.50
1:C:297:MET:HB3	1:C:304:LEU:HD11	1.94	0.49
1:D:26:TRP:CE2	1:D:245:LYS:HD3	2.48	0.49
1:A:22:ASP:OD1	4:A:2015:HOH:O	2.20	0.49
1:D:223:HIS:HB2	1:D:241[A]:ARG:HB2	1.95	0.49
1:A:94:LYS:HD3	1:A:110:GLY:HA3	1.94	0.48
1:A:349[B]:ARG:HB2	1:A:366:ALA:HB3	1.95	0.48
1:C:271:ALA:HB2	1:C:324:PHE:HD1	1.78	0.48
1:A:394:ARG:HB3	2:E:1:ACE:H3	1.96	0.48
2:G:2:MVA:HA	2:G:3:MP8:HD	1.73	0.47
1:B:391:MET:HB2	2:F:5:LEU:HD23	1.97	0.47
1:C:305:LYS:NZ	1:C:319[B]:ASP:OD1	2.37	0.47
1:C:90:ALA:O	4:C:2042:HOH:O	2.19	0.47
1:D:137:ILE:HD11	1:D:141:LEU:HG	1.97	0.46
1:B:284[A]:ARG:HD2	1:B:318:GLU:OE1	2.14	0.46
2:F:10:MLU:HCN3	2:F:10:MLU:HB2	1.57	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:148:GLN:O	1:D:151:VAL:HG12	2.15	0.46
1:D:272:THR:HG22	1:D:349[A]:ARG:HE	1.81	0.46
1:C:78:LEU:HD23	1:C:119:LEU:HB2	1.97	0.45
1:D:27:VAL:HG21	1:D:54:ILE:HG22	1.98	0.45
1:C:30:SER:OG	4:C:2011:HOH:O	2.21	0.45
1:C:290:ASP:O	1:C:293:ALA:N	2.50	0.45
1:C:280:GLU:OE2	1:C:284:ARG:NH1	2.27	0.45
2:E:10:MLU:HA	2:E:10:MLU:HD23	1.72	0.45
1:D:25:ALA:O	1:D:29:ARG:HG3	2.17	0.45
1:D:130:LEU:HD13	1:D:246[B]:ARG:NH2	2.31	0.45
1:C:181:ARG:HD3	2:G:10:MLU:HD22	1.99	0.44
2:F:10:MLU:HD23	2:F:10:MLU:HA	1.70	0.44
1:D:133:GLU:HB3	1:D:194:THR:HG21	2.00	0.44
2:H:10:MLU:HCN3	2:H:10:MLU:HB2	1.65	0.44
2:G:3:MP8:HA	2:G:4:NZC:H40	1.68	0.44
1:C:139:SER:HB3	1:C:222:VAL:HG13	2.00	0.44
1:C:262:LEU:HD11	2:G:7:LEU:CD2	2.48	0.44
1:B:46:THR:HB	1:B:53[A]:THR:OG1	2.18	0.44
1:A:31:LEU:HD11	1:A:43:VAL:HG23	1.99	0.44
1:D:43[B]:VAL:HG23	1:D:81:GLY:HA2	1.99	0.44
1:C:57:PHE:HB2	1:C:127:LEU:HD12	1.99	0.44
1:A:148:GLN:HG2	1:A:344:SER:O	2.18	0.43
2:G:4:NZC:O	2:G:4:NZC:H40B	2.18	0.43
1:C:58:ASP:O	1:C:59:TYR:HB2	2.19	0.43
1:A:111:SER:HB2	4:A:2079:HOH:O	2.18	0.43
1:D:43[A]:VAL:HG22	1:D:81:GLY:HA2	2.01	0.43
1:A:263:LEU:HA	1:A:264:PRO:HD3	1.84	0.43
1:A:100:VAL:HG22	1:A:105:VAL:HG22	1.99	0.43
1:B:57:PHE:HB2	1:B:127:LEU:HD23	2.01	0.43
1:C:168:GLU:HB2	1:C:175:VAL:HB	2.00	0.43
2:F:3:MP8:HA	2:F:4:NZC:H40	1.68	0.42
1:C:258:LYS:HD2	1:C:258:LYS:HA	1.64	0.42
1:B:58:ASP:O	1:B:59:TYR:HB2	2.19	0.42
1:D:108[A]:THR:HG23	1:D:113:ARG:HG2	2.01	0.42
1:C:263:LEU:HA	1:C:264:PRO:HD3	1.88	0.42
1:A:58:ASP:O	1:A:59:TYR:HB2	2.19	0.42
1:B:385:ASP:OD1	4:B:2186:HOH:O	2.21	0.42
2:E:10:MLU:HCN3	2:E:10:MLU:HB2	1.63	0.42
1:A:89:LYS:HG3	4:A:2014:HOH:O	2.20	0.41
1:C:17:ARG:HG3	1:C:18:GLU:N	2.36	0.41
1:B:79[B]:VAL:CG1	1:B:105:VAL:HG11	2.47	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:159:LEU:CD1	1:C:161:MET:HG3	2.50	0.41
2:G:7:LEU:HA	2:G:8:MVA:HN1	1.80	0.41
1:C:180:ASP:O	1:C:181:ARG:HB2	2.20	0.41
1:D:139:SER:OG	1:D:217:THR:OG1	2.37	0.41
4:A:2183:HOH:O	1:D:161:MET:HG2	2.21	0.41
2:H:7:LEU:HA	2:H:8:MVA:HN1	1.81	0.41
1:C:105:VAL:HG23	1:C:118:THR:HG22	2.02	0.41
1:B:161:MET:HG2	4:B:2099:HOH:O	2.20	0.41
1:B:180:ASP:O	1:B:181:ARG:HB2	2.21	0.41
1:A:52:LEU:HB3	1:A:67[B]:VAL:HG22	2.02	0.41
1:C:57:PHE:CZ	1:C:59:TYR:HA	2.57	0.40
1:C:262:LEU:HD11	2:G:7:LEU:HD23	2.03	0.40
1:A:306:LEU:O	1:A:317:GLU:HA	2.21	0.40
1:C:298:GLU:HG3	1:C:330:THR:OG1	2.22	0.40
1:D:57:PHE:CZ	1:D:59:TYR:HA	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/401 (96%)	376 (98%)	7 (2%)	2 (0%)	34	29
1	B	379/401 (94%)	371 (98%)	8 (2%)	0	100	100
1	C	378/401 (94%)	367 (97%)	11 (3%)	0	100	100
1	D	380/401 (95%)	371 (98%)	9 (2%)	0	100	100
2	E	3/11 (27%)	2 (67%)	1 (33%)	0	100	100
2	F	3/11 (27%)	1 (33%)	1 (33%)	1 (33%)	0	0
2	G	3/11 (27%)	1 (33%)	1 (33%)	1 (33%)	0	0
2	H	3/11 (27%)	1 (33%)	1 (33%)	1 (33%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1534/1648 (93%)	1490 (97%)	39 (2%)	5 (0%)	46	44

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	9	PRO
2	G	9	PRO
2	H	9	PRO
1	A	327[A]	GLU
1	A	327[B]	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/306 (97%)	290 (98%)	7 (2%)	57	60
1	B	293/306 (96%)	285 (97%)	8 (3%)	52	55
1	C	287/306 (94%)	271 (94%)	16 (6%)	26	22
1	D	283/306 (92%)	271 (96%)	12 (4%)	36	35
2	E	3/3 (100%)	1 (33%)	2 (67%)	0	0
2	F	3/3 (100%)	2 (67%)	1 (33%)	0	0
2	G	3/3 (100%)	1 (33%)	2 (67%)	0	0
2	H	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	1172/1236 (95%)	1123 (96%)	49 (4%)	37	35

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	LEU
1	A	183	ARG
1	A	191	TRP
1	A	237	LEU
1	A	253	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	291	ARG
1	A	313	VAL
1	B	119	LEU
1	B	121	VAL
1	B	183	ARG
1	B	191	TRP
1	B	199	GLU
1	B	296	ARG
1	B	312	ASP
1	B	358	SER
1	C	17	ARG
1	C	50	GLU
1	C	101	GLU
1	C	159	LEU
1	C	161	MET
1	C	183	ARG
1	C	191	TRP
1	C	231	SER
1	C	237	LEU
1	C	258	LYS
1	C	312	ASP
1	C	319[A]	ASP
1	C	319[B]	ASP
1	C	329	LEU
1	C	376	SER
1	C	395	LEU
1	D	17	ARG
1	D	36	THR
1	D	43[A]	VAL
1	D	43[B]	VAL
1	D	119	LEU
1	D	155	ARG
1	D	183	ARG
1	D	191	TRP
1	D	198	VAL
1	D	258	LYS
1	D	327	GLU
1	D	329	LEU
2	E	5	LEU
2	E	7	LEU
2	F	5	LEU
2	G	5	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	G	7	LEU
2	H	5	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

24 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MLU	E	10	2	7,8,9	0.54	0	4,9,11	0.96	0
2	MVA	E	2	2	6,7,8	1.10	1 (16%)	6,8,10	2.05	1 (16%)
2	MP8	E	3	2	7,8,9	0.69	0	5,10,12	1.32	1 (20%)
2	NZC	E	4	2	6,7,8	0.85	0	6,8,10	1.07	0
2	MP8	E	6	2	7,8,9	0.50	0	5,10,12	1.48	0
2	MVA	E	8	2	6,7,8	0.87	0	6,8,10	1.13	1 (16%)
2	MLU	F	10	2	7,8,9	0.39	0	4,9,11	1.00	0
2	MVA	F	2	2	6,7,8	0.90	0	6,8,10	1.59	2 (33%)
2	MP8	F	3	2	7,8,9	0.51	0	5,10,12	1.10	1 (20%)
2	NZC	F	4	2	6,7,8	0.90	0	6,8,10	1.19	0
2	MP8	F	6	2	7,8,9	0.56	0	5,10,12	1.23	1 (20%)
2	MVA	F	8	2	6,7,8	0.74	0	6,8,10	1.22	1 (16%)
2	MLU	G	10	2	7,8,9	0.55	0	4,9,11	1.32	1 (25%)
2	MVA	G	2	2	6,7,8	0.81	0	6,8,10	2.16	1 (16%)
2	MP8	G	3	2	7,8,9	0.70	0	5,10,12	1.43	1 (20%)
2	NZC	G	4	2	6,7,8	0.95	0	6,8,10	0.96	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MP8	G	6	2	7,8,9	0.69	0	5,10,12	1.53	1 (20%)
2	MVA	G	8	2	6,7,8	0.99	0	6,8,10	1.56	1 (16%)
2	MLU	H	10	2	7,8,9	0.55	0	4,9,11	0.97	0
2	MVA	H	2	2	6,7,8	0.89	0	6,8,10	2.47	2 (33%)
2	MP8	H	3	2	7,8,9	0.98	1 (14%)	5,10,12	1.85	2 (40%)
2	NZC	H	4	2	6,7,8	0.91	0	6,8,10	1.00	0
2	MP8	H	6	2	7,8,9	0.46	0	5,10,12	1.24	1 (20%)
2	MVA	H	8	2	6,7,8	0.90	0	6,8,10	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLU	E	10	2	-	0/4/8/10	0/0/0/0
2	MVA	E	2	2	-	0/5/8/10	0/0/0/0
2	MP8	E	3	2	-	0/0/11/13	0/1/1/1
2	NZC	E	4	2	-	0/5/8/10	0/0/0/0
2	MP8	E	6	2	-	0/0/11/13	0/1/1/1
2	MVA	E	8	2	-	0/5/8/10	0/0/0/0
2	MLU	F	10	2	-	0/4/8/10	0/0/0/0
2	MVA	F	2	2	-	0/5/8/10	0/0/0/0
2	MP8	F	3	2	-	0/0/11/13	0/1/1/1
2	NZC	F	4	2	-	0/5/8/10	0/0/0/0
2	MP8	F	6	2	-	0/0/11/13	0/1/1/1
2	MVA	F	8	2	-	0/5/8/10	0/0/0/0
2	MLU	G	10	2	-	0/4/8/10	0/0/0/0
2	MVA	G	2	2	-	0/5/8/10	0/0/0/0
2	MP8	G	3	2	-	0/0/11/13	0/1/1/1
2	NZC	G	4	2	-	0/5/8/10	0/0/0/0
2	MP8	G	6	2	-	0/0/11/13	0/1/1/1
2	MVA	G	8	2	-	0/5/8/10	0/0/0/0
2	MLU	H	10	2	-	0/4/8/10	0/0/0/0
2	MVA	H	2	2	-	0/5/8/10	0/0/0/0
2	MP8	H	3	2	-	0/0/11/13	0/1/1/1
2	NZC	H	4	2	-	0/5/8/10	0/0/0/0
2	MP8	H	6	2	-	0/0/11/13	0/1/1/1
2	MVA	H	8	2	-	0/5/8/10	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2	MVA	CB-CA	-2.27	1.52	1.54
2	H	3	MP8	CA-N	2.45	1.50	1.47

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	2	MVA	CB-CA-N	-5.33	102.34	111.57
2	E	2	MVA	CB-CA-N	-4.36	104.02	111.57
2	G	8	MVA	CB-CA-N	-3.29	105.88	111.57
2	F	2	MVA	CB-CA-N	-3.02	106.35	111.57
2	G	6	MP8	CG-CD-N	-3.00	98.96	104.48
2	G	10	MLU	O-C-CA	-2.55	118.71	125.44
2	F	8	MVA	CB-CA-N	-2.48	107.28	111.57
2	H	3	MP8	O-C-CA	-2.14	119.80	125.44
2	E	8	MVA	CB-CA-N	-2.12	107.89	111.57
2	F	6	MP8	O-C-CA	-2.08	119.94	125.44
2	H	2	MVA	O-C-CA	-2.07	119.38	125.74
2	H	6	MP8	CG-CD-N	-2.07	100.68	104.48
2	F	2	MVA	O-C-CA	-2.02	119.53	125.74
2	F	3	MP8	CB-CG-CD	2.04	104.93	102.21
2	E	3	MP8	CB-CG-CD	2.04	104.93	102.21
2	G	3	MP8	CB-CG-CD	2.08	104.98	102.21
2	H	3	MP8	CB-CG-CD	3.21	106.49	102.21
2	G	2	MVA	CG2-CB-CA	4.85	118.84	111.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	10	MLU	3	0
2	F	10	MLU	3	0
2	F	3	MP8	1	0
2	F	4	NZC	1	0
2	G	10	MLU	1	0
2	G	2	MVA	1	0
2	G	3	MP8	2	0
2	G	4	NZC	2	0
2	G	8	MVA	1	0
2	H	10	MLU	2	0
2	H	8	MVA	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/401 (94%)	0.25	7 (1%) 71 76	21, 36, 68, 94	0
1	B	377/401 (94%)	0.25	10 (2%) 58 65	22, 37, 67, 96	0
1	C	382/401 (95%)	0.30	10 (2%) 59 66	25, 43, 75, 104	0
1	D	379/401 (94%)	0.37	11 (2%) 55 63	27, 48, 80, 102	0
2	E	4/11 (36%)	0.35	0 100 100	28, 33, 39, 50	0
2	F	4/11 (36%)	0.05	0 100 100	31, 32, 47, 56	0
2	G	4/11 (36%)	0.25	0 100 100	31, 36, 46, 52	0
2	H	4/11 (36%)	0.06	0 100 100	32, 33, 48, 56	0
All	All	1534/1648 (93%)	0.29	38 (2%) 61 67	21, 41, 73, 104	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	374	GLY	5.7
1	D	292	GLY	5.3
1	C	290	ASP	4.0
1	B	36	THR	3.9
1	D	290	ASP	3.7
1	C	157	ASP	3.2
1	B	32	PRO	3.2
1	D	194	THR	2.9
1	B	292	GLY	2.9
1	B	33	THR	2.7
1	A	301	ASP	2.7
1	B	30	SER	2.7
1	B	396	PRO	2.6
1	D	291	ARG	2.6
1	A	290	ASP	2.6
1	B	367	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	217	THR	2.5
1	A	311	ASP	2.5
1	C	310	ALA	2.5
1	D	16	VAL	2.5
1	A	33	THR	2.4
1	A	375	GLY	2.4
1	C	270	VAL	2.4
1	C	158	THR	2.3
1	A	217	THR	2.3
1	C	49	ASP	2.2
1	C	8	LEU	2.2
1	C	33	THR	2.2
1	A	215	ALA	2.2
1	D	198	VAL	2.1
1	D	246[A]	ARG	2.1
1	C	36	THR	2.1
1	B	37	ILE	2.1
1	C	31	LEU	2.1
1	B	290	ASP	2.1
1	D	15	VAL	2.1
1	B	311	ASP	2.0
1	D	293	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NZC	F	4	8/9	0.94	0.12	-	28,31,32,33	0
2	MVA	G	2	8/9	0.87	0.16	-	31,33,36,37	0
2	MP8	F	3	8/9	0.94	0.15	-	27,28,30,31	0
2	MVA	E	8	8/9	0.90	0.16	-	44,48,52,55	0
2	MVA	F	8	8/9	0.85	0.17	-	53,56,56,57	0
2	MP8	G	6	8/9	0.93	0.09	-	39,41,45,46	0
2	MP8	H	6	8/9	0.92	0.13	-	35,38,41,42	0
2	MP8	G	3	8/9	0.95	0.12	-	26,29,32,32	0
2	MVA	H	2	8/9	0.92	0.10	-	33,35,37,38	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MVA	G	8	8/9	0.91	0.19	-	50,57,58,63	0
2	NZC	E	4	8/9	0.94	0.14	-	28,29,30,30	0
2	MP8	F	6	8/9	0.93	0.10	-	36,39,42,43	0
2	MP8	E	3	8/9	0.95	0.14	-	25,28,29,29	0
2	MLU	G	10	9/10	0.94	0.19	-	46,48,53,54	0
2	MLU	E	10	9/10	0.95	0.10	-	40,49,52,53	0
2	MVA	H	8	8/9	0.87	0.20	-	54,61,63,65	0
2	MP8	H	3	8/9	0.93	0.13	-	25,29,30,32	0
2	NZC	H	4	8/9	0.98	0.12	-	27,28,31,31	0
2	MVA	F	2	8/9	0.95	0.11	-	31,33,34,35	0
2	NZC	G	4	8/9	0.94	0.12	-	24,27,29,32	0
2	MLU	H	10	9/10	0.93	0.16	-	35,48,54,55	0
2	MVA	E	2	8/9	0.93	0.15	-	30,32,35,36	0
2	MLU	F	10	9/10	0.96	0.14	-	36,44,48,50	0
2	MP8	E	6	8/9	0.94	0.15	-	29,34,36,38	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NA	C	1398	1/1	0.93	0.42	15.50	56,56,56,56	0
3	NA	A	1398	1/1	0.92	0.19	8.07	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.