



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:28 AM GMT

PDB ID : 2AJ8
Title : Porcine dipeptidyl peptidase IV (CD26) in complex with 7-Benzyl-1,3-dimethyl-8-piperazin-1-yl-3,7-dihydro-purine-2,6-dione (BDPX)
Authors : Engel, M.; Hoffmann, T.; Manhart, S.; Heiser, U.; Chambre, S.; Huber, R.; Demuth, H.U.; Bode, W.
Deposited on : 2005-08-01
Resolution : 2.11 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4587 (2.14-2.10)
Clashscore	102246	5132 (2.14-2.10)
Ramachandran outliers	100387	5080 (2.14-2.10)
Sidechain outliers	100360	5081 (2.14-2.10)
RSRZ outliers	91569	4597 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>21%</div> </div> <div>•</div> </div>
1	B	728	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>17%</div> </div> <div>•</div> </div>
1	C	728	<div> <div>4%</div> <div> <div></div> <div>79%</div> <div>19%</div> </div> <div>•</div> </div>
1	D	728	<div> <div>3%</div> <div> <div></div> <div>79%</div> <div>19%</div> </div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	C	772(A)	-	-	-	X
6	SC3	D	1606	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 26222 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	728	Total	C	N	O	S	38	0	0
			5966	3825	986	1132	23			
1	B	728	Total	C	N	O	S	42	0	0
			5966	3825	986	1132	23			
1	C	728	Total	C	N	O	S	46	0	0
			5966	3825	986	1132	23			
1	D	728	Total	C	N	O	S	36	0	0
			5966	3825	986	1132	23			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	3	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	B	2	Total	C	N	O	0	0
			28	16	2	10		
3	C	2	Total	C	N	O	0	0
			28	16	2	10		
3	D	2	Total	C	N	O	0	0
			28	16	2	10		
3	D	2	Total	C	N	O	0	0
			28	16	2	10		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	3	Total	C	N	O	0	0
			39	22	2	15		
4	D	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 7-BENZYL-1,3-DIMETHYL-8-PIPERAZIN-1-YL-3,7-DIHYDRO-PURINE-2,6-DIONE (three-letter code: SC3) (formula: C₁₈H₂₃N₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			26	18	6	2		
6	B	1	Total	C	N	O	0	0
			26	18	6	2		
6	C	1	Total	C	N	O	0	0
			26	18	6	2		
6	D	1	Total	C	N	O	0	0
			26	18	6	2		
6	B	1	Total	C	N	O	0	0
			26	18	6	2		
6	D	1	Total	C	N	O	0	0
			26	18	6	2		
6	C	1	Total	C	N	O	0	0
			26	18	6	2		
6	A	1	Total	C	N	O	0	0
			26	18	6	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	394	Total	O	0	0
			394	394		
7	B	481	Total	O	0	0
			481	481		
7	C	395	Total	O	0	0
			395	395		
7	D	376	Total	O	0	0
			376	376		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dipeptidyl peptidase 4





• Molecule 1: Dipeptidyl peptidase 4







• Molecule 1: Dipeptidyl peptidase 4



• Molecule 1: Dipeptidyl peptidase 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	63.11Å 118.96Å 133.76Å 112.79° 95.74° 90.63°	Depositor
Resolution (Å)	39.45 – 2.11 39.45 – 2.11	Depositor EDS
% Data completeness (in resolution range)	97.7 (39.45-2.11) 85.1 (39.45-2.11)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.12Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.197 , 0.239 0.203 , 0.243	Depositor DCC
R_{free} test set	9993 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	27.2	Xtriage
Anisotropy	0.386	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 200628 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	26222	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, SC3, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/6141	0.73	2/8353 (0.0%)
1	B	0.59	0/6141	0.76	4/8353 (0.0%)
1	C	0.56	0/6141	0.74	1/8353 (0.0%)
1	D	0.53	0/6141	0.72	2/8353 (0.0%)
All	All	0.55	0/24564	0.74	9/33412 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	669	ARG	NE-CZ-NH2	-7.16	116.72	120.30
1	A	669	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	B	415	LEU	CA-CB-CG	5.79	128.62	115.30
1	A	669	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	B	669	ARG	NE-CZ-NH1	5.51	123.06	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	700	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5966	0	5663	124	0
1	B	5966	0	5663	99	0
1	C	5966	0	5663	110	0
1	D	5966	0	5662	116	0
2	A	56	0	52	1	0
2	B	56	0	52	1	0
2	C	70	0	65	1	0
2	D	28	0	26	1	0
3	A	56	0	50	1	0
3	B	28	0	25	0	0
3	C	28	0	25	0	0
3	D	84	0	75	0	0
4	B	39	0	34	0	0
4	D	39	0	34	1	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
5	C	5	0	0	0	0
5	D	5	0	0	0	0
6	A	52	0	46	2	0
6	B	52	0	46	1	0
6	C	52	0	46	1	0
6	D	52	0	46	1	0
7	A	394	0	0	10	0
7	B	481	0	0	8	0
7	C	395	0	0	9	0
7	D	376	0	0	6	0
All	All	26222	0	23273	433	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 433 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:536:LYS:HG2	1:C:537:SER:H	1.24	1.01

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:65:ASP:HB2	1:B:466:LYS:HD2	1.45	0.98
1:C:535:ASP:C	1:C:536:LYS:HD3	1.81	0.98
1:B:41:ARG:HD3	1:B:42:THR:O	1.64	0.97
1:C:74:ASN:HB3	1:C:92:ASN:HB2	1.55	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	726/728 (100%)	689 (95%)	35 (5%)	2 (0%)	46	44
1	B	726/728 (100%)	695 (96%)	28 (4%)	3 (0%)	39	36
1	C	726/728 (100%)	692 (95%)	30 (4%)	4 (1%)	30	24
1	D	726/728 (100%)	696 (96%)	28 (4%)	2 (0%)	46	44
All	All	2904/2912 (100%)	2772 (96%)	121 (4%)	11 (0%)	39	36

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	40	ARG
1	C	535	ASP
1	B	82	GLU
1	B	491	LEU
1	C	536	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	652/652 (100%)	627 (96%)	25 (4%)	40	39
1	B	652/652 (100%)	623 (96%)	29 (4%)	35	32
1	C	652/652 (100%)	614 (94%)	38 (6%)	25	21
1	D	652/652 (100%)	622 (95%)	30 (5%)	33	30
All	All	2608/2608 (100%)	2486 (95%)	122 (5%)	32	29

5 of 122 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	91	GLU
1	C	367	ASP
1	D	561	LEU
1	C	119	ASN
1	C	192	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 96 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	718	GLN
1	C	176	ASN
1	D	572	ASN
1	B	731	GLN
1	C	123	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	769(A)	1,3	14,14,15	0.60	0	15,19,21	0.69	1 (6%)
3	NAG	A	770(B)	3	14,14,15	0.49	0	15,19,21	0.85	1 (6%)
3	NAG	A	773(A)	1,3	14,14,15	0.61	0	15,19,21	0.78	1 (6%)
3	NAG	A	774(B)	3	14,14,15	0.64	0	15,19,21	0.82	1 (6%)
4	NAG	B	768(A)	1,4	14,14,15	0.61	0	15,19,21	1.02	2 (13%)
4	NAG	B	769(B)	4	14,14,15	0.66	0	15,19,21	0.77	0
4	BMA	B	770(C)	4	11,11,12	0.60	0	14,15,17	1.37	3 (21%)
3	NAG	B	774(A)	1,3	14,14,15	0.63	0	15,19,21	0.93	1 (6%)
3	NAG	B	775(B)	3	14,14,15	0.69	0	15,19,21	0.92	1 (6%)
3	NAG	C	772(A)	1,3	14,14,15	0.59	0	15,19,21	0.78	1 (6%)
3	NAG	C	773(B)	3	14,14,15	0.59	0	15,19,21	1.00	2 (13%)
4	NAG	D	768(A)	1,4	14,14,15	0.57	0	15,19,21	0.73	0
4	NAG	D	769(B)	4	14,14,15	0.56	0	15,19,21	0.72	0
4	BMA	D	770(C)	4	11,11,12	0.62	0	14,15,17	0.70	1 (7%)
3	NAG	D	771(A)	1,3	14,14,15	0.56	0	15,19,21	0.67	1 (6%)
3	NAG	D	772(B)	3	14,14,15	0.49	0	15,19,21	0.66	0
3	NAG	D	774(A)	1,3	14,14,15	0.55	0	15,19,21	1.04	2 (13%)
3	NAG	D	775(B)	3	14,14,15	0.66	0	15,19,21	0.85	1 (6%)
3	NAG	D	776(A)	1,3	14,14,15	0.72	0	15,19,21	0.65	0
3	NAG	D	777(B)	3	14,14,15	0.59	0	15,19,21	1.13	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	769(A)	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	770(B)	3	-	0/6/23/26	0/1/1/1
3	NAG	A	773(A)	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	774(B)	3	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	768(A)	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	769(B)	4	-	0/6/23/26	0/1/1/1
4	BMA	B	770(C)	4	-	0/2/19/22	0/1/1/1
3	NAG	B	774(A)	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	775(B)	3	-	0/6/23/26	0/1/1/1
3	NAG	C	772(A)	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	773(B)	3	-	0/6/23/26	0/1/1/1
4	NAG	D	768(A)	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	769(B)	4	-	0/6/23/26	0/1/1/1
4	BMA	D	770(C)	4	-	0/2/19/22	0/1/1/1
3	NAG	D	771(A)	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	772(B)	3	-	0/6/23/26	0/1/1/1
3	NAG	D	774(A)	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	775(B)	3	-	0/6/23/26	0/1/1/1
3	NAG	D	776(A)	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	777(B)	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	774(A)	NAG	C2-N2-C7	-2.84	119.38	123.04
4	B	770(C)	BMA	C6-C5-C4	-2.84	106.02	113.02
3	A	770(B)	NAG	C2-N2-C7	-2.62	119.67	123.04
3	D	774(A)	NAG	C2-N2-C7	-2.47	119.87	123.04
3	B	775(B)	NAG	C2-N2-C7	-2.44	119.91	123.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	774(B)	NAG	1	0
4	D	768(A)	NAG	1	0
4	D	769(B)	NAG	1	0

5.6 Ligand geometry ⓘ

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	SO4	A	1500	-	4,4,4	0.63	0	6,6,6	0.58	0
6	SC3	A	1601	-	22,29,29	3.14	6 (27%)	21,42,42	1.56	2 (9%)
6	SC3	A	1608	-	22,29,29	2.83	9 (40%)	21,42,42	1.58	2 (9%)
2	NAG	A	767(A)	1	14,14,15	0.65	0	15,19,21	0.78	1 (6%)
2	NAG	A	768(A)	1	14,14,15	0.58	0	15,19,21	0.78	1 (6%)
2	NAG	A	771(A)	1	14,14,15	0.47	0	15,19,21	0.86	1 (6%)
2	NAG	A	772(A)	1	14,14,15	0.48	0	15,19,21	1.24	3 (20%)
5	SO4	B	1501	-	4,4,4	0.73	0	6,6,6	0.73	0
6	SC3	B	1602	-	22,29,29	3.28	8 (36%)	21,42,42	1.56	2 (9%)
6	SC3	B	1605	-	22,29,29	3.28	10 (45%)	21,42,42	1.40	3 (14%)
2	NAG	B	767(A)	1	14,14,15	0.62	0	15,19,21	0.65	0
2	NAG	B	771(A)	1	14,14,15	0.45	0	15,19,21	0.78	0
2	NAG	B	772(A)	1	14,14,15	0.63	0	15,19,21	0.66	0
2	NAG	B	773(A)	1	14,14,15	0.59	0	15,19,21	0.92	1 (6%)
5	SO4	C	1502	-	4,4,4	0.73	0	6,6,6	0.72	0
6	SC3	C	1603	-	22,29,29	3.18	7 (31%)	21,42,42	1.50	2 (9%)
6	SC3	C	1607	-	22,29,29	3.18	8 (36%)	21,42,42	1.45	2 (9%)
2	NAG	C	767(A)	1	14,14,15	0.49	0	15,19,21	0.75	0
2	NAG	C	768(A)	1	14,14,15	0.50	0	15,19,21	0.82	1 (6%)
2	NAG	C	769(A)	1	14,14,15	0.52	0	15,19,21	0.79	0
2	NAG	C	770(A)	1	14,14,15	0.63	0	15,19,21	0.67	0
2	NAG	C	771(A)	1	14,14,15	0.54	0	15,19,21	0.82	0
5	SO4	D	1503	-	4,4,4	0.45	0	6,6,6	0.70	0
6	SC3	D	1604	-	22,29,29	3.30	6 (27%)	21,42,42	1.52	2 (9%)
6	SC3	D	1606	-	22,29,29	3.28	10 (45%)	21,42,42	1.48	2 (9%)
2	NAG	D	767(A)	1	14,14,15	0.84	1 (7%)	15,19,21	0.80	1 (6%)
2	NAG	D	773(A)	1	14,14,15	0.70	0	15,19,21	1.07	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SO4	A	1500	-	-	0/0/0/0	0/0/0/0
6	SC3	A	1601	-	-	0/4/16/16	0/4/4/4
6	SC3	A	1608	-	-	0/4/16/16	1/4/4/4
2	NAG	A	767(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	A	768(A)	1	-	1/6/23/26	0/1/1/1
2	NAG	A	771(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	A	772(A)	1	-	0/6/23/26	0/1/1/1
5	SO4	B	1501	-	-	0/0/0/0	0/0/0/0
6	SC3	B	1602	-	-	0/4/16/16	0/4/4/4
6	SC3	B	1605	-	-	0/4/16/16	0/4/4/4
2	NAG	B	767(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	B	771(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	B	772(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	B	773(A)	1	-	0/6/23/26	0/1/1/1
5	SO4	C	1502	-	-	0/0/0/0	0/0/0/0
6	SC3	C	1603	-	-	0/4/16/16	0/4/4/4
6	SC3	C	1607	-	-	0/4/16/16	1/4/4/4
2	NAG	C	767(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	C	768(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	C	769(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	C	770(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	C	771(A)	1	-	0/6/23/26	0/1/1/1
5	SO4	D	1503	-	-	0/0/0/0	0/0/0/0
6	SC3	D	1604	-	-	0/4/16/16	0/4/4/4
6	SC3	D	1606	-	-	0/4/16/16	1/4/4/4
2	NAG	D	767(A)	1	-	0/6/23/26	0/1/1/1
2	NAG	D	773(A)	1	-	0/6/23/26	0/1/1/1

The worst 5 of 65 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	1606	SC3	C8-N9	-5.46	1.26	1.34
6	C	1607	SC3	C8-N9	-4.76	1.27	1.34
6	B	1605	SC3	C8-N9	-4.56	1.27	1.34
6	D	1604	SC3	C8-N9	-4.08	1.28	1.34
6	A	1608	SC3	C8-N9	-4.05	1.28	1.34

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1608	SC3	C6-C5-C4	-5.29	116.14	119.93
6	D	1606	SC3	C6-C5-C4	-4.78	116.50	119.93
6	A	1601	SC3	C6-C5-C4	-4.67	116.59	119.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1602	SC3	C6-C5-C4	-4.66	116.59	119.93
6	C	1603	SC3	C6-C5-C4	-4.41	116.77	119.93

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	768(A)	NAG	O7-C7-N2-C2

All (3) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	1607	SC3	C11-C12-C14-C15-N10-N13
6	D	1606	SC3	C11-C12-C14-C15-N10-N13
6	A	1608	SC3	C11-C12-C14-C15-N10-N13

8 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1601	SC3	2	0
2	A	768(A)	NAG	1	0
6	B	1602	SC3	1	0
2	B	772(A)	NAG	1	0
6	C	1603	SC3	1	0
2	C	768(A)	NAG	1	0
6	D	1604	SC3	1	0
2	D	773(A)	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	728/728 (100%)	0.08	23 (3%) 51 60	15, 30, 59, 87	11 (1%)
1	B	728/728 (100%)	-0.17	9 (1%) 81 85	13, 26, 49, 72	12 (1%)
1	C	728/728 (100%)	-0.02	27 (3%) 45 54	15, 29, 60, 99	12 (1%)
1	D	728/728 (100%)	0.03	21 (2%) 55 63	16, 31, 57, 75	10 (1%)
All	All	2912/2912 (100%)	-0.02	80 (2%) 58 65	13, 29, 56, 99	45 (1%)

The worst 5 of 80 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	100	TYR	8.1
1	C	39	SER	6.5
1	C	95	PHE	6.1
1	A	83	TYR	6.0
1	C	101	SER	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	C	772(A)	14/15	0.95	0.13	2.38	31,35,40,40	0
3	NAG	D	774(A)	14/15	0.84	0.18	1.37	55,61,65,66	0
3	NAG	D	771(A)	14/15	0.92	0.11	0.11	32,37,46,50	0
3	NAG	A	773(A)	14/15	0.93	0.11	0.09	29,37,44,46	0
3	NAG	A	769(A)	14/15	0.93	0.10	0.06	37,42,46,51	0
3	NAG	D	776(A)	14/15	0.93	0.10	-0.21	28,35,43,44	0
4	NAG	D	768(A)	14/15	0.83	0.15	-0.69	71,73,74,76	0
4	NAG	B	768(A)	14/15	0.91	0.12	-0.72	42,47,51,51	0
3	NAG	B	774(A)	14/15	0.96	0.07	-1.07	26,34,41,42	0
3	NAG	D	772(B)	14/15	0.77	0.27	-	58,62,66,66	0
4	NAG	D	769(B)	14/15	0.79	0.29	-	76,77,81,82	0
3	NAG	A	774(B)	14/15	0.90	0.22	-	44,49,53,57	0
3	NAG	D	775(B)	14/15	0.77	0.31	-	66,68,69,69	0
4	BMA	D	770(C)	11/12	0.72	0.30	-	76,77,77,77	0
3	NAG	D	777(B)	14/15	0.88	0.16	-	47,52,55,56	0
3	NAG	C	773(B)	14/15	0.89	0.19	-	45,48,51,51	0
4	BMA	B	770(C)	11/12	0.80	0.15	-	50,52,54,54	0
3	NAG	B	775(B)	14/15	0.89	0.20	-	37,45,50,56	0
3	NAG	A	770(B)	14/15	0.83	0.26	-	58,63,65,65	0
4	NAG	B	769(B)	14/15	0.92	0.12	-	45,46,48,48	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	SC3	D	1606	26/26	0.94	0.12	4.02	26,30,42,43	0
6	SC3	A	1601	26/26	0.94	0.19	1.35	22,28,31,33	0
6	SC3	C	1603	26/26	0.94	0.17	1.13	24,30,35,36	0
6	SC3	D	1604	26/26	0.94	0.17	1.12	24,31,33,34	0
2	NAG	B	773(A)	14/15	0.91	0.11	0.87	36,40,45,45	0
2	NAG	B	767(A)	14/15	0.72	0.24	0.81	71,73,74,74	0
2	NAG	A	767(A)	14/15	0.72	0.29	0.77	80,81,83,83	0
2	NAG	C	771(A)	14/15	0.88	0.13	0.64	42,48,50,52	0
6	SC3	B	1602	26/26	0.94	0.15	0.62	21,26,30,32	0
2	NAG	B	771(A)	14/15	0.90	0.12	0.57	29,33,40,40	0
6	SC3	B	1605	26/26	0.95	0.11	0.41	26,28,39,40	0
2	NAG	C	767(A)	14/15	0.87	0.14	0.32	54,57,60,60	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	B	772(A)	14/15	0.88	0.11	0.23	39,41,44,47	0
6	SC3	A	1608	26/26	0.95	0.11	0.04	26,35,42,45	0
2	NAG	C	769(A)	14/15	0.94	0.10	-0.04	32,36,39,40	0
2	NAG	A	772(A)	14/15	0.91	0.12	-0.41	44,48,50,52	0
5	SO4	D	1503	5/5	0.98	0.09	-0.62	38,40,42,43	0
5	SO4	C	1502	5/5	0.99	0.09	-0.81	28,28,31,31	0
2	NAG	A	768(A)	14/15	0.75	0.15	-0.91	73,74,74,75	0
6	SC3	C	1607	26/26	0.97	0.09	-0.97	30,34,39,40	0
5	SO4	A	1500	5/5	0.98	0.10	-1.31	33,33,34,34	0
5	SO4	B	1501	5/5	1.00	0.07	-2.34	26,26,27,27	0
2	NAG	C	770(A)	14/15	0.80	0.14	-	40,41,45,45	0
2	NAG	C	768(A)	14/15	0.59	0.24	-	85,88,89,89	0
2	NAG	D	773(A)	14/15	0.68	0.23	-	60,64,67,68	0
2	NAG	A	771(A)	14/15	0.84	0.12	-	51,53,55,58	3
2	NAG	D	767(A)	14/15	0.32	0.45	-	79,81,84,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.