



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:23 AM GMT

PDB ID : 3ALM
Title : Crystal structure of 2-methyl-3-hydroxypyridine-5-carboxylic acid oxygenase, mutant C294A
Authors : Kobayashi, J.; Yoshida, H.; Yoshikane, Y.; Kamitori, S.; Yagi, T.
Deposited on : 2010-08-04
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6655 (1.80-1.76)
Clashscore	102246	7658 (1.80-1.76)
Ramachandran outliers	100387	7570 (1.80-1.76)
Sidechain outliers	100360	7569 (1.80-1.76)
RSRZ outliers	91569	6671 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 89%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 89% 9% </div> </div>
1	B	379	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 11%, green 86%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 86% 11% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	381	-	-	-	X
3	GOL	B	381	-	-	-	X
3	GOL	B	382	-	-	-	X
3	GOL	B	384	-	-	X	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6232 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-methyl-3-hydroxypyridine-5-carboxylic acid oxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	370	Total	C	N	O	S	0	3	0
			2888	1823	516	536	13			
1	B	370	Total	C	N	O	S	0	1	0
			2877	1815	514	535	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	294	ALA	CYS	ENGINEERED MUTATION	UNP Q988D3
B	294	ALA	CYS	ENGINEERED MUTATION	UNP Q988D3

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	160	Total	O	0	0
			160	160		
4	B	171	Total	O	0	0
			171	171		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-methyl-3-hydroxypyridine-5-carboxylic acid oxygenase



- Molecule 1: 2-methyl-3-hydroxypyridine-5-carboxylic acid oxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.90 Å 129.80 Å 88.97 Å 90.00° 122.18° 90.00°	Depositor
Resolution (Å)	35.42 – 1.77 35.42 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.2 (35.42-1.77) 99.2 (35.42-1.77)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.16 (at 1.77 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.193 , 0.228 0.193 , 0.226	Depositor DCC
R_{free} test set	5076 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 101640 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6232	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	5/2963 (0.2%)	1.11	13/4021 (0.3%)
1	B	1.36	9/2946 (0.3%)	1.10	12/3999 (0.3%)
All	All	1.35	14/5909 (0.2%)	1.10	25/8020 (0.3%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	259	ALA	CA-CB	7.01	1.67	1.52
1	B	271	GLU	CB-CG	-6.96	1.39	1.52
1	A	203	TRP	CE3-CZ3	6.88	1.50	1.38
1	B	114	VAL	CB-CG1	6.57	1.66	1.52
1	B	223	TYR	CD1-CE1	6.06	1.48	1.39
1	A	195	GLU	CB-CG	5.97	1.63	1.52
1	A	290	ALA	C-O	5.73	1.34	1.23
1	A	259	ALA	CA-CB	5.42	1.63	1.52
1	A	114	VAL	CB-CG1	5.29	1.64	1.52
1	B	290	ALA	C-O	5.28	1.33	1.23
1	B	203	TRP	CE3-CZ3	5.25	1.47	1.38
1	B	83	GLU	CD-OE1	5.17	1.31	1.25
1	B	270	TYR	CG-CD1	5.07	1.45	1.39
1	B	196	TRP	CE3-CZ3	5.04	1.47	1.38

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	276	ASP	CB-CG-OD1	12.49	129.54	118.30
1	A	276	ASP	CB-CG-OD1	11.18	128.36	118.30
1	A	333	ARG	NE-CZ-NH2	-10.23	115.19	120.30
1	B	276	ASP	CB-CG-OD2	-8.32	110.81	118.30
1	B	333	ARG	NE-CZ-NH2	-8.16	116.22	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	61	ARG	NE-CZ-NH2	-7.54	116.53	120.30
1	B	226	LEU	CB-CG-CD2	7.38	123.55	111.00
1	A	276	ASP	CB-CG-OD2	-7.18	111.83	118.30
1	A	99	LEU	CB-CG-CD2	-6.71	99.59	111.00
1	B	134	ASP	CB-CG-OD2	-6.61	112.35	118.30
1	B	275	LEU	CB-CG-CD1	-6.56	99.85	111.00
1	B	208	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	367	ARG	NE-CZ-NH2	-5.73	117.43	120.30
1	A	338	ARG	NE-CZ-NH1	-5.73	117.44	120.30
1	B	12	ARG	NE-CZ-NH2	5.61	123.11	120.30
1	B	69	LEU	CB-CG-CD1	5.54	120.41	111.00
1	A	323	ASP	CB-CG-OD1	5.44	123.20	118.30
1	B	78	THR	CA-CB-CG2	-5.43	104.80	112.40
1	B	134	ASP	CB-CG-OD1	5.40	123.16	118.30
1	B	61	ARG	NE-CZ-NH2	-5.40	117.60	120.30
1	A	333	ARG	NH1-CZ-NH2	5.26	125.18	119.40
1	A	118	ARG	NE-CZ-NH1	-5.24	117.68	120.30
1	A	78	THR	CA-CB-CG2	-5.18	105.14	112.40
1	A	372	ARG	NE-CZ-NH2	-5.15	117.72	120.30
1	A	12	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2888	0	2861	15	0
1	B	2877	0	2842	17	0
2	A	53	0	31	1	0
2	B	53	0	31	2	0
3	A	6	0	8	1	0
3	B	24	0	32	6	0
4	A	160	0	0	1	0
4	B	171	0	0	2	0
All	All	6232	0	5805	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:173:TRP:CZ3	3:B:384:GOL:H11	1.96	1.00
1:A:170:GLN:HE22	1:B:172:ARG:HH11	1.27	0.82
1:A:87:HIS:CE1	1:A:205:PHE:H	2.01	0.78
1:A:295:PRO:HG2	3:A:381:GOL:H12	1.72	0.72
1:B:15:GLU:OE1	1:B:40:HIS:HE1	1.72	0.72
1:A:87:HIS:HE1	1:A:206:TRP:H	1.38	0.71
1:A:172:ARG:HH11	1:B:170:GLN:HE22	1.38	0.71
1:A:87:HIS:HE1	1:A:205:PHE:H	1.38	0.70
1:A:87:HIS:CE1	1:A:206:TRP:H	2.17	0.61
1:B:44:SER:HB3	1:B:127:ASN:HD21	1.67	0.60
1:A:137:GLY:C	1:A:138:ARG:HG2	2.22	0.58
1:B:266:ARG:NH2	1:B:268:ASP:OD1	2.34	0.58
1:A:138:ARG:N	1:A:152:ILE:HD11	2.19	0.57
1:B:40:HIS:HD2	1:B:128:SER:OG	1.88	0.56
1:B:173:TRP:CE3	3:B:384:GOL:H11	2.41	0.54
1:B:173:TRP:CZ3	3:B:384:GOL:C1	2.84	0.53
1:B:42:LYS:HE2	2:B:380:FAD:C8A	2.40	0.52
1:A:87:HIS:HE1	1:A:205:PHE:N	2.07	0.51
1:B:294:ALA:HB2	3:B:384:GOL:O1	2.11	0.51
1:B:142:GLN:NE2	3:B:381:GOL:O3	2.43	0.51
4:A:455:HOH:O	1:B:269:LYS:HD3	2.11	0.49
3:B:382:GOL:H31	4:B:549:HOH:O	2.13	0.49
1:A:97[B]:ASN:HD22	1:A:379:GLN:NE2	2.10	0.49
1:A:300:GLY:HA3	2:A:380:FAD:H1'2	1.95	0.48
1:B:152:ILE:HB	1:B:283:VAL:HG22	1.98	0.45
1:A:342:LEU:HD23	1:A:362:ALA:HA	1.98	0.44
1:A:137:GLY:C	1:A:152:ILE:HD11	2.40	0.42
1:A:269:LYS:HD3	4:B:443:HOH:O	2.19	0.42
1:B:300:GLY:HA3	2:B:380:FAD:H1'2	2.00	0.42
1:B:238:VAL:HA	1:B:239:PRO:C	2.40	0.42
1:B:183:ILE:HD13	1:B:223:TYR:HA	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/379 (98%)	360 (97%)	11 (3%)	0	100	100
1	B	369/379 (97%)	357 (97%)	11 (3%)	1 (0%)	46	28
All	All	740/758 (98%)	717 (97%)	22 (3%)	1 (0%)	56	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	160	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/304 (99%)	296 (99%)	4 (1%)	76	66
1	B	298/304 (98%)	292 (98%)	6 (2%)	63	47
All	All	598/608 (98%)	588 (98%)	10 (2%)	68	54

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LYS
1	A	82	TYR
1	A	138	ARG
1	A	142	GLN
1	B	10	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	45	GLU
1	B	82	TYR
1	B	142	GLN
1	B	223	TYR
1	B	266	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	87	HIS
1	A	115	ASN
1	A	127	ASN
1	A	142	GLN
1	A	170	GLN
1	A	198	ASN
1	A	307	ASN
1	A	379	GLN
1	B	33	ASN
1	B	40	HIS
1	B	97	ASN
1	B	127	ASN
1	B	142	GLN
1	B	170	GLN
1	B	198	ASN
1	B	307	ASN
1	B	379	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	380	-	48,58,58	1.25	6 (12%)	54,89,89	2.35	11 (20%)
3	GOL	A	381	-	5,5,5	0.71	0	5,5,5	1.21	0
2	FAD	B	380	-	48,58,58	1.67	7 (14%)	54,89,89	2.19	10 (18%)
3	GOL	B	381	-	5,5,5	0.50	0	5,5,5	0.89	0
3	GOL	B	382	-	5,5,5	0.56	0	5,5,5	1.01	0
3	GOL	B	383	-	5,5,5	0.44	0	5,5,5	0.71	0
3	GOL	B	384	-	5,5,5	0.77	0	5,5,5	1.35	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	380	-	-	0/30/50/50	0/6/6/6
3	GOL	A	381	-	-	0/4/4/4	0/0/0/0
2	FAD	B	380	-	-	0/30/50/50	0/6/6/6
3	GOL	B	381	-	-	0/4/4/4	0/0/0/0
3	GOL	B	382	-	-	0/4/4/4	0/0/0/0
3	GOL	B	383	-	-	0/4/4/4	0/0/0/0
3	GOL	B	384	-	-	0/4/4/4	0/0/0/0

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	380	FAD	C5X-N5	2.00	1.38	1.35
2	A	380	FAD	C7M-C7	2.02	1.55	1.51
2	A	380	FAD	PA-O5B	2.04	1.68	1.59
2	B	380	FAD	C4-N3	2.09	1.37	1.33
2	B	380	FAD	C5X-N5	2.11	1.38	1.35
2	B	380	FAD	C10-N1	2.19	1.39	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	380	FAD	C9A-N10	2.29	1.41	1.38
2	B	380	FAD	C8M-C8	2.60	1.56	1.51
2	B	380	FAD	C4X-N5	2.97	1.38	1.33
2	A	380	FAD	O4B-C1B	3.16	1.45	1.41
2	A	380	FAD	C8M-C8	3.21	1.57	1.51
2	B	380	FAD	C9A-N10	3.27	1.43	1.38
2	B	380	FAD	O4B-C1B	6.97	1.50	1.41

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	380	FAD	N3A-C2A-N1A	-13.05	118.90	128.89
2	B	380	FAD	N3A-C2A-N1A	-9.95	121.28	128.89
2	B	380	FAD	O4B-C1B-N9A	-4.44	98.80	108.10
2	B	380	FAD	P-O3P-PA	-3.43	123.10	132.73
2	A	380	FAD	C4B-O4B-C1B	-2.72	106.73	109.72
2	B	380	FAD	C4X-C4-N3	-2.44	120.26	123.59
2	A	380	FAD	O4B-C4B-C5B	-2.39	100.78	109.32
2	B	380	FAD	O4B-C4B-C5B	-2.39	100.79	109.32
2	A	380	FAD	C1B-N9A-C4A	-2.34	123.41	126.94
2	B	380	FAD	C4-C4X-C10	-2.33	118.45	119.94
2	A	380	FAD	C9A-C5X-N5	-2.11	119.23	122.36
2	A	380	FAD	C8M-C8-C9	-2.09	114.59	120.28
2	B	380	FAD	C2A-N1A-C6A	2.32	122.92	118.77
3	B	384	GOL	C3-C2-C1	2.54	121.08	111.12
2	A	380	FAD	O3P-P-O5'	2.58	109.78	102.94
2	A	380	FAD	C4X-N5-C5X	2.83	120.02	116.76
2	A	380	FAD	C4-N3-C2	3.37	118.16	115.25
2	B	380	FAD	C1'-N10-C9A	3.49	122.78	118.86
2	A	380	FAD	C1'-N10-C9A	3.69	123.01	118.86
2	A	380	FAD	C5X-C9A-N10	4.20	120.81	117.62
2	B	380	FAD	C5X-C9A-N10	4.30	120.88	117.62
2	B	380	FAD	C4-N3-C2	5.57	120.06	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	380	FAD	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	381	GOL	1	0
2	B	380	FAD	2	0
3	B	381	GOL	1	0
3	B	382	GOL	1	0
3	B	384	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/379 (97%)	-0.24	5 (1%) 78 77	12, 19, 31, 54	0
1	B	370/379 (97%)	-0.26	4 (1%) 82 82	13, 19, 32, 56	0
All	All	740/758 (97%)	-0.25	9 (1%) 81 81	12, 19, 32, 56	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	10	LYS	4.6
1	A	10	LYS	4.2
1	A	11	THR	3.6
1	B	11	THR	2.9
1	B	318	GLY	2.8
1	B	303	CYS	2.6
1	A	303	CYS	2.4
1	A	143	THR	2.3
1	A	318	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	B	384	6/6	0.86	0.28	9.71	32,46,52,52	0
3	GOL	A	381	6/6	0.81	0.16	4.87	35,45,45,49	0
3	GOL	B	381	6/6	0.81	0.28	4.32	44,47,48,50	0
3	GOL	B	382	6/6	0.86	0.19	2.73	33,46,49,50	0
3	GOL	B	383	6/6	0.93	0.11	1.72	26,40,43,49	0
2	FAD	B	380	53/53	0.96	0.08	-0.52	13,17,21,22	0
2	FAD	A	380	53/53	0.96	0.07	-0.86	14,18,21,23	0

6.5 Other polymers

There are no such residues in this entry.