



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 29, 2016 – 12:53 PM EDT

PDB ID : 5AOD
Title : Crystal structure of wild type pneumolysin.
Authors : van Pee, K.; Yildiz, O.
Deposited on : 2015-09-10
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3777 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PNEUMOLYSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	474	Total	C	N	O	S	0	0	0
			3751	2364	637	743	7			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP Q04IN8
A	-1	SER	-	EXPRESSION TAG	UNP Q04IN8
A	0	HIS	-	EXPRESSION TAG	UNP Q04IN8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	26	Total	O	0	0
			26	26		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PNEUMOLYSIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	24.28Å 113.26Å 156.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.17 – 2.40 39.17 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.2 (39.17-2.40) 99.2 (39.17-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 2.39Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.218 , 0.277 0.217 , 0.276	Depositor DCC
R_{free} test set	888 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	60.9	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 63.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3777	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/3825	0.59	3/5198 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	450	LYS	CA-CB-CG	5.37	125.21	113.40
1	A	469	GLU	OE1-CD-OE2	-5.20	117.06	123.30
1	A	158	MET	CG-SD-CE	5.06	108.29	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3751	0	3692	85	1
2	A	26	0	0	2	0
All	All	3777	0	3692	85	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:HIS:HB2	1:A:322:ALA:HB3	1.39	1.02
1:A:450:LYS:HD2	1:A:469:GLU:CD	1.94	0.88
1:A:73:GLY:O	1:A:94:ARG:NH2	2.11	0.83
1:A:301:ARG:HH21	1:A:303:VAL:HG11	1.46	0.80
1:A:450:LYS:HD2	1:A:469:GLU:CG	2.19	0.72
1:A:469:GLU:O	1:A:469:GLU:HG2	1.88	0.72
1:A:446:PRO:O	1:A:451:ARG:NH2	2.22	0.71
1:A:152:LYS:NZ	1:A:264:GLU:OE2	2.24	0.69
1:A:450:LYS:HD2	1:A:469:GLU:HG3	1.76	0.68
1:A:150:TYR:CE2	1:A:195:PHE:CE1	2.82	0.68
1:A:28:ILE:HD11	1:A:50:LYS:HD2	1.76	0.67
1:A:181:ASN:OD1	1:A:182:SER:N	2.29	0.66
1:A:450:LYS:N	1:A:469:GLU:OE2	2.28	0.66
1:A:450:LYS:CE	1:A:469:GLU:HG3	2.29	0.63
1:A:450:LYS:CD	1:A:469:GLU:HG3	2.30	0.62
1:A:469:GLU:OE1	1:A:469:GLU:N	2.33	0.62
1:A:295:ASP:HB3	1:A:296:PRO:HD2	1.83	0.59
1:A:150:TYR:CZ	1:A:195:PHE:HE1	2.20	0.59
1:A:469:GLU:O	1:A:469:GLU:CG	2.51	0.59
1:A:450:LYS:HE3	1:A:469:GLU:HG3	1.85	0.57
1:A:175:SER:O	1:A:288:LYS:NZ	2.34	0.57
1:A:62:VAL:O	2:A:2008:HOH:O	2.18	0.57
1:A:62:VAL:HG21	1:A:70:LEU:HD13	1.87	0.56
1:A:277:GLU:O	1:A:281:ILE:HG23	2.08	0.54
1:A:362:ASP:N	1:A:449:ARG:O	2.29	0.54
1:A:150:TYR:CE2	1:A:195:PHE:HE1	2.24	0.54
1:A:319:ARG:HE	1:A:321:THR:HG22	1.72	0.54
1:A:29:GLU:O	1:A:31:ARG:HG2	2.09	0.53
1:A:215:GLN:NE2	2:A:2013:HOH:O	2.38	0.53
1:A:367:HIS:HB3	1:A:408:PHE:HB3	1.90	0.53
1:A:448:VAL:HG21	1:A:468:VAL:HG13	1.91	0.53
1:A:258:GLU:HB3	1:A:261:ALA:HB3	1.90	0.52
1:A:362:ASP:O	1:A:450:LYS:HA	2.09	0.52
1:A:189:GLN:HE21	1:A:255:LYS:HA	1.74	0.52
1:A:255:LYS:HB2	1:A:357:ALA:HB2	1.92	0.52
1:A:106:LEU:HD21	1:A:134:TRP:HD1	1.74	0.52
1:A:292:LEU:HB3	1:A:327:LEU:HD21	1.92	0.52
1:A:259:VAL:HG22	1:A:281:ILE:HD12	1.93	0.51
1:A:246:VAL:HG22	1:A:291:ILE:HG13	1.93	0.51
1:A:277:GLU:HG3	1:A:280:GLN:OE1	2.12	0.50
1:A:39:LEU:HB3	1:A:40:PRO:HD2	1.94	0.50
1:A:32:PHE:HD1	1:A:33:ILE:N	2.11	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:ILE:O	1:A:50:LYS:NZ	2.46	0.48
1:A:368:SER:HB3	1:A:407:HIS:CD2	2.49	0.48
1:A:189:GLN:NE2	1:A:255:LYS:HA	2.29	0.48
1:A:124:ARG:NH2	1:A:328:PRO:HD3	2.29	0.47
1:A:53:LEU:O	1:A:347:SER:HA	2.14	0.47
1:A:104:PRO:HG2	1:A:144:VAL:HG11	1.96	0.47
1:A:108:SER:O	1:A:111:SER:OG	2.26	0.47
1:A:62:VAL:HG13	1:A:64:ALA:H	1.80	0.47
1:A:432:ALA:O	1:A:433:TRP:HB3	2.16	0.46
1:A:47:GLU:O	1:A:353:THR:HA	2.15	0.46
1:A:158:MET:O	1:A:162:LYS:HG3	2.15	0.46
1:A:450:LYS:HD2	1:A:469:GLU:OE1	2.14	0.46
1:A:16:ASP:OD1	1:A:19:LYS:HG2	2.17	0.45
1:A:249:LYS:NZ	1:A:251:GLU:OE1	2.30	0.45
1:A:150:TYR:CE2	1:A:195:PHE:CZ	3.05	0.45
1:A:47:GLU:HB3	1:A:49:LYS:HZ2	1.82	0.44
1:A:262:ALA:HB3	1:A:281:ILE:HD11	1.99	0.44
1:A:263:PHE:O	1:A:267:ILE:HG12	2.18	0.44
1:A:439:VAL:CG1	1:A:464:VAL:HG22	2.48	0.44
1:A:58:SER:HB3	1:A:343:THR:HG22	2.00	0.44
1:A:23:HIS:HB3	1:A:346:ASN:HD22	1.83	0.43
1:A:266:LEU:HD21	1:A:310:VAL:HG21	2.00	0.43
1:A:385:ASP:OD2	1:A:389:LYS:HE3	2.18	0.43
1:A:99:TYR:CD2	1:A:204:VAL:HG12	2.53	0.43
1:A:375:TYR:CD1	1:A:404:LEU:HD12	2.54	0.42
1:A:47:GLU:HB3	1:A:49:LYS:NZ	2.35	0.42
1:A:103:LEU:HB2	1:A:106:LEU:HD11	2.01	0.42
1:A:106:LEU:HD21	1:A:134:TRP:CD1	2.54	0.42
1:A:282:LEU:HD23	1:A:282:LEU:HA	1.91	0.42
1:A:194:ASN:OD1	1:A:196:LYS:HE2	2.20	0.42
1:A:290:VAL:HG22	1:A:302:VAL:HG22	2.02	0.42
1:A:195:PHE:O	1:A:245:GLN:HA	2.20	0.41
1:A:273:ALA:HA	1:A:274:PRO:HD3	1.93	0.41
1:A:229:SER:OG	1:A:231:GLU:HG2	2.19	0.41
1:A:104:PRO:HG2	1:A:144:VAL:CG1	2.50	0.41
1:A:199:TYR:CD2	1:A:243:GLY:HA2	2.55	0.41
1:A:272:VAL:HG13	1:A:279:LYS:NZ	2.35	0.41
1:A:51:ARG:HB2	1:A:350:TYR:CZ	2.55	0.41
1:A:153:ILE:HD12	1:A:153:ILE:HA	1.92	0.41
1:A:368:SER:HB3	1:A:407:HIS:HD2	1.85	0.41
1:A:72:PRO:HB2	1:A:99:TYR:CE2	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:ALA:HB3	1:A:237:ILE:HD12	2.03	0.40
1:A:255:LYS:HG3	1:A:357:ALA:HB2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:SER:OG	1:A:275:GLN:NE2[3_555]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/474 (100%)	452 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	418/418 (100%)	402 (96%)	16 (4%)	40	60

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	32	PHE
1	A	130	LEU
1	A	157	SER
1	A	160	GLN
1	A	167	SER
1	A	182	SER
1	A	203	SER
1	A	284	ASN
1	A	319	ARG
1	A	324	HIS
1	A	359	ARG
1	A	403	ASP
1	A	410	THR
1	A	450	LYS
1	A	471	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	149	GLN
1	A	189	GLN
1	A	245	GLN
1	A	346	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	474/474 (100%)	0.40	34 (7%)	18 18	41, 72, 119, 181	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	433	TRP	8.9
1	A	386	HIS	7.7
1	A	388	GLY	5.5
1	A	33	ILE	4.4
1	A	105	GLY	4.3
1	A	46	ILE	4.1
1	A	471	ASP	4.1
1	A	30	ASN	4.1
1	A	195	PHE	3.8
1	A	297	SER	3.7
1	A	294	GLY	3.5
1	A	355	VAL	3.3
1	A	28	ILE	3.1
1	A	108	SER	3.1
1	A	469	GLU	3.1
1	A	141	VAL	2.9
1	A	47	GLU	2.9
1	A	107	ALA	2.9
1	A	91	ALA	2.8
1	A	296	PRO	2.7
1	A	450	LYS	2.7
1	A	31	ARG	2.6
1	A	150	TYR	2.6
1	A	389	LYS	2.5
1	A	356	THR	2.5
1	A	188	LYS	2.4
1	A	323	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	318	SER	2.3
1	A	136	GLN	2.3
1	A	324	HIS	2.2
1	A	435	TRP	2.1
1	A	135	HIS	2.1
1	A	295	ASP	2.0
1	A	66	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.