



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:34 AM GMT

PDB ID : 2APJ
Title : X-Ray Structure of Protein from Arabidopsis Thaliana AT4G34215 at 1.6 Angstrom Resolution
Authors : Wesenberg, G.E.; Phillips Jr., G.N.; McCoy, J.G.; Bitto, E.; Bingman, C.A.; Allard, S.T.; Center for Eukaryotic Structural Genomics (CESG)
Deposited on : 2005-08-16
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	
1	B	260	
1	C	260	
1	D	260	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	SEB	A	31	-	-	X	-
1	SEB	B	31	-	-	X	-

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative Esterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	244	Total	C	N	O	S	0	6	0
			1906	1201	336	358	11			
1	B	243	Total	C	N	O	S	0	8	0
			1901	1197	335	358	11			
1	C	241	Total	C	N	O	S	0	2	0
			1869	1172	333	353	11			
1	D	238	Total	C	N	O	S	0	2	0
			1845	1159	330	345	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	SEB	SER	MODIFIED RESIDUE	UNP O49483
B	31	SEB	SER	MODIFIED RESIDUE	UNP O49483
C	31	SEB	SER	MODIFIED RESIDUE	UNP O49483
D	31	SEB	SER	MODIFIED RESIDUE	UNP O49483


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	464	Total	O	0	0
			464	464		
2	B	443	Total	O	0	0
			443	443		
2	C	348	Total	O	0	0
			348	348		
2	D	356	Total	O	0	0
			356	356		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Putative Esterase




• Molecule 1: Putative Esterase



• Molecule 1: Putative Esterase



• Molecule 1: Putative Esterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.69 Å 71.92 Å 93.21 Å 108.16° 93.25° 90.40°	Depositor
Resolution (Å)	35.85 – 1.60 35.85 – 1.60	Depositor EDS
% Data completeness (in resolution range)	94.4 (35.85-1.60) 85.1 (35.85-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.74 (at 1.60 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.146 , 0.183 0.148 , 0.184	Depositor DCC
R_{free} test set	6226 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 30.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 124841 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9132	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.96	4/1952 (0.2%)	0.89	5/2645 (0.2%)
1	B	0.90	4/1954 (0.2%)	0.87	2/2647 (0.1%)
1	C	0.72	0/1898	0.73	1/2568 (0.0%)
1	D	0.73	0/1872	0.74	1/2531 (0.0%)
All	All	0.84	8/7676 (0.1%)	0.81	9/10391 (0.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	242	GLU	CG-CD	7.70	1.63	1.51
1	A	242	GLU	CG-CD	6.50	1.61	1.51
1	A	105	ARG	CB-CG	6.29	1.69	1.52
1	B	242	GLU	CB-CG	6.26	1.64	1.52
1	B	126	LYS	CD-CE	6.02	1.66	1.51
1	B	126	LYS	CE-NZ	5.29	1.62	1.49
1	A	223[A]	VAL	CB-CG2	-5.04	1.42	1.52
1	A	223[B]	VAL	CB-CG2	-5.04	1.42	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	126	LYS	CD-CE-NZ	6.18	125.91	111.70
1	B	158	TYR	CB-CG-CD2	-6.06	117.36	121.00
1	A	126	LYS	CD-CE-NZ	5.98	125.45	111.70
1	A	47	ARG	NE-CZ-NH1	-5.87	117.37	120.30
1	C	166	ASP	CB-CG-OD1	5.59	123.33	118.30
1	A	72	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	A	36	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	D	126	LYS	CD-CE-NZ	5.20	123.66	111.70
1	A	235	ASP	CB-CG-OD1	5.12	122.91	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1906	0	1892	37	0
1	B	1901	0	1892	34	0
1	C	1869	0	1854	20	0
1	D	1845	0	1834	15	0
2	A	464	0	0	17	1
2	B	443	0	0	24	2
2	C	348	0	0	13	0
2	D	356	0	0	7	0
All	All	9132	0	7472	106	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:168:HIS:CE1	2:D:289:HOH:O	1.89	1.22
1:A:31:SEB:HE3	2:A:451:HOH:O	1.49	1.12
1:A:168:HIS:CE1	2:A:361:HOH:O	2.04	1.10
1:B:82[B]:VAL:HG22	2:B:540:HOH:O	1.48	1.09
1:B:31:SEB:HE3	2:B:345:HOH:O	1.51	1.09
1:A:31:SEB:HI1	2:A:475:HOH:O	1.52	1.07
1:C:168:HIS:CE1	2:C:291:HOH:O	2.06	1.06
1:B:168:HIS:CE1	2:B:355:HOH:O	2.07	1.06
1:D:168:HIS:NE2	2:D:289:HOH:O	1.87	1.03
1:A:31:SEB:CH2	1:A:237:LEU:HD21	1.88	1.03
1:C:168:HIS:NE2	2:C:291:HOH:O	1.91	1.03
1:A:168:HIS:NE2	2:A:361:HOH:O	1.98	0.94
1:A:31:SEB:HE2	1:A:238:HIS:NE2	1.85	0.92
1:A:82[B]:VAL:HG12	2:A:455:HOH:O	1.69	0.91
1:A:105:ARG:HD3	2:A:574:HOH:O	1.71	0.90

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:168:HIS:NE2	2:B:355:HOH:O	2.04	0.90
1:B:31:SEB:HJ	2:B:639:HOH:O	1.71	0.88
1:B:31:SEB:HE2	1:B:238:HIS:NE2	1.89	0.88
1:A:105:ARG:HD2	1:A:256:LEU:HD13	1.56	0.86
1:B:106:LEU:O	1:B:108[A]:THR:HG23	1.80	0.81
1:D:20:PRO:N	2:D:485:HOH:O	2.13	0.79
1:A:31:SEB:HH2	1:A:237:LEU:HD11	1.62	0.78
1:B:108[A]:THR:HG21	2:B:519:HOH:O	1.84	0.76
1:B:106:LEU:HB3	1:B:108[B]:THR:HG22	1.67	0.75
1:C:20:PRO:N	2:C:317:HOH:O	2.18	0.75
1:C:144:GLU:OE1	2:C:453:HOH:O	2.06	0.74
1:A:31:SEB:HH2	1:A:237:LEU:HD21	1.68	0.73
1:B:31:SEB:HH1	2:B:446:HOH:O	1.88	0.72
1:A:72:ARG:NH2	2:A:541:HOH:O	2.23	0.71
1:A:246:GLN:NE2	2:A:593:HOH:O	2.12	0.71
1:A:242:GLU:OE2	2:A:721:HOH:O	2.09	0.69
1:C:182:LYS:HG3	2:C:485:HOH:O	1.91	0.69
1:B:237[A]:LEU:HG	2:B:439:HOH:O	1.91	0.69
1:B:237[A]:LEU:CD1	2:B:439:HOH:O	2.40	0.68
1:A:82[B]:VAL:HG13	2:A:691:HOH:O	1.93	0.68
1:D:47:ARG:HD2	2:D:498:HOH:O	1.92	0.68
1:C:31:SEB:HI2	2:C:406:HOH:O	1.95	0.65
1:B:109:ASP:HB2	2:B:414:HOH:O	1.98	0.63
1:B:106:LEU:HB3	1:B:108[B]:THR:CG2	2.28	0.63
1:B:31:SEB:CJ	2:B:639:HOH:O	2.36	0.62
1:B:31:SEB:HH2	2:B:439:HOH:O	1.98	0.61
1:D:110:SER:HB2	2:D:566:HOH:O	2.00	0.61
1:A:31:SEB:HH2	1:A:237:LEU:CD1	2.30	0.60
1:B:137:ARG:HB2	2:B:654:HOH:O	2.02	0.59
1:B:237[A]:LEU:HD11	2:B:439:HOH:O	2.02	0.58
1:C:144:GLU:HG2	2:C:510:HOH:O	2.03	0.58
1:B:237[A]:LEU:CG	2:B:439:HOH:O	2.51	0.58
1:B:31:SEB:CJ	2:B:665:HOH:O	2.51	0.58
1:A:106:LEU:O	1:A:108[A]:THR:HG23	2.05	0.57
1:A:31:SEB:HH2	1:A:237:LEU:CD2	2.35	0.56
1:A:31:SEB:CH2	1:A:237:LEU:HD11	2.35	0.54
1:A:31:SEB:HH2	1:A:237:LEU:CG	2.38	0.53
1:C:138:MET:HE3	1:C:184:LEU:HD11	1.90	0.52
1:C:228:LYS:HD2	2:C:594:HOH:O	2.09	0.52
1:C:72:ARG:HD3	2:C:596:HOH:O	2.09	0.52
1:B:109:ASP:HB2	2:B:545:HOH:O	2.09	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:ILE:CG2	2:A:621:HOH:O	2.59	0.50
1:D:31:SEB:HE2	1:D:238:HIS:NE2	2.25	0.50
1:B:31:SEB:HJ	2:B:665:HOH:O	2.10	0.50
1:B:109:ASP:CG	2:B:414:HOH:O	2.50	0.50
1:A:40[B]:PHE:HE1	1:A:51:ASP:OD2	1.94	0.50
1:C:31:SEB:HE2	1:C:238:HIS:NE2	2.26	0.50
1:C:105:ARG:HG2	1:C:256:LEU:HD13	1.94	0.49
1:A:138:MET:HE3	1:A:184:LEU:HD21	1.95	0.49
1:C:41:LYS:HD2	2:C:529:HOH:O	2.12	0.48
1:B:106:LEU:CB	1:B:108[B]:THR:HG22	2.42	0.48
1:A:82[B]:VAL:CG1	2:A:691:HOH:O	2.54	0.48
1:A:31:SEB:CE	1:A:238:HIS:NE2	2.69	0.48
1:A:32:ASN:ND2	2:A:707:HOH:O	2.47	0.47
1:C:138:MET:HE2	1:C:184:LEU:HD21	1.96	0.47
1:A:106:LEU:HB3	1:A:108[B]:THR:HG23	1.97	0.47
1:B:108[B]:THR:HG21	2:B:576:HOH:O	2.15	0.46
1:A:19:ILE:HG23	2:A:621:HOH:O	2.15	0.46
1:D:255:TYR:CE2	1:D:260:CYS:SG	3.09	0.46
1:A:17:SER:N	2:A:658:HOH:O	2.49	0.46
1:B:137:ARG:CB	2:B:654:HOH:O	2.62	0.46
1:D:23:GLN:OE1	1:D:153:LYS:HE3	2.15	0.45
1:B:31:SEB:CE	1:B:238:HIS:NE2	2.70	0.45
1:D:31:SEB:HH2	1:D:161:GLU:CB	2.46	0.45
1:B:42:ASP:OD1	1:B:44:HIS:HB3	2.17	0.45
1:A:72:ARG:HB2	1:A:72:ARG:CZ	2.47	0.44
1:A:19:ILE:O	1:A:65:LEU:HD11	2.17	0.44
1:D:142:THR:O	1:D:146:ARG:HG3	2.17	0.44
1:D:192:SER:OG	2:D:536:HOH:O	2.21	0.44
1:B:32:ASN:HB3	1:B:158:TYR:CE2	2.53	0.44
1:C:41:LYS:HE3	2:C:479:HOH:O	2.18	0.44
1:C:246:GLN:HG2	2:C:512:HOH:O	2.17	0.44
1:A:138:MET:CE	1:A:184:LEU:HD21	2.48	0.44
1:C:138:MET:CE	1:C:184:LEU:HD21	2.48	0.43
1:A:108[A]:THR:HG22	2:A:713:HOH:O	2.19	0.43
1:D:32:ASN:HB3	1:D:158:TYR:CE2	2.53	0.43
1:D:168:HIS:HE1	2:D:289:HOH:O	1.61	0.42
1:A:106:LEU:O	1:A:108[B]:THR:HG22	2.19	0.42
1:D:31:SEB:HH2	1:D:161:GLU:HB3	2.02	0.42
1:D:196:ILE:HG12	1:D:223[A]:VAL:HG22	2.01	0.42
1:B:108[A]:THR:OG1	1:B:111:ALA:HB2	2.20	0.42
1:B:109:ASP:CB	2:B:545:HOH:O	2.66	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:108[B]:THR:HG23	1:B:111:ALA:HB2	2.02	0.42
1:A:108[B]:THR:HG21	2:A:683:HOH:O	2.19	0.42
1:C:32:ASN:HB3	1:C:158:TYR:CE2	2.55	0.41
1:B:31:SEB:HI1	2:B:688:HOH:O	2.19	0.41
1:C:31:SEB:HH1	2:C:497:HOH:O	2.21	0.41
1:A:32:ASN:HB3	1:A:158:TYR:CE2	2.56	0.41
1:B:109:ASP:CB	2:B:414:HOH:O	2.61	0.40
1:C:196:ILE:HG23	1:C:223:VAL:HG23	2.04	0.40
1:A:42:ASP:OD1	1:A:44:HIS:HB3	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:526:HOH:O	2:B:405:HOH:O[1_455]	2.18	0.02
2:B:475:HOH:O	2:B:629:HOH:O[1_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/260 (95%)	243 (98%)	4 (2%)	0	100	100
1	B	248/260 (95%)	242 (98%)	6 (2%)	0	100	100
1	C	240/260 (92%)	236 (98%)	4 (2%)	0	100	100
1	D	235/260 (90%)	231 (98%)	4 (2%)	0	100	100
All	All	970/1040 (93%)	952 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/215 (97%)	207 (100%)	1 (0%)	92	85
1	B	209/215 (97%)	208 (100%)	1 (0%)	92	85
1	C	201/215 (94%)	199 (99%)	2 (1%)	82	67
1	D	198/215 (92%)	195 (98%)	3 (2%)	72	50
All	All	816/860 (95%)	809 (99%)	7 (1%)	84	71

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	158	TYR
1	B	158	TYR
1	C	158	TYR
1	C	242	GLU
1	D	110	SER
1	D	158	TYR
1	D	192	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	133	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEB	A	31	1	14,16,17	2.15	3 (21%)	15,21,23	3.26	7 (46%)
1	SEB	B	31	1	14,16,17	1.96	4 (28%)	15,21,23	1.35	2 (13%)
1	SEB	C	31	1	14,16,17	2.00	3 (21%)	15,21,23	2.07	4 (26%)
1	SEB	D	31	1	14,16,17	1.84	4 (28%)	15,21,23	1.55	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEB	A	31	1	-	2/10/13/15	0/1/1/1
1	SEB	B	31	1	-	0/10/13/15	0/1/1/1
1	SEB	C	31	1	-	0/10/13/15	0/1/1/1
1	SEB	D	31	1	-	0/10/13/15	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	31	SEB	OG-CB	-2.71	1.34	1.46
1	D	31	SEB	OG-CB	-2.23	1.36	1.46
1	B	31	SEB	CE-CZ	2.72	1.55	1.50
1	D	31	SEB	OD2-SD	2.83	1.51	1.44
1	C	31	SEB	OD2-SD	2.93	1.51	1.44
1	A	31	SEB	OD2-SD	3.44	1.53	1.44
1	B	31	SEB	OD2-SD	3.55	1.53	1.44
1	C	31	SEB	OD1-SD	3.57	1.53	1.44
1	D	31	SEB	CE-CZ	3.79	1.57	1.50
1	D	31	SEB	OD1-SD	3.89	1.54	1.44
1	A	31	SEB	OD1-SD	4.10	1.55	1.44
1	B	31	SEB	OD1-SD	4.78	1.56	1.44
1	C	31	SEB	CE-CZ	5.11	1.60	1.50
1	A	31	SEB	CE-CZ	5.39	1.60	1.50

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	31	SEB	O-C-CA	-3.14	117.30	125.49
1	B	31	SEB	O-C-CA	-3.08	117.46	125.49
1	D	31	SEB	O-C-CA	-2.99	117.71	125.49
1	D	31	SEB	CB-OG-SD	-2.77	112.76	119.10
1	A	31	SEB	O-C-CA	-2.77	118.27	125.49
1	A	31	SEB	CB-OG-SD	-2.58	113.20	119.10
1	A	31	SEB	CE-CZ-CH2	-2.16	117.58	120.52
1	C	31	SEB	CB-OG-SD	-2.10	114.31	119.10
1	A	31	SEB	CJ-CI1-CH1	2.18	123.38	120.19
1	A	31	SEB	CI2-CH2-CZ	2.18	124.12	120.65
1	B	31	SEB	CZ-CE-SD	2.71	115.67	112.58
1	C	31	SEB	CE-CZ-CH1	2.87	124.44	120.52
1	D	31	SEB	CZ-CE-SD	2.95	115.94	112.58
1	A	31	SEB	CE-CZ-CH1	4.10	126.13	120.52
1	C	31	SEB	CZ-CE-SD	5.48	118.81	112.58
1	A	31	SEB	CZ-CE-SD	10.27	124.27	112.58

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	31	SEB	CH1-CZ-CE-SD
1	A	31	SEB	CH2-CZ-CE-SD

There are no ring outliers.

4 monomers are involved in 27 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	31	SEB	11	0
1	B	31	SEB	10	0
1	C	31	SEB	3	0
1	D	31	SEB	3	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/260 (93%)	0.31	2 (0%) 87 87	2, 5, 11, 16	0
1	B	242/260 (93%)	0.29	5 (2%) 67 65	2, 5, 11, 18	0
1	C	240/260 (92%)	0.26	11 (4%) 36 33	2, 5, 13, 27	0
1	D	237/260 (91%)	0.35	13 (5%) 29 25	2, 5, 12, 19	0
All	All	962/1040 (92%)	0.30	31 (3%) 51 48	2, 5, 12, 27	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	108	THR	5.1
1	C	43	HIS	4.7
1	C	44	HIS	4.4
1	C	107	GLU	4.0
1	A	40[A]	PHE	4.0
1	D	148	CYS	4.0
1	D	260	CYS	3.6
1	D	48	TRP	3.4
1	C	47	ARG	3.3
1	B	109	ASP	3.2
1	D	47	ARG	3.1
1	B	43	HIS	3.0
1	D	46	ASN	3.0
1	C	48	TRP	2.9
1	D	40	PHE	2.8
1	C	105	ARG	2.7
1	C	45	ASN	2.7
1	C	109	ASP	2.7
1	D	20	PRO	2.7
1	D	110	SER	2.5
1	D	44	HIS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	43	HIS	2.4
1	B	249	LEU	2.3
1	C	260	CYS	2.3
1	D	245	VAL	2.3
1	B	225[A]	VAL	2.3
1	D	111	ALA	2.2
1	B	223[A]	VAL	2.1
1	C	148	CYS	2.1
1	A	198	VAL	2.0
1	D	41	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	SEB	A	31	16/17	0.93	0.15	-	4,11,28,29	0
1	SEB	B	31	16/17	0.94	0.15	-	3,11,29,29	0
1	SEB	C	31	16/17	0.92	0.14	-	4,14,27,29	0
1	SEB	D	31	16/17	0.92	0.15	-	4,15,30,31	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.