



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:34 AM GMT

PDB ID : 2ASL
Title : oxoG-modified Postinsertion Binary Complex
Authors : Rechko, O.; Malinina, L.; Cheng, Y.; Kuryavyi, V.; Broyde, S.; Geacintov, N.E.; Patel, D.J.
Deposited on : 2005-08-23
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3152 (2.70-2.62)
Clashscore	102246	3524 (2.70-2.62)
Ramachandran outliers	100387	3469 (2.70-2.62)
Sidechain outliers	100360	3469 (2.70-2.62)
RSRZ outliers	91569	3161 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	14	<div> <div></div> <div>57%43%</div> </div>
1	H	14	<div> <div></div> <div>71%29%</div> </div>
2	E	19	<div> <div>11%</div> <div>58%21%11%11%</div> </div>
2	J	19	<div> <div>11%</div> <div>74%11%5%11%</div> </div>
3	A	360	<div> <div>%</div> <div>79%13%• 5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	B	360	 2% 81% 12% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	8OG	E	906	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*GP*TP*TP*GP*GP*AP*TP*GP*GP*TP*AP*GP*(DOC))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	14	Total	C	N	O	P	0	0	0
			291	139	56	83	13			
1	H	14	Total	C	N	O	P	0	0	0
			291	139	56	83	13			

- Molecule 2 is a DNA chain called 5'-D(*CP*T*AP*AP*CP*(8OG)P*CP*TP*AP*CP*CP*AP*TP*CP*CP*AP*AP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	17	Total	C	N	O	P	0	0	0
			338	162	63	97	16			
2	J	17	Total	C	N	O	P	0	0	0
			338	162	63	97	16			

- Molecule 3 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	341	Total	C	N	O	S	0	0	0
			2739	1757	472	504	6			
3	B	341	Total	C	N	O	S	0	0	0
			2739	1757	472	504	6			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	CLONING ARTIFACT	UNP Q97W02
A	-6	SER	-	CLONING ARTIFACT	UNP Q97W02
A	-5	HIS	-	CLONING ARTIFACT	UNP Q97W02
A	-4	MET	-	CLONING ARTIFACT	UNP Q97W02
A	-3	GLY	-	CLONING ARTIFACT	UNP Q97W02
A	-2	GLY	-	CLONING ARTIFACT	UNP Q97W02

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	CLONING ARTIFACT	UNP Q97W02
A	0	GLY	-	CLONING ARTIFACT	UNP Q97W02
A	1	GLY	-	CLONING ARTIFACT	UNP Q97W02
B	-7	GLY	-	CLONING ARTIFACT	UNP Q97W02
B	-6	SER	-	CLONING ARTIFACT	UNP Q97W02
B	-5	HIS	-	CLONING ARTIFACT	UNP Q97W02
B	-4	MET	-	CLONING ARTIFACT	UNP Q97W02
B	-3	GLY	-	CLONING ARTIFACT	UNP Q97W02
B	-2	GLY	-	CLONING ARTIFACT	UNP Q97W02
B	-1	GLY	-	CLONING ARTIFACT	UNP Q97W02
B	0	GLY	-	CLONING ARTIFACT	UNP Q97W02
B	1001	GLY	-	CLONING ARTIFACT	UNP Q97W02

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ca 1 1	0	0
4	A	1	Total Ca 1 1	0	0
4	D	1	Total Ca 1 1	0	0
4	J	1	Total Ca 1 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	66	Total O 66 66	0	0
5	B	60	Total O 60 60	0	0
5	D	14	Total O 14 14	0	0
5	E	15	Total O 15 15	0	0
5	H	12	Total O 12 12	0	0
5	J	14	Total O 14 14	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*GP*GP*TP*TP*GP*GP*AP*TP*GP*GP*TP*AP*GP*(DOC))-3'

Chain D: 



- Molecule 1: 5'-D(*GP*GP*TP*TP*GP*GP*AP*TP*GP*GP*TP*AP*GP*(DOC))-3'

Chain H: 




- Molecule 2: 5'-D(*CP*T*AP*AP*CP*(8OG)P*CP*TP*AP*CP*CP*AP*TP*CP*CP*AP*AP*CP*C)-3'

Chain E: 





- Molecule 2: 5'-D(*CP*T*AP*AP*CP*(8OG)P*CP*TP*AP*CP*CP*AP*TP*CP*CP*AP*AP*CP*C)-3'


Chain J: 





- Molecule 3: DNA polymerase IV

Chain A: 





● Molecule 3: DNA polymerase IV



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.22Å 101.02Å 84.92Å 90.00° 97.16° 90.00°	Depositor
Resolution (Å)	20.00 – 2.65 25.14 – 2.65	Depositor EDS
% Data completeness (in resolution range)	91.8 (20.00-2.65) 91.7 (25.14-2.65)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 2.64Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.232 , 0.287 0.213 , 0.260	Depositor DCC
R_{free} test set	3290 reflections (11.22%)	DCC
Wilson B-factor (Å ²)	44.0	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 32665 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6921	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.21 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.9181e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 8OG, CA, DOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.85	0/307	1.53	5/475 (1.1%)
1	H	0.78	0/307	1.57	4/475 (0.8%)
2	E	0.90	0/351	1.60	7/534 (1.3%)
2	J	0.85	0/351	1.57	5/534 (0.9%)
3	A	0.51	0/2778	0.74	9/3731 (0.2%)
3	B	0.48	0/2778	0.73	8/3731 (0.2%)
All	All	0.58	0/6872	0.98	38/9480 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	E	1	0
3	A	0	1
3	B	0	1
All	All	1	2

There are no bond length outliers.

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	809	DG	O4'-C1'-N9	9.59	114.71	108.00
2	J	1905	DC	O4'-C1'-N1	8.86	114.20	108.00
1	H	1803	DT	O4'-C1'-N1	8.05	113.64	108.00
2	E	919	DC	O4'-C1'-N1	8.03	113.62	108.00
2	J	1905	DC	C1'-O4'-C4'	-7.72	102.38	110.10
3	B	1007	ASP	CB-CG-OD2	6.84	124.46	118.30
1	H	1809	DG	O4'-C1'-N9	6.77	112.74	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1105	ASP	CB-CG-OD2	6.59	124.23	118.30
3	A	294	ASP	CB-CG-OD2	6.39	124.05	118.30
3	B	1292	ASP	CB-CG-OD2	6.25	123.92	118.30
3	B	1294	ASP	CB-CG-OD2	6.08	123.78	118.30
2	J	1918	DC	O4'-C1'-N1	6.08	112.26	108.00
1	H	1808	DT	P-O3'-C3'	6.07	126.98	119.70
2	E	918	DC	O4'-C4'-C3'	-6.06	102.08	104.50
2	E	910	DC	O4'-C1'-N1	6.03	112.22	108.00
1	D	811	DT	O4'-C1'-N1	5.95	112.16	108.00
2	E	918	DC	O4'-C1'-N1	5.92	112.14	108.00
3	A	182	ASP	CB-CG-OD2	5.72	123.45	118.30
3	A	292	ASP	CB-CG-OD2	5.65	123.39	118.30
1	D	813	DG	O4'-C1'-N9	5.63	111.94	108.00
3	A	117	ASP	CB-CG-OD2	5.61	123.35	118.30
3	A	277	ASP	CB-CG-OD2	5.53	123.27	118.30
3	A	110	ASP	CB-CG-OD2	5.51	123.26	118.30
2	J	1918	DC	C1'-O4'-C4'	-5.47	104.62	110.10
3	A	105	ASP	CB-CG-OD2	5.34	123.11	118.30
2	J	1904	DA	O4'-C1'-N9	5.33	111.73	108.00
3	A	204	ASP	CB-CG-OD2	5.33	123.10	118.30
2	E	919	DC	C6-N1-C2	-5.32	118.17	120.30
1	D	810	DG	O4'-C1'-N9	5.29	111.70	108.00
3	B	1168	ASP	CB-CG-OD2	5.28	123.05	118.30
2	E	913	DT	C6-C5-C7	-5.14	119.81	122.90
3	B	1182	ASP	CB-CG-OD2	5.14	122.93	118.30
3	A	211	ASP	CB-CG-OD2	5.13	122.92	118.30
3	B	1117	ASP	CB-CG-OD2	5.11	122.89	118.30
2	E	918	DC	C1'-O4'-C4'	-5.07	105.03	110.10
3	B	1277	ASP	CB-CG-OD2	5.05	122.85	118.30
1	H	1802	DG	O4'-C1'-N9	5.04	111.53	108.00
1	D	803	DT	O4'-C1'-N1	5.01	111.51	108.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	E	906	8OG	C1'

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	159	LYS	Peptide
3	B	1159	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	291	0	160	1	0
1	H	291	0	160	0	0
2	E	338	0	188	4	0
2	J	338	0	189	1	0
3	A	2739	0	2883	29	0
3	B	2739	0	2880	30	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
4	J	1	0	0	0	0
5	A	66	0	0	15	0
5	B	60	0	0	12	0
5	D	14	0	0	0	0
5	E	15	0	0	0	0
5	H	12	0	0	0	0
5	J	14	0	0	0	0
All	All	6921	0	6460	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1224:TYR:OH	5:B:128:HOH:O	1.74	1.04
3:B:1010:TYR:HB3	5:B:79:HOH:O	1.60	1.00
3:A:14:GLN:HE22	3:A:139:THR:H	1.22	0.88
3:A:10:TYR:HB3	5:A:553:HOH:O	1.75	0.85
3:B:1014:GLN:HE22	3:B:1139:THR:H	1.29	0.81
3:A:336:ARG:CZ	5:A:565:HOH:O	2.28	0.79
3:B:1049:GLU:HG3	5:B:135:HOH:O	1.82	0.79
3:A:48:TYR:HB3	5:A:517:HOH:O	1.84	0.77
3:B:1048:TYR:CZ	3:B:1160:PRO:HD3	2.20	0.76
3:A:52:LYS:NZ	5:A:517:HOH:O	2.22	0.73
3:B:1273:TYR:OH	3:B:1306:ILE:O	2.07	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1048:TYR:HB3	5:B:114:HOH:O	1.91	0.71
3:B:1105:ASP:OD2	5:B:1503:HOH:O	2.13	0.66
2:E:918:DC:H2''	2:E:919:DC:H5'	1.80	0.64
3:B:1239:THR:HA	5:B:70:HOH:O	1.98	0.63
3:A:48:TYR:CD1	5:A:553:HOH:O	2.51	0.62
3:A:48:TYR:CE1	3:A:160:PRO:HD3	2.36	0.61
3:B:1048:TYR:OH	3:B:1160:PRO:HD3	2.02	0.59
3:B:1038:GLU:O	3:B:1039:ASP:HB2	2.03	0.58
3:B:1014:GLN:NE2	3:B:1139:THR:H	1.99	0.57
3:A:267:ARG:HD2	5:A:528:HOH:O	2.04	0.55
3:A:158:ALA:O	3:A:159:LYS:O	2.23	0.55
2:E:918:DC:H2''	2:E:919:DC:C5'	2.38	0.54
3:B:1010:TYR:HD1	3:B:1013:ALA:HB3	1.73	0.53
3:A:49:GLU:HG3	5:A:554:HOH:O	2.09	0.53
3:A:304:HIS:HB2	5:A:504:HOH:O	2.09	0.53
3:A:226:ILE:O	3:A:230:ARG:HG3	2.10	0.52
3:A:336:ARG:NH2	5:A:565:HOH:O	2.36	0.52
3:A:17:GLU:OE1	3:A:24:LYS:NZ	2.34	0.52
3:B:1048:TYR:CE1	5:B:79:HOH:O	2.54	0.51
3:A:48:TYR:CE1	5:A:553:HOH:O	2.63	0.50
3:B:1159:LYS:HB3	3:B:1160:PRO:CD	2.41	0.50
3:B:1046:ALA:HB1	3:B:1050:ALA:HB3	1.93	0.50
3:A:243:LYS:HD3	5:A:549:HOH:O	2.12	0.50
1:D:814:DOC:H5'	5:A:502:HOH:O	2.13	0.49
3:A:159:LYS:HB3	3:A:160:PRO:CD	2.42	0.49
2:J:1904:DA:N1	3:B:1066:LYS:NZ	2.58	0.49
3:B:1009:ASP:O	3:B:1010:TYR:C	2.52	0.47
3:B:1052:LYS:NZ	5:B:114:HOH:O	2.47	0.47
3:A:9:ASP:O	3:A:10:TYR:C	2.52	0.47
3:A:273:TYR:OH	3:A:306:ILE:O	2.29	0.47
3:A:48:TYR:CZ	3:A:160:PRO:HD3	2.50	0.47
3:B:1048:TYR:CZ	3:B:1160:PRO:CD	2.95	0.46
3:B:1248:ILE:HB	3:B:1334:GLY:HA3	1.97	0.46
2:E:918:DC:C2'	2:E:919:DC:H5'	2.46	0.46
3:B:1180:ILE:HD12	3:B:1202:LEU:HA	1.98	0.46
3:B:1014:GLN:HE22	3:B:1139:THR:N	2.06	0.46
3:A:12:TYR:HB2	3:A:45:THR:CG2	2.47	0.45
3:A:296:VAL:HG11	3:A:322:ILE:HG23	1.99	0.44
3:B:1045:THR:HG21	5:B:10:HOH:O	2.16	0.44
3:A:285:HIS:ND1	3:A:299:GLY:HA3	2.33	0.44
3:A:285:HIS:HD2	5:A:515:HOH:O	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1010:TYR:CD1	3:B:1010:TYR:O	2.71	0.43
3:B:1048:TYR:HE2	5:B:132:HOH:O	2.03	0.42
3:B:1007:ASP:OD1	5:B:1503:HOH:O	2.21	0.42
3:A:12:TYR:HB2	3:A:45:THR:HG21	2.02	0.42
3:B:1041:GLY:HA3	3:B:1061:ILE:HD12	2.02	0.41
3:A:196:LYS:HG3	5:A:552:HOH:O	2.19	0.41
3:A:106:GLU:CD	5:A:503:HOH:O	2.58	0.41
3:A:170:GLU:O	3:A:174:LEU:HG	2.21	0.41
2:E:906:8OG:H2'	2:E:907:DC:O5'	2.21	0.40
3:A:8:PHE:N	3:A:8:PHE:CD1	2.88	0.40
3:B:1304:HIS:HB2	5:B:4:HOH:O	2.20	0.40
3:B:1095:TYR:CD1	3:B:1124:LEU:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	339/360 (94%)	322 (95%)	15 (4%)	2 (1%)	30	54
3	B	339/360 (94%)	324 (96%)	12 (4%)	3 (1%)	21	44
All	All	678/720 (94%)	646 (95%)	27 (4%)	5 (1%)	26	51

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	159	LYS
3	A	160	PRO
3	B	1160	PRO
3	B	1159	LYS
3	B	1010	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	299/311 (96%)	277 (93%)	22 (7%)	17	35
3	B	299/311 (96%)	281 (94%)	18 (6%)	24	47
All	All	598/622 (96%)	558 (93%)	40 (7%)	20	41

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	20	ASN
3	A	23	LEU
3	A	24	LYS
3	A	36	ARG
3	A	45	THR
3	A	52	LYS
3	A	63	GLU
3	A	78	LYS
3	A	83	GLN
3	A	114	LYS
3	A	145	SER
3	A	170	GLU
3	A	230	ARG
3	A	238	ARG
3	A	242	ARG
3	A	248	ILE
3	A	252	LYS
3	A	262	LYS
3	A	273	TYR
3	A	323	LEU
3	A	327	GLU
3	A	331	ARG
3	B	1023	LEU
3	B	1026	LYS
3	B	1045	THR
3	B	1079	GLU
3	B	1093	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	B	1145	SER
3	B	1179	ASP
3	B	1195	LYS
3	B	1200	ASN
3	B	1212	LYS
3	B	1238	ARG
3	B	1242	ARG
3	B	1243	LYS
3	B	1248	ILE
3	B	1252	LYS
3	B	1273	TYR
3	B	1292	ASP
3	B	1336	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	14	GLN
3	A	20	ASN
3	A	285	HIS
3	B	1014	GLN
3	B	1285	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	DOC	D	814	1,2,4	11,19,20	0.79	0	14,26,29	1.11	1 (7%)
2	8OG	E	906	1,2	16,25,26	3.26	4 (25%)	21,37,40	4.72	13 (61%)
1	DOC	H	1814	1,2,4	11,19,20	0.85	0	14,26,29	1.62	3 (21%)
2	8OG	J	1906	1,2	16,25,26	3.25	4 (25%)	21,37,40	4.45	11 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DOC	D	814	1,2,4	-	0/3/18/19	0/2/2/2
2	8OG	E	906	1,2	1/1/4/4	0/3/21/22	0/3/3/3
1	DOC	H	1814	1,2,4	-	0/3/18/19	0/2/2/2
2	8OG	J	1906	1,2	-	0/3/21/22	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	1906	8OG	O3'-C3'	-8.89	1.23	1.43
2	E	906	8OG	O3'-C3'	-8.87	1.23	1.43
2	E	906	8OG	C3'-C4'	-7.60	1.31	1.53
2	J	1906	8OG	C3'-C4'	-7.55	1.31	1.53
2	J	1906	8OG	C5-C4	3.42	1.48	1.40
2	J	1906	8OG	C6-C5	3.54	1.48	1.41
2	E	906	8OG	C5-C4	3.67	1.48	1.40
2	E	906	8OG	C6-C5	3.86	1.49	1.41

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	906	8OG	C4'-O4'-C1'	-5.63	95.23	109.47
2	J	1906	8OG	C4'-O4'-C1'	-5.25	96.19	109.47
2	E	906	8OG	C6-C5-C4	-4.71	115.26	120.90
2	J	1906	8OG	C6-C5-C4	-4.48	115.54	120.90
2	J	1906	8OG	C5-C6-N1	-4.34	117.66	123.59
2	E	906	8OG	C5-C6-N1	-4.29	117.72	123.59
2	J	1906	8OG	N3-C2-N1	-3.41	122.25	127.44
2	J	1906	8OG	C3'-C2'-C1'	-2.57	96.21	102.40
2	E	906	8OG	N3-C2-N1	-2.56	123.55	127.44
2	E	906	8OG	C3'-C2'-C1'	-2.34	96.76	102.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	906	8OG	C1'-N9-C4	-2.04	124.21	127.37
2	J	1906	8OG	C2'-C3'-C4'	2.02	106.96	102.77
1	H	1814	DOC	O4'-C1'-C2'	2.12	108.96	106.67
2	E	906	8OG	C2'-C3'-C4'	2.16	107.26	102.77
2	E	906	8OG	O4'-C4'-C5'	2.38	117.84	109.32
2	E	906	8OG	O4'-C1'-C2'	2.78	111.82	106.27
1	H	1814	DOC	C3'-C2'-C1'	3.00	106.06	102.71
2	J	1906	8OG	O4'-C1'-C2'	3.16	112.58	106.27
1	D	814	DOC	C2-N3-C4	3.49	120.54	115.61
1	H	1814	DOC	C2-N3-C4	3.61	120.70	115.61
2	J	1906	8OG	O3'-C3'-C2'	3.71	123.03	110.74
2	E	906	8OG	C6-N1-C2	4.89	122.73	115.94
2	J	1906	8OG	C6-N1-C2	5.29	123.28	115.94
2	E	906	8OG	O3'-C3'-C2'	5.44	128.74	110.74
2	J	1906	8OG	O4'-C1'-N9	8.01	114.99	108.22
2	E	906	8OG	C2'-C1'-N9	10.06	125.91	115.83
2	J	1906	8OG	C2'-C1'-N9	13.77	129.64	115.83
2	E	906	8OG	O4'-C1'-N9	13.98	120.04	108.22

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	E	906	8OG	C1'

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	814	DOC	1	0
2	E	906	8OG	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	13/14 (92%)	-0.77	0 100 100	7, 12, 16, 16	0
1	H	13/14 (92%)	-0.39	0 100 100	12, 15, 17, 18	0
2	E	16/19 (84%)	-0.19	2 (12%) 5 4	11, 14, 21, 25	0
2	J	16/19 (84%)	0.12	2 (12%) 5 4	11, 15, 20, 23	0
3	A	341/360 (94%)	-0.14	5 (1%) 76 75	8, 15, 20, 24	0
3	B	341/360 (94%)	0.06	8 (2%) 64 62	9, 15, 20, 25	0
All	All	740/786 (94%)	-0.06	17 (2%) 64 62	7, 15, 20, 25	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	1038	GLU	3.9
3	B	1274	TYR	3.7
3	A	274	TYR	3.3
2	J	1903	DA	3.1
2	E	904	DA	3.0
3	A	38	GLU	2.9
3	B	1188	ASN	2.9
3	B	1304	HIS	2.8
2	J	1904	DA	2.7
3	A	304	HIS	2.6
3	B	1277	ASP	2.6
3	B	1010	TYR	2.5
3	A	10	TYR	2.4
3	B	1001	GLY	2.4
2	E	903	DA	2.3
3	B	1210	PHE	2.3
3	A	39	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	DOC	D	814	18/19	0.93	0.14	-	24,29,33,33	0
1	DOC	H	1814	18/19	0.92	0.14	-	29,32,36,36	0
2	8OG	E	906	23/24	0.98	0.14	-	4,8,9,9	0
2	8OG	J	1906	23/24	0.97	0.16	-	4,7,8,8	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CA	B	1415	1/1	0.86	0.21	-	14,14,14,14	0
4	CA	J	1416	1/1	0.97	0.29	-	30,30,30,30	0
4	CA	D	416	1/1	0.96	0.20	-	36,36,36,36	0
4	CA	A	415	1/1	0.97	0.05	-	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.