



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:28 AM GMT

PDB ID : 3AWF  
Title : Crystal structure of Pten-like domain of Ci-VSP (236-576)  
Authors : Matsuda, M.; Sakata, S.; Takeshita, K.; Suzuki, M.; Yamashita, E.; Okamura, Y.; Nakagawa, A.  
Deposited on : 2011-03-19  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	<div> <div></div> <div>76% 11% • 12%</div> </div>
1	B	346	<div> <div></div> <div>77% 10% • 11%</div> </div>
1	C	346	<div> <div>3%</div> <div>78% 9% • 10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	C	1	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Voltage-sensor containing phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	5	0
			2464	1569	415	465	15			
1	B	308	Total	C	N	O	S	0	4	0
			2510	1596	429	470	15			
1	C	310	Total	C	N	O	S	0	4	0
			2516	1596	431	474	15			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	231	GLY	-	EXPRESSION TAG	UNP Q4W8A1
A	232	PRO	-	EXPRESSION TAG	UNP Q4W8A1
A	233	LEU	-	EXPRESSION TAG	UNP Q4W8A1
A	234	GLY	-	EXPRESSION TAG	UNP Q4W8A1
A	235	SER	-	EXPRESSION TAG	UNP Q4W8A1
B	231	GLY	-	EXPRESSION TAG	UNP Q4W8A1
B	232	PRO	-	EXPRESSION TAG	UNP Q4W8A1
B	233	LEU	-	EXPRESSION TAG	UNP Q4W8A1
B	234	GLY	-	EXPRESSION TAG	UNP Q4W8A1
B	235	SER	-	EXPRESSION TAG	UNP Q4W8A1
C	231	GLY	-	EXPRESSION TAG	UNP Q4W8A1
C	232	PRO	-	EXPRESSION TAG	UNP Q4W8A1
C	233	LEU	-	EXPRESSION TAG	UNP Q4W8A1
C	234	GLY	-	EXPRESSION TAG	UNP Q4W8A1
C	235	SER	-	EXPRESSION TAG	UNP Q4W8A1

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	195	Total	O	0	0
			195	195		
4	B	176	Total	O	0	0
			176	176		
4	C	164	Total	O	0	0
			164	164		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Voltage-sensor containing phosphatase



- Molecule 1: Voltage-sensor containing phosphatase



- Molecule 1: Voltage-sensor containing phosphatase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.56 Å   177.44 Å   50.94 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	44.17 – 1.99 44.17 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.8 (44.17-1.99) 98.8 (44.17-1.99)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.13 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.5.0071	Depositor
R, $R_{free}$	0.179   ,   0.223 0.179   ,   0.225	Depositor DCC
$R_{free}$ test set	4094 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 54.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 81627 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8081	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.23	5/2525 (0.2%)	0.96	3/3407 (0.1%)
1	B	1.16	2/2566 (0.1%)	0.96	5/3462 (0.1%)
1	C	1.16	5/2572 (0.2%)	0.91	1/3468 (0.0%)
All	All	1.18	12/7663 (0.2%)	0.95	9/10337 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	2
All	All	0	5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	495[A]	CYS	CB-SG	-5.84	1.72	1.81
1	C	495[B]	CYS	CB-SG	-5.84	1.72	1.81
1	A	312	GLU	CB-CG	5.83	1.63	1.52
1	B	312	GLU	CG-CD	5.61	1.60	1.51
1	A	310	CYS	CB-SG	-5.32	1.73	1.81
1	C	452	VAL	CB-CG2	5.28	1.64	1.52
1	A	349	VAL	CB-CG2	5.25	1.63	1.52
1	A	444	VAL	CB-CG2	5.24	1.63	1.52
1	A	312	GLU	CD-OE2	5.13	1.31	1.25
1	C	480	TYR	CD1-CE1	5.06	1.47	1.39
1	B	560	TYR	CD2-CE2	5.03	1.46	1.39
1	C	527	PHE	CE1-CZ	5.01	1.46	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	367	LYS	N-CA-C	5.76	126.56	111.00
1	B	433	LEU	CA-CB-CG	5.75	128.53	115.30
1	A	405	ASP	CB-CG-OD1	5.49	123.24	118.30
1	A	369	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	C	433	LEU	CA-CB-CG	5.34	127.59	115.30
1	B	480	TYR	CA-CB-CG	5.29	123.45	113.40
1	B	331	ASP	CB-CG-OD2	5.25	123.03	118.30
1	B	365	GLY	N-CA-C	-5.11	100.33	113.10
1	A	454	ARG	CG-CD-NE	-5.03	101.23	111.80

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	362	HIS	Peptide
1	B	362	HIS	Peptide
1	B	366	GLY	Peptide
1	C	362	HIS	Peptide
1	C	571	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2464	0	2417	22	0
1	B	2510	0	2461	18	0
1	C	2516	0	2459	22	0
2	A	15	0	0	0	0
2	B	15	0	0	0	0
2	C	20	0	0	0	0
3	C	6	0	8	2	0
4	A	195	0	0	4	0
4	B	176	0	0	1	0
4	C	164	0	0	2	0
All	All	8081	0	7345	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:466:SER:O	1:B:469:GLN:HG2	1.77	0.83
1:A:546[A]:GLU:HG2	4:A:578:HOH:O	1.81	0.79
1:C:424:LYS:HD3	1:C:535:LEU:HD22	1.69	0.73
1:A:421:TYR:CD2	1:A:535[B]:LEU:HD22	2.24	0.72
1:C:572:ASP:OD1	1:C:572:ASP:C	2.29	0.71
1:C:405:ASP:HB3	1:C:408:GLN:HG3	1.78	0.64
1:B:313:ARG:NH1	4:B:229:HOH:O	2.32	0.62
1:C:341:LEU:HD12	1:C:433:LEU:HD13	1.83	0.60
1:A:380:GLU:OE1	1:A:426:LYS:HE3	2.01	0.60
1:B:424:LYS:HD3	1:B:535:LEU:HD22	1.84	0.59
1:B:341:LEU:HD12	1:B:433:LEU:HD13	1.84	0.59
1:C:303:LYS:HD3	1:C:356:ASP:CB	2.34	0.58
1:B:572:ASP:OD1	1:B:572:ASP:C	2.40	0.58
1:A:456:ASN:C	1:A:456:ASN:HD22	2.08	0.57
1:C:366:GLY:HA2	4:C:641:HOH:O	2.05	0.57
1:C:303:LYS:HD3	1:C:356:ASP:HB3	1.89	0.55
1:C:478:GLU:HG3	1:C:480:TYR:HB2	1.89	0.55
1:A:287:ASN:HB3	1:A:292:VAL:CG2	2.38	0.54
1:C:545:ARG:HB3	3:C:1:GOL:H12	1.89	0.53
1:A:367:LYS:HD3	1:A:398:ARG:CZ	2.40	0.51
1:A:333:ASN:OD1	1:A:508:LYS:HE3	2.10	0.51
1:C:516:LYS:HE3	1:C:516:LYS:H	1.75	0.51
1:B:303:LYS:HD3	1:B:356:ASP:HB2	1.93	0.51
1:A:421:TYR:CG	1:A:535[B]:LEU:HD22	2.46	0.51
1:B:341:LEU:CD1	1:B:433:LEU:HD13	2.41	0.50
1:B:367:LYS:HG3	1:B:368:GLY:N	2.25	0.50
1:A:437:LYS:HB2	1:A:534:SER:HA	1.94	0.50
1:A:276:PHE:HB2	1:A:364:LYS:HD3	1.94	0.50
1:C:549:ASP:O	1:C:550:ASN:HB2	2.12	0.49
1:A:367:LYS:HG3	1:A:411:GLU:HG3	1.94	0.49
1:C:270:HIS:HA	1:C:358:VAL:HG12	1.94	0.49
1:A:405:ASP:HB3	1:A:408:GLN:HG2	1.94	0.49
1:A:366:GLY:HA2	4:A:39:HOH:O	2.13	0.49
1:C:475:LYS:HB2	1:C:480:TYR:HB3	1.95	0.48
1:A:348:LYS:HD2	1:A:380:GLU:OE2	2.14	0.48
1:C:367:LYS:HG2	1:C:398:ARG:CZ	2.44	0.47
1:A:456:ASN:ND2	1:A:458:SER:H	2.13	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:303:LYS:HD3	1:C:356:ASP:HB2	1.96	0.47
1:A:330:ASP:OD1	1:A:468:ARG:NH2	2.47	0.47
1:C:511:PHE:O	1:C:526:PRO:HD2	2.15	0.46
1:B:494[A]:THR:HG22	1:B:495[A]:CYS:N	2.30	0.46
1:B:367:LYS:HG3	1:B:411:GLU:HG3	1.97	0.46
1:A:485:GLN:HG3	4:A:675:HOH:O	2.17	0.45
1:B:301:PRO:O	1:B:302:ASP:HB2	2.17	0.45
1:C:550:ASN:HB2	1:C:551:PRO:HD3	1.99	0.45
1:B:445:THR:HG23	1:B:494[B]:THR:HG22	1.98	0.44
1:B:494[A]:THR:CG2	1:B:495[A]:CYS:N	2.81	0.44
1:C:545:ARG:HB3	3:C:1:GOL:C1	2.47	0.44
1:B:287:ASN:HB3	1:B:292:VAL:CG2	2.48	0.43
1:A:518:LEU:HD12	1:A:519:PRO:HD2	2.00	0.43
1:C:437:LYS:HD3	1:C:439:LEU:HD21	2.00	0.43
1:A:397:ARG:HB3	1:A:397:ARG:HE	1.62	0.43
1:C:301:PRO:O	1:C:302:ASP:HB2	2.18	0.43
1:C:253:ARG:NH1	1:C:256:ARG:HH22	2.17	0.42
1:A:278:SER:O	1:A:288:PRO:HA	2.20	0.41
1:A:437:LYS:HB3	1:A:437:LYS:HE2	1.56	0.41
1:B:443:GLY:HA2	1:B:495[B]:CYS:O	2.20	0.41
1:C:408:GLN:HG2	4:C:665:HOH:O	2.19	0.41
1:B:337:LEU:HD21	1:B:425:ILE:HD11	2.02	0.41
1:A:469:GLN:NE2	4:A:683:HOH:O	2.54	0.41
1:B:541[B]:VAL:HG13	1:B:570:PHE:HE1	1.86	0.40
1:B:448:ALA:HB3	1:B:564:PHE:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	303/346 (88%)	294 (97%)	8 (3%)	1 (0%)	46 41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	306/346 (88%)	297 (97%)	8 (3%)	1 (0%)	46 41
1	C	308/346 (89%)	302 (98%)	5 (2%)	1 (0%)	46 41
All	All	917/1038 (88%)	893 (97%)	21 (2%)	3 (0%)	46 41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	550	ASN
1	B	550	ASN
1	C	550	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	275/307 (90%)	272 (99%)	3 (1%)	80 83
1	B	279/307 (91%)	271 (97%)	8 (3%)	50 49
1	C	279/307 (91%)	266 (95%)	13 (5%)	32 27
All	All	833/921 (90%)	809 (97%)	24 (3%)	50 49

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	LEU
1	A	426	LYS
1	A	456	ASN
1	B	257	LYS
1	B	278	SER
1	B	398	ARG
1	B	433	LEU
1	B	467	GLU
1	B	480	TYR
1	B	518	LEU
1	B	572	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	253	ARG
1	C	279	SER
1	C	302	ASP
1	C	313	ARG
1	C	364	LYS
1	C	367	LYS
1	C	398	ARG
1	C	426	LYS
1	C	437	LYS
1	C	516	LYS
1	C	520	ARG
1	C	523	ASP
1	C	572	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	332	HIS
1	A	428	ASN
1	A	456	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1	-	4,4,4	0.74	0	6,6,6	0.63	0
2	SO4	A	6	-	4,4,4	0.23	0	6,6,6	1.18	1 (16%)
2	SO4	A	9	-	4,4,4	0.17	0	6,6,6	0.33	0
2	SO4	B	2	-	4,4,4	1.30	1 (25%)	6,6,6	0.46	0
2	SO4	B	5	-	4,4,4	0.35	0	6,6,6	0.84	0
2	SO4	B	8	-	4,4,4	0.72	0	6,6,6	0.77	0
3	GOL	C	1	-	5,5,5	0.75	0	5,5,5	2.12	3 (60%)
2	SO4	C	10	-	4,4,4	0.12	0	6,6,6	0.21	0
2	SO4	C	3	-	4,4,4	0.31	0	6,6,6	0.31	0
2	SO4	C	4	-	4,4,4	0.09	0	6,6,6	0.37	0
2	SO4	C	7	-	4,4,4	0.44	0	6,6,6	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1	-	-	0/0/0/0	0/0/0/0
2	SO4	A	6	-	-	0/0/0/0	0/0/0/0
2	SO4	A	9	-	-	0/0/0/0	0/0/0/0
2	SO4	B	2	-	-	0/0/0/0	0/0/0/0
2	SO4	B	5	-	-	0/0/0/0	0/0/0/0
2	SO4	B	8	-	-	0/0/0/0	0/0/0/0
3	GOL	C	1	-	-	0/4/4/4	0/0/0/0
2	SO4	C	10	-	-	0/0/0/0	0/0/0/0
2	SO4	C	3	-	-	0/0/0/0	0/0/0/0
2	SO4	C	4	-	-	0/0/0/0	0/0/0/0
2	SO4	C	7	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	SO4	O1-S	-2.23	1.39	1.47

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	C	1	GOL	C3-C2-C1	-2.59	100.97	111.12
2	A	6	SO4	O2-S-O1	-2.36	102.03	109.50
3	C	1	GOL	O3-C3-C2	2.24	121.04	110.18
3	C	1	GOL	O2-C2-C3	2.36	119.45	108.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	304/346 (87%)	0.04	2 (0%) 89 89	14, 23, 40, 56	0
1	B	308/346 (89%)	0.11	5 (1%) 74 75	16, 26, 42, 53	0
1	C	310/346 (89%)	0.13	11 (3%) 48 49	15, 28, 45, 55	0
All	All	922/1038 (88%)	0.09	18 (1%) 68 69	14, 26, 43, 56	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	253	ARG	3.5
1	C	313	ARG	2.9
1	C	404	GLY	2.9
1	C	279	SER	2.6
1	C	294	ARG	2.5
1	B	489	THR	2.4
1	B	450	GLN	2.4
1	B	537	GLU	2.3
1	C	365	GLY	2.3
1	C	280	GLY	2.2
1	C	572	ASP	2.2
1	A	489	THR	2.2
1	C	432	GLN	2.1
1	C	302	ASP	2.1
1	C	293	SER	2.1
1	A	404	GLY	2.1
1	B	538	GLY	2.0
1	B	469	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	C	1	6/6	0.95	0.14	3.46	21,30,34,35	0
2	SO4	B	5	5/5	0.93	0.14	0.79	56,56,58,62	0
2	SO4	C	4	5/5	0.85	0.16	0.68	84,85,86,86	0
2	SO4	B	2	5/5	0.99	0.11	-0.16	23,25,28,29	0
2	SO4	A	9	5/5	0.89	0.13	-0.21	71,71,74,74	0
2	SO4	C	7	5/5	0.92	0.12	-0.43	29,38,41,44	0
2	SO4	B	8	5/5	0.94	0.11	-0.63	33,36,44,46	0
2	SO4	C	3	5/5	0.98	0.11	-0.68	48,50,53,53	0
2	SO4	A	6	5/5	0.99	0.08	-1.44	28,32,35,37	0
2	SO4	A	1	5/5	0.99	0.08	-1.61	34,34,37,38	0
2	SO4	C	10	5/5	0.97	0.17	-	57,60,61,61	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.