



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:29 PM GMT

PDB ID : 1B0H
Title : OLIGO-PEPTIDE BINDING PROTEIN COMPLEXED WITH LYSYL-NA
PTHYLALANYL-LYSINE
Authors : Davies, T.G.; Tame, J.R.H.
Deposited on : 1998-11-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	517	 87% 12% •
2	B	3	 67% 33%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PERIPLASMIC OLIGOPEPTIDE-BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	517	Total	C	N	O	S	34	0	0
			4165	2666	700	794	5			

- Molecule 2 is a protein called LYS-ALN-LYS PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			34	25	5	4			

- Molecule 3 is URANIUM ATOM (three-letter code: U1) (formula: U).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	8	Total	U	0	0
			8	8		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	412	Total	O	0	0
			412	412		
4	B	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PERIPLASMIC OLIGOPEPTIDE-BINDING PROTEIN



• Molecule 2: LYS-ALN-LYS PEPTIDE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	109.96 Å 75.46 Å 70.58 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.90	Depositor
% Data completeness (in resolution range)	99.0 (15.00-1.90)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.185 , 0.224	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4622	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ALN, U1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	5/4276 (0.1%)	1.26	26/5830 (0.4%)
2	B	0.61	0/17	1.46	0/16
All	All	0.67	5/4293 (0.1%)	1.26	26/5846 (0.4%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	177	ASP	CB-CG	15.53	1.84	1.51
1	A	384	LYS	CG-CD	-9.21	1.21	1.52
1	A	342	GLU	CG-CD	7.64	1.63	1.51
1	A	78	GLU	CB-CG	7.35	1.66	1.52
1	A	338	GLN	CG-CD	-5.22	1.39	1.51

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	206	ARG	NE-CZ-NH2	-20.96	109.82	120.30
1	A	413	ARG	NE-CZ-NH2	-14.43	113.08	120.30
1	A	413	ARG	CD-NE-CZ	10.38	138.13	123.60
1	A	201	ARG	NE-CZ-NH2	9.76	125.18	120.30
1	A	384	LYS	CB-CG-CD	9.70	136.83	111.60
1	A	206	ARG	NH1-CZ-NH2	9.23	129.55	119.40
1	A	433	ASP	CB-CG-OD1	9.14	126.52	118.30
1	A	84	ASP	CB-CG-OD1	7.71	125.24	118.30
1	A	485	TYR	CA-CB-CG	7.39	127.44	113.40
1	A	8	GLN	CA-CB-CG	7.12	129.06	113.40
1	A	141	ASP	CB-CG-OD2	7.05	124.64	118.30
1	A	433	ASP	CB-CG-OD2	-7.03	111.97	118.30
1	A	413	ARG	NH1-CZ-NH2	6.97	127.07	119.40
1	A	410	ASP	CB-CG-OD1	-6.52	112.44	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	265	ARG	NE-CZ-NH2	-6.31	117.14	120.30
1	A	177	ASP	CB-CG-OD1	-5.98	112.92	118.30
1	A	393	GLU	OE1-CD-OE2	-5.91	116.21	123.30
1	A	269	TYR	CB-CG-CD2	-5.59	117.65	121.00
1	A	102	ASP	CB-CG-OD1	-5.48	113.37	118.30
1	A	99	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	A	201	ARG	CD-NE-CZ	5.39	131.15	123.60
1	A	286	ASP	CB-CG-OD2	5.39	123.15	118.30
1	A	206	ARG	N-CA-CB	-5.30	101.07	110.60
1	A	338	GLN	CB-CG-CD	5.22	125.18	111.60
1	A	290	ARG	NE-CZ-NH2	5.15	122.88	120.30
1	A	362	ASP	CB-CG-OD2	-5.01	113.79	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4165	0	4076	27	0
2	B	34	0	35	1	0
3	A	8	0	0	0	0
4	A	412	0	0	4	0
4	B	3	0	0	0	0
All	All	4622	0	4111	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:ILE:HA	1:A:305:LYS:HD2	1.61	0.83
1:A:62:GLU:HG3	1:A:63:LYS:HD2	1.77	0.67
1:A:229:GLU:HB3	1:A:249:PRO:HD3	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:460:THR:O	1:A:464:GLU:HG3	2.07	0.55
1:A:62:GLU:HG3	1:A:63:LYS:CD	2.36	0.54
1:A:382:TRP:HB3	1:A:388:VAL:HG22	1.89	0.54
1:A:115:TYR:CE1	1:A:428:ASN:HB3	2.44	0.52
1:A:117:HIS:HE1	4:A:1057:HOH:O	1.93	0.52
1:A:51:ASP:OD1	1:A:55:HIS:HD2	1.94	0.51
1:A:114:GLN:NE2	4:A:1309:HOH:O	2.44	0.50
1:A:117:HIS:HD2	4:A:1118:HOH:O	1.95	0.50
1:A:382:TRP:HB3	1:A:388:VAL:CG2	2.44	0.48
1:A:43:LEU:O	1:A:187:THR:HB	2.15	0.47
1:A:402:ASP:O	1:A:406:GLN:HG3	2.15	0.47
1:A:453:THR:HG21	1:A:466:TYR:CE2	2.51	0.46
1:A:31:ILE:HB	1:A:36:GLU:HB3	1.98	0.45
1:A:4:PRO:O	1:A:7:VAL:HG13	2.16	0.45
1:A:52:VAL:HG22	4:A:1041:HOH:O	2.17	0.44
1:A:274:TYR:OH	2:B:602:ALN:HE3	2.18	0.43
1:A:294:LYS:HA	1:A:480:VAL:HG13	1.99	0.43
1:A:119:ALA:O	1:A:120:ASN:HB2	2.18	0.43
1:A:43:LEU:HD21	1:A:204:LEU:HD22	2.01	0.42
1:A:265:ARG:O	1:A:488:ALA:HA	2.20	0.42
1:A:3:VAL:HA	1:A:4:PRO:HD3	1.90	0.42
1:A:307:LYS:HE2	1:A:370:LEU:HD11	2.02	0.41
1:A:230:VAL:HG13	1:A:252:LEU:HD11	2.02	0.41
1:A:323:ASP:O	1:A:423:PRO:HD3	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	515/517 (100%)	499 (97%)	15 (3%)	1 (0%)	52 42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/455 (100%)	446 (98%)	9 (2%)	63	57
2	B	2/2 (100%)	2 (100%)	0	100	100
All	All	457/457 (100%)	448 (98%)	9 (2%)	63	57

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	VAL
1	A	52	VAL
1	A	63	LYS
1	A	67	LYS
1	A	109	TYR
1	A	177	ASP
1	A	312	LEU
1	A	384	LYS
1	A	413	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	HIS
1	A	117	HIS
1	A	195	ASN
1	A	199	ASN
1	A	279	ASN
1	A	304	ASN
1	A	406	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ALN	B	602	2	15,16,17	1.00	0	15,21,23	2.03	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALN	B	602	2	-	0/4/6/8	0/2/2/2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	602	ALN	CZ1-CE2-CD2	-4.44	113.09	119.10
2	B	602	ALN	CE1-CZ1-CE2	2.05	123.73	120.44
2	B	602	ALN	CD3-CG2-CD2	2.98	125.11	120.88
2	B	602	ALN	CG1-CD2-CE2	3.99	124.19	118.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	602	ALN	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.