



# Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 07:29 AM GMT

PDB ID : 3B1U  
Title : Crystal structure of human peptidylarginine deiminase 4 in complex with o-F-amidine  
Authors : Causey, C.P.; Jones, J.E.; Slack, J.L.; Kamei, D.; Jones Jr, L.E.; Subramanian, V.; Knuckley, B.; Ebrahimi, P.; Chumanovich, A.A.; Luo, Y.; Hashimoto, H.; Shimizu, T.; Sato, M.; Hofseth, L.J.; Thompson, P.R.  
Deposited on : 2011-07-13  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	YFF	A	800	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	906	-	-	-	X

## 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5164 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein-arginine deiminase type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	628	4947	3158	829	925	35	0	0	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	EXPRESSION TAG	UNP Q9UM07
A	-6	PRO	-	EXPRESSION TAG	UNP Q9UM07
A	-5	LEU	-	EXPRESSION TAG	UNP Q9UM07
A	-4	GLY	-	EXPRESSION TAG	UNP Q9UM07
A	-3	SER	-	EXPRESSION TAG	UNP Q9UM07
A	-2	PRO	-	EXPRESSION TAG	UNP Q9UM07
A	-1	GLU	-	EXPRESSION TAG	UNP Q9UM07
A	0	PHE	-	EXPRESSION TAG	UNP Q9UM07
A	55	SER	GLY	SEE REMARK 999	UNP Q9UM07
A	82	ALA	VAL	SEE REMARK 999	UNP Q9UM07
A	112	ALA	GLY	SEE REMARK 999	UNP Q9UM07

- Molecule 2 is 2-{|(2S)-1-AMINO-5-{|(1Z)-2-FLUOROETHANIMIDOYL|AMINO}-1-O XOPENTAN-2-YL|CARBAMOYL}BENZOIC ACID (three-letter code: YFF) (formula: C<sub>15</sub>H<sub>19</sub>FN<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	23	15	4	4	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	O	S			
3	A	1	5	4	1		0	0
3	A	1	5	4	1		0	0
3	A	1	5	4	1		0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	5	Total Ca 5 5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	174	Total O 174 174	0	0

### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein-arginine deiminase type-4



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	146.42Å    60.47Å    115.51Å 90.00°    124.43°    90.00°	Depositor
Resolution (Å)	20.00 – 2.10 19.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.10) 91.5 (19.88-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.47 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
$R$ , $R_{free}$	0.219 , 0.250 0.230 , 0.263	Depositor DCC
$R_{free}$ test set	2254 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	51.7	Xtriage
Anisotropy	0.222	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.33$	Xtriage
Outliers	0 of 44717 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5164	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4, YFF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/5066	0.49	0/6874

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4947	0	4905	67	0
2	A	23	0	15	0	0
3	A	15	0	0	0	0
4	A	5	0	0	0	0
5	A	174	0	0	0	0
All	All	5164	0	4920	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:313:ILE:HG12	1:A:314:PHE:N	1.58	1.14
1:A:313:ILE:CG1	1:A:314:PHE:H	1.71	1.04
1:A:313:ILE:HG12	1:A:314:PHE:HD2	1.31	0.95
1:A:313:ILE:HG12	1:A:314:PHE:H	0.79	0.90
1:A:313:ILE:CG1	1:A:314:PHE:HD2	1.87	0.88
1:A:425:LEU:HD12	1:A:456:VAL:HG13	1.58	0.86
1:A:367:VAL:HG21	1:A:384:VAL:CG1	2.12	0.80
1:A:367:VAL:HG21	1:A:384:VAL:HG12	1.67	0.77
1:A:42:ILE:HG21	1:A:50:VAL:HG11	1.66	0.76
1:A:6:LEU:HD21	1:A:28:ASP:HB2	1.68	0.73
1:A:313:ILE:HG12	1:A:314:PHE:CD2	2.21	0.72
1:A:298:MET:HE1	1:A:412:VAL:HG22	1.71	0.72
1:A:27:LEU:HD13	1:A:78:LEU:HD13	1.71	0.71
1:A:623:LEU:HB2	1:A:625:LEU:HD23	1.73	0.70
1:A:201:LEU:HD13	1:A:229:LEU:HD11	1.73	0.69
1:A:314:PHE:O	1:A:316:ASN:N	2.28	0.67
1:A:354:ILE:HD12	1:A:367:VAL:HG22	1.79	0.65
1:A:425:LEU:HD12	1:A:456:VAL:CG1	2.26	0.64
1:A:6:LEU:HD21	1:A:28:ASP:CB	2.28	0.63
1:A:52:ILE:HD12	1:A:76:VAL:HG22	1.81	0.62
1:A:313:ILE:CG1	1:A:314:PHE:CD2	2.78	0.60
1:A:185:LEU:HD22	1:A:187:THR:HG23	1.84	0.59
1:A:313:ILE:HD11	1:A:314:PHE:CD2	2.37	0.59
1:A:314:PHE:H	1:A:314:PHE:HD2	1.52	0.57
1:A:108:LEU:HG	1:A:110:LEU:HD13	1.86	0.57
1:A:201:LEU:HD13	1:A:229:LEU:CD1	2.34	0.57
1:A:367:VAL:HG21	1:A:384:VAL:HG11	1.84	0.56
1:A:71:ASP:O	1:A:74:VAL:HG12	2.06	0.56
1:A:462:LEU:CD2	1:A:552:LEU:HD22	2.38	0.54
1:A:123:ARG:HD3	1:A:659:TRP:CD1	2.43	0.54
1:A:367:VAL:HG23	1:A:389:PHE:CE2	2.44	0.53
1:A:10:THR:HG22	1:A:32:SER:CB	2.39	0.52
1:A:74:VAL:HG13	1:A:74:VAL:O	2.10	0.52
1:A:497:CYS:SG	1:A:570:LEU:HD13	2.50	0.51
1:A:313:ILE:CD1	1:A:314:PHE:CD2	2.93	0.51
1:A:216:ALA:O	1:A:217:THR:HG22	2.11	0.51
1:A:462:LEU:HD23	1:A:552:LEU:HD22	1.93	0.51
1:A:428:ILE:HD13	1:A:451:LEU:HD22	1.93	0.50
1:A:33:ALA:CB	1:A:70:LEU:HD11	2.41	0.50
1:A:166:CYS:HB2	1:A:254:LEU:HD22	1.95	0.49
1:A:27:LEU:HD11	1:A:78:LEU:HD22	1.96	0.48
1:A:326:LEU:HD23	1:A:592:LEU:HD22	1.96	0.48

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:354:ILE:CD1	1:A:367:VAL:HG22	2.45	0.47
1:A:456:VAL:HG21	1:A:658:TRP:HB2	1.97	0.47
1:A:198:THR:HG23	1:A:272:LEU:HB2	1.96	0.46
1:A:594:LYS:HA	1:A:625:LEU:HD13	1.96	0.46
1:A:10:THR:HG22	1:A:32:SER:HB3	1.97	0.46
1:A:314:PHE:O	1:A:315:GLU:C	2.55	0.46
1:A:48:VAL:HG23	1:A:90:GLN:HE21	1.81	0.45
1:A:337:CYS:SG	1:A:342:ASN:ND2	2.90	0.45
1:A:466:TRP:CZ3	1:A:542:VAL:HG13	2.52	0.44
1:A:401:ILE:HG23	1:A:402:SER:N	2.32	0.44
1:A:313:ILE:O	1:A:314:PHE:O	2.35	0.44
1:A:427:ARG:HD2	1:A:458:ALA:O	2.18	0.44
1:A:313:ILE:C	1:A:314:PHE:O	2.56	0.44
1:A:334:LEU:HD11	1:A:336:ILE:HG13	1.99	0.44
1:A:467:LEU:HD13	1:A:474:GLU:HB2	1.99	0.43
1:A:40:PHE:CE1	1:A:42:ILE:HD13	2.53	0.43
1:A:52:ILE:HG23	1:A:68:TRP:CZ2	2.53	0.43
1:A:33:ALA:HB1	1:A:70:LEU:HD11	2.00	0.43
1:A:45:SER:OG	1:A:90:GLN:NE2	2.52	0.43
1:A:312:SER:HA	1:A:320:LEU:HD11	2.01	0.43
1:A:298:MET:CE	1:A:412:VAL:HG22	2.43	0.42
1:A:298:MET:HB3	1:A:412:VAL:HG21	1.99	0.42
1:A:113:VAL:HB	1:A:189:THR:HG22	2.01	0.42
1:A:152:VAL:HA	1:A:253:ALA:HB3	2.02	0.41
1:A:184:THR:HG22	1:A:248:ASP:OD2	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	618/671 (92%)	598 (97%)	17 (3%)	3 (0%)	34 30

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	314	PHE
1	A	315	GLU
1	A	458	ALA

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	559/594 (94%)	531 (95%)	28 (5%)	30   27

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	TRP
1	A	80	MET
1	A	110	LEU
1	A	117	LEU
1	A	150	LEU
1	A	159	LEU
1	A	169	ASP
1	A	185	LEU
1	A	201	LEU
1	A	206	SER
1	A	278	GLU
1	A	313	ILE
1	A	314	PHE
1	A	315	GLU
1	A	317	GLU
1	A	343	MET
1	A	364	LEU
1	A	374	ARG
1	A	389	PHE
1	A	412	VAL
1	A	429	LEU
1	A	456	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	489	LEU
1	A	500	LEU
1	A	514	LEU
1	A	557	LEU
1	A	570	LEU
1	A	592	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	90	GLN
1	A	136	GLN
1	A	178	GLN
1	A	303	GLN
1	A	342	ASN
1	A	448	GLN
1	A	504	GLN
1	A	505	GLN
1	A	524	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 9 ligands modelled in this entry, 5 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	YFF	A	800	1	19,23,24	0.90	1 (5%)	23,30,31	1.15	2 (8%)
3	SO4	A	905	-	4,4,4	0.23	0	6,6,6	0.09	0
3	SO4	A	906	-	4,4,4	0.26	0	6,6,6	0.06	0
3	SO4	A	907	-	4,4,4	0.23	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	YFF	A	800	1	-	0/18/23/25	0/1/1/1
3	SO4	A	905	-	-	0/0/0/0	0/0/0/0
3	SO4	A	906	-	-	0/0/0/0	0/0/0/0
3	SO4	A	907	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	800	YFF	CAV-NAC	3.48	1.32	1.27

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	800	YFF	CAO-CAZ-NAS	-3.58	104.05	110.87
2	A	800	YFF	CAM-CAN-NAR	-2.56	104.70	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	628/671 (93%)	0.94	106 (16%) <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">2</span> <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">3</span>	56, 78, 110, 137	0

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	217	THR	12.2
1	A	159	LEU	8.0
1	A	314	PHE	7.5
1	A	74	VAL	6.3
1	A	137	ARG	5.6
1	A	522	LYS	5.4
1	A	136	GLN	5.0
1	A	576	PHE	4.9
1	A	519	LYS	4.7
1	A	160	GLU	4.6
1	A	125	GLY	4.5
1	A	213	VAL	4.4
1	A	419	ARG	4.2
1	A	520	LYS	4.1
1	A	70	LEU	4.1
1	A	100	LYS	4.1
1	A	328	MET	4.0
1	A	315	GLU	4.0
1	A	251	VAL	4.0
1	A	94	ILE	4.0
1	A	12	GLU	3.9
1	A	67	THR	3.9
1	A	162	SER	3.9
1	A	104	VAL	3.8
1	A	658	TRP	3.8
1	A	4	GLY	3.8
1	A	81	LYS	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	78	LEU	3.7
1	A	68	TRP	3.7
1	A	163	ALA	3.7
1	A	161	SER	3.6
1	A	13	GLN	3.6
1	A	383	ARG	3.6
1	A	612	LEU	3.6
1	A	570	LEU	3.5
1	A	663	PRO	3.5
1	A	10	THR	3.4
1	A	232	LYS	3.4
1	A	24	LEU	3.4
1	A	31	SER	3.3
1	A	340	GLU	3.2
1	A	92	VAL	3.2
1	A	172	LEU	3.2
1	A	276	ASN	3.1
1	A	583	PHE	3.0
1	A	151	LEU	3.0
1	A	49	VAL	3.0
1	A	138	THR	3.0
1	A	575	GLU	3.0
1	A	38	THR	2.9
1	A	149	ILE	2.9
1	A	144	CYS	2.9
1	A	339	GLU	2.9
1	A	514	LEU	2.9
1	A	35	GLU	2.9
1	A	233	TRP	2.8
1	A	146	GLN	2.8
1	A	105	LYS	2.8
1	A	41	SER	2.8
1	A	466	TRP	2.8
1	A	229	LEU	2.7
1	A	244	LYS	2.7
1	A	33	ALA	2.7
1	A	310	ALA	2.7
1	A	439	ASP	2.7
1	A	489	LEU	2.7
1	A	126	LYS	2.6
1	A	52	ILE	2.6
1	A	22	GLY	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	294	ALA	2.5
1	A	399	GLY	2.5
1	A	649	VAL	2.5
1	A	400	GLY	2.5
1	A	376	LEU	2.5
1	A	357	ILE	2.5
1	A	71	ASP	2.5
1	A	507	GLU	2.5
1	A	150	LEU	2.5
1	A	201	LEU	2.5
1	A	216	ALA	2.4
1	A	46	PRO	2.4
1	A	643	VAL	2.4
1	A	607	ASN	2.3
1	A	297	ILE	2.3
1	A	295	PRO	2.3
1	A	113	VAL	2.3
1	A	467	LEU	2.3
1	A	571	PHE	2.2
1	A	48	VAL	2.2
1	A	367	VAL	2.2
1	A	124	THR	2.2
1	A	366	VAL	2.2
1	A	608	GLY	2.2
1	A	214	PHE	2.2
1	A	250	TYR	2.2
1	A	19	CYS	2.1
1	A	278	GLU	2.1
1	A	40	PHE	2.1
1	A	501	PHE	2.1
1	A	51	ASP	2.1
1	A	18	VAL	2.1
1	A	239	MET	2.1
1	A	96	TYR	2.1
1	A	344	ASP	2.1
1	A	401	ILE	2.0
1	A	275	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	906	5/5	0.96	0.29	3.90	94,94,94,94	0
2	YFF	A	800	23/24	0.79	0.27	2.77	61,63,64,64	0
3	SO4	A	905	5/5	0.94	0.22	1.51	92,92,92,92	0
4	CA	A	904	1/1	0.97	0.15	-0.40	54,54,54,54	0
4	CA	A	902	1/1	0.98	0.10	-1.03	54,54,54,54	0
4	CA	A	901	1/1	0.97	0.10	-2.07	54,54,54,54	0
4	CA	A	900	1/1	0.99	0.03	-2.45	49,49,49,49	0
4	CA	A	903	1/1	0.97	0.12	-	59,59,59,59	0
3	SO4	A	907	5/5	0.88	0.32	-	87,87,87,87	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.