



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 8, 2016 – 10:41 AM EDT

PDB ID : 5B1Y  
Title : Crystal structure of NADPH bound carbonyl reductase from *Aeropyrum*  
          pernix  
Authors : Yoneda, K.; Sakuraba, H.; Fukuda, Y.; Araki, T.; Ohshima, T.  
Deposited on : 2015-12-22  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027674  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027674


## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>13%</div> <div>•</div> <div>13%</div> </div> </div>
1	B	270	<div> <div>6%</div> <div> <div></div> <div>73%</div> <div>14%</div> <div>•</div> <div>10%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NDP	A	300	-	-	-	X
2	NDP	B	300	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-[acyl-carrier-protein] reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	234	Total	C	N	O	S	0	0	0
			1760	1112	315	330	3			
1	B	243	Total	C	N	O	S	0	0	0
			1833	1158	330	342	3			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q9Y8Y1
A	-18	GLY	-	expression tag	UNP Q9Y8Y1
A	-17	SER	-	expression tag	UNP Q9Y8Y1
A	-16	SER	-	expression tag	UNP Q9Y8Y1
A	-15	HIS	-	expression tag	UNP Q9Y8Y1
A	-14	HIS	-	expression tag	UNP Q9Y8Y1
A	-13	HIS	-	expression tag	UNP Q9Y8Y1
A	-12	HIS	-	expression tag	UNP Q9Y8Y1
A	-11	HIS	-	expression tag	UNP Q9Y8Y1
A	-10	HIS	-	expression tag	UNP Q9Y8Y1
A	-9	SER	-	expression tag	UNP Q9Y8Y1
A	-8	SER	-	expression tag	UNP Q9Y8Y1
A	-7	GLY	-	expression tag	UNP Q9Y8Y1
A	-6	LEU	-	expression tag	UNP Q9Y8Y1
A	-5	VAL	-	expression tag	UNP Q9Y8Y1
A	-4	PRO	-	expression tag	UNP Q9Y8Y1
A	-3	ARG	-	expression tag	UNP Q9Y8Y1
A	-2	GLY	-	expression tag	UNP Q9Y8Y1
A	-1	SER	-	expression tag	UNP Q9Y8Y1
A	0	HIS	-	expression tag	UNP Q9Y8Y1
B	-19	MET	-	expression tag	UNP Q9Y8Y1
B	-18	GLY	-	expression tag	UNP Q9Y8Y1
B	-17	SER	-	expression tag	UNP Q9Y8Y1
B	-16	SER	-	expression tag	UNP Q9Y8Y1
B	-15	HIS	-	expression tag	UNP Q9Y8Y1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP Q9Y8Y1
B	-13	HIS	-	expression tag	UNP Q9Y8Y1
B	-12	HIS	-	expression tag	UNP Q9Y8Y1
B	-11	HIS	-	expression tag	UNP Q9Y8Y1
B	-10	HIS	-	expression tag	UNP Q9Y8Y1
B	-9	SER	-	expression tag	UNP Q9Y8Y1
B	-8	SER	-	expression tag	UNP Q9Y8Y1
B	-7	GLY	-	expression tag	UNP Q9Y8Y1
B	-6	LEU	-	expression tag	UNP Q9Y8Y1
B	-5	VAL	-	expression tag	UNP Q9Y8Y1
B	-4	PRO	-	expression tag	UNP Q9Y8Y1
B	-3	ARG	-	expression tag	UNP Q9Y8Y1
B	-2	GLY	-	expression tag	UNP Q9Y8Y1
B	-1	SER	-	expression tag	UNP Q9Y8Y1
B	0	HIS	-	expression tag	UNP Q9Y8Y1

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	
			48	21	7	17	3	0
2	B	1	Total	C	N	O	P	
			48	21	7	17	3	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	133	Total 133	O 133	0	0
3	B	111	Total 111	O 111	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.63Å 91.63Å 177.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.10 – 2.09 33.12 – 2.09	Depositor EDS
% Data completeness (in resolution range)	99.7 (33.10-2.09) 99.6 (33.12-2.09)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.55 (at 2.10Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.230 , 0.242 0.229 , 0.242	Depositor DCC
$R_{free}$ test set	2273 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 56.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3933	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/1784	0.60	0/2420
1	B	0.42	0/1858	0.62	0/2519
All	All	0.44	0/3642	0.61	0/4939

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1760	0	1831	34	0
1	B	1833	0	1910	62	0
2	A	48	0	26	15	0
2	B	48	0	26	9	0
3	A	133	0	0	5	0
3	B	111	0	0	3	1
All	All	3933	0	3793	98	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:300:NDP:H52N	3:A:413:HOH:O	1.28	1.28
1:A:35:LYS:HB3	2:A:300:NDP:O1X	1.46	1.15
1:B:190:THR:HG21	2:B:300:NDP:O1N	1.46	1.15
1:B:35:LYS:HB3	2:B:300:NDP:O1X	1.55	1.06
1:B:215:ILE:H	1:B:215:ILE:HD13	1.18	1.03
1:A:63:VAL:H	2:A:300:NDP:H62A	0.96	0.92
1:A:192:MET:SD	2:A:300:NDP:O1A	2.28	0.92
1:A:63:VAL:N	2:A:300:NDP:H62A	1.69	0.89
2:A:300:NDP:H52A	3:A:413:HOH:O	1.72	0.89
1:A:35:LYS:CB	2:A:300:NDP:O1X	2.21	0.89
1:B:5:TYR:H	1:B:84:ASN:HD22	1.24	0.86
1:B:215:ILE:O	1:B:217:LYS:HE3	1.78	0.84
1:B:215:ILE:N	1:B:215:ILE:HD13	1.93	0.83
1:B:215:ILE:CG1	1:B:217:LYS:NZ	2.44	0.81
1:B:215:ILE:HG12	1:B:217:LYS:NZ	1.98	0.79
1:A:5:TYR:H	1:A:84:ASN:HD22	1.31	0.78
1:B:190:THR:CG2	1:B:191:ASP:N	2.50	0.75
1:B:192:MET:HB3	3:B:457:HOH:O	1.88	0.73
1:B:215:ILE:H	1:B:215:ILE:CD1	1.99	0.72
1:A:140:ALA:O	2:A:300:NDP:H6N	1.90	0.71
1:B:190:THR:CG2	1:B:191:ASP:H	2.05	0.70
1:B:186:SER:HB2	1:B:187:PHE:CE2	2.27	0.69
1:B:215:ILE:HG13	1:B:217:LYS:HZ1	1.58	0.68
1:B:197:LEU:HD12	1:B:197:LEU:N	2.09	0.68
1:B:215:ILE:N	1:B:215:ILE:CD1	2.56	0.68
1:B:63:VAL:H	2:B:300:NDP:H62A	1.39	0.68
1:B:190:THR:HG22	1:B:191:ASP:N	2.09	0.67
1:B:190:THR:HG23	1:B:191:ASP:H	1.60	0.66
1:B:150:VAL:HG23	1:B:150:VAL:O	1.95	0.65
1:B:215:ILE:CG1	1:B:217:LYS:HZ1	2.07	0.64
1:B:215:ILE:HG12	1:B:217:LYS:HZ2	1.63	0.64
1:A:189:GLU:HB2	1:A:215:ILE:HD11	1.78	0.63
1:B:12:SER:O	1:B:13:ARG:HG2	2.00	0.62
1:A:75:ARG:HG2	1:A:75:ARG:HH11	1.65	0.62
1:B:197:LEU:H	1:B:197:LEU:HD12	1.64	0.62
1:B:215:ILE:HG13	1:B:217:LYS:NZ	2.13	0.61
1:B:215:ILE:HG12	1:B:217:LYS:CE	2.30	0.61
1:A:232:ARG:HD2	3:A:421:HOH:O	2.02	0.60
1:B:72:MET:O	1:B:76:VAL:HG23	2.02	0.60
1:A:63:VAL:HG22	2:A:300:NDP:N6A	2.18	0.58
1:B:233:ARG:HG2	1:B:233:ARG:HH11	1.68	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:LYS:CB	2:B:300:NDP:O1X	2.42	0.58
1:B:215:ILE:HG12	1:B:217:LYS:HE3	1.87	0.56
1:A:12:SER:O	1:A:13:ARG:HG3	2.05	0.56
1:A:215:ILE:HG23	1:A:217:LYS:NZ	2.20	0.56
1:A:231:PRO:O	1:A:235:ARG:HB3	2.07	0.55
1:B:187:PHE:HE2	1:B:248:ARG:HH21	1.55	0.54
1:B:233:ARG:NH1	3:B:402:HOH:O	2.41	0.54
1:B:59:VAL:HA	1:B:75:ARG:NH1	2.22	0.54
2:A:300:NDP:C5B	3:A:413:HOH:O	2.39	0.53
1:B:186:SER:HB2	1:B:187:PHE:CD2	2.44	0.52
1:B:233:ARG:NH1	1:B:233:ARG:HG2	2.23	0.52
1:B:63:VAL:N	2:B:300:NDP:H62A	2.09	0.51
1:B:215:ILE:CG1	1:B:217:LYS:HZ2	2.20	0.50
1:A:10:GLY:HA2	2:A:300:NDP:H1B	1.93	0.50
1:B:140:ALA:O	2:B:300:NDP:H6N	2.12	0.48
1:A:215:ILE:HG23	1:A:217:LYS:HZ1	1.79	0.48
1:B:197:LEU:N	1:B:197:LEU:CD1	2.77	0.47
1:B:186:SER:O	1:B:188:VAL:HG23	2.15	0.47
1:A:63:VAL:HG13	2:A:300:NDP:N1A	2.30	0.47
1:B:172:GLN:NE2	3:B:403:HOH:O	2.48	0.47
1:A:22:ARG:NH2	1:A:26:GLU:OE2	2.48	0.46
1:A:210:HIS:HD2	1:A:212:LEU:O	1.99	0.46
1:B:103:ILE:O	1:B:107:GLU:HG3	2.15	0.46
1:A:93:LEU:HD12	1:A:94:GLN:H	1.80	0.46
1:B:115:THR:HG22	1:B:119:LEU:HD22	1.98	0.46
1:B:9:THR:HA	1:B:33:ALA:HB3	1.97	0.45
1:B:187:PHE:HE1	1:B:210:HIS:ND1	2.15	0.45
1:B:215:ILE:O	1:B:215:ILE:HG12	2.16	0.45
1:B:84:ASN:O	1:B:134:ALA:HA	2.17	0.44
1:B:187:PHE:HE2	1:B:248:ARG:HE	1.65	0.44
1:A:230:ASP:HA	1:A:231:PRO:HD2	1.89	0.44
1:B:150:VAL:CG2	1:B:150:VAL:O	2.63	0.44
1:B:62:ASP:HA	2:B:300:NDP:H62A	1.81	0.44
1:A:91:GLY:H	2:A:300:NDP:H3D	1.83	0.44
1:A:49:ARG:HD3	3:A:496:HOH:O	2.16	0.44
1:A:4:THR:HA	1:A:84:ASN:ND2	2.33	0.43
1:B:59:VAL:HG13	1:B:75:ARG:NE	2.33	0.43
1:B:4:THR:HA	1:B:84:ASN:ND2	2.33	0.43
1:B:212:LEU:HB2	1:B:246:ALA:HB1	1.99	0.43
1:A:191:ASP:O	1:A:192:MET:HG3	2.19	0.43
1:B:22:ARG:NH1	1:B:26:GLU:HG2	2.34	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:196:PHE:O	1:B:197:LEU:C	2.57	0.43
1:B:22:ARG:HH12	1:B:26:GLU:HG2	1.84	0.43
1:B:35:LYS:HB2	2:B:300:NDP:C6A	2.49	0.43
1:A:84:ASN:O	1:A:134:ALA:HA	2.19	0.43
1:A:230:ASP:OD2	1:A:233:ARG:NH1	2.52	0.43
1:A:125:LEU:HD23	1:B:99:GLU:CG	2.49	0.42
1:A:91:GLY:N	2:A:300:NDP:H3D	2.34	0.42
1:A:9:THR:HA	1:A:33:ALA:HB3	2.00	0.42
1:B:192:MET:SD	1:B:192:MET:O	2.78	0.42
1:A:22:ARG:HD3	1:A:223:GLU:OE2	2.19	0.41
1:A:210:HIS:HA	1:A:211:PRO:HD3	1.89	0.41
1:A:91:GLY:CA	2:A:300:NDP:H3D	2.50	0.41
1:B:190:THR:HG21	2:B:300:NDP:PN	2.54	0.41
1:B:230:ASP:HA	1:B:231:PRO:HD2	1.93	0.41
1:A:83:LEU:O	1:A:128:LEU:HD23	2.21	0.41
1:B:72:MET:C	1:B:72:MET:SD	3.00	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:402:HOH:O	3:B:402:HOH:O[8_555]	1.30	0.90

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/270 (85%)	223 (97%)	7 (3%)	0	100	100
1	B	239/270 (88%)	230 (96%)	9 (4%)	0	100	100
All	All	469/540 (87%)	453 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/217 (85%)	182 (98%)	3 (2%)	70	76
1	B	193/217 (89%)	181 (94%)	12 (6%)	23	19
All	All	378/434 (87%)	363 (96%)	15 (4%)	38	38

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	ARG
1	A	215	ILE
1	A	216	LEU
1	B	1	MET
1	B	60	ARG
1	B	119	LEU
1	B	187	PHE
1	B	192	MET
1	B	193	THR
1	B	194	ARG
1	B	210	HIS
1	B	215	ILE
1	B	217	LYS
1	B	233	ARG
1	B	248	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN
1	A	172	GLN
1	A	210	HIS
1	B	84	ASN
1	B	172	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NDP	A	300	-	44,52,52	1.89	13 (29%)	55,80,80	3.31	14 (25%)
2	NDP	B	300	-	44,52,52	1.90	12 (27%)	55,80,80	3.31	14 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	300	-	-	0/30/77/77	0/5/5/5
2	NDP	B	300	-	-	0/30/77/77	0/5/5/5

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	300	NDP	C4N-C5N	-4.50	1.39	1.49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	NDP	C4N-C5N	-4.48	1.39	1.49
2	A	300	NDP	C5A-C4A	-3.44	1.32	1.40
2	B	300	NDP	C5A-C4A	-3.44	1.32	1.40
2	B	300	NDP	O4B-C4B	-3.30	1.37	1.45
2	A	300	NDP	O4B-C4B	-3.30	1.37	1.45
2	B	300	NDP	P2B-O2X	-2.61	1.45	1.54
2	A	300	NDP	P2B-O2X	-2.60	1.45	1.54
2	A	300	NDP	P2B-O3X	-2.38	1.46	1.54
2	B	300	NDP	P2B-O3X	-2.37	1.46	1.54
2	A	300	NDP	PA-O2A	-2.05	1.46	1.55
2	A	300	NDP	C2B-C1B	-2.05	1.47	1.53
2	B	300	NDP	PA-O2A	-2.04	1.46	1.55
2	B	300	NDP	C2B-C1B	-2.03	1.47	1.53
2	A	300	NDP	C6A-C5A	-2.01	1.32	1.42
2	A	300	NDP	C3B-C2B	2.17	1.58	1.53
2	B	300	NDP	C3B-C2B	2.19	1.58	1.53
2	A	300	NDP	O2B-C2B	2.61	1.52	1.44
2	B	300	NDP	O2B-C2B	2.67	1.52	1.44
2	A	300	NDP	C6N-C5N	2.75	1.38	1.33
2	B	300	NDP	C6N-C5N	2.77	1.38	1.33
2	A	300	NDP	C2A-N1A	3.04	1.39	1.33
2	B	300	NDP	C2A-N1A	3.17	1.40	1.33
2	B	300	NDP	C2A-N3A	5.40	1.41	1.32
2	A	300	NDP	C2A-N3A	5.47	1.41	1.32

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	NDP	N3A-C2A-N1A	-13.83	118.00	128.87
2	B	300	NDP	N3A-C2A-N1A	-13.81	118.03	128.87
2	B	300	NDP	C4B-O4B-C1B	-12.71	96.17	109.64
2	A	300	NDP	C4B-O4B-C1B	-12.68	96.21	109.64
2	B	300	NDP	C1B-N9A-C4A	-9.18	116.56	126.81
2	A	300	NDP	C1B-N9A-C4A	-9.18	116.56	126.81
2	B	300	NDP	C3B-C2B-C1B	-5.10	92.86	102.63
2	A	300	NDP	C3B-C2B-C1B	-5.10	92.87	102.63
2	A	300	NDP	O4B-C1B-N9A	-3.99	100.56	108.11
2	B	300	NDP	O4B-C1B-N9A	-3.99	100.57	108.11
2	A	300	NDP	O2B-P2B-O1X	-3.32	99.55	107.48
2	B	300	NDP	O2B-P2B-O1X	-3.31	99.58	107.48
2	B	300	NDP	O4D-C1D-C2D	-3.06	99.69	106.61
2	A	300	NDP	O4D-C1D-C2D	-3.05	99.70	106.61

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	NDP	C3N-C2N-N1N	-2.16	120.08	123.24
2	A	300	NDP	C5B-C4B-C3B	-2.11	107.05	115.20
2	B	300	NDP	C5B-C4B-C3B	-2.10	107.06	115.20
2	B	300	NDP	C3N-C2N-N1N	-2.10	120.16	123.24
2	B	300	NDP	O3X-P2B-O2X	2.26	115.73	107.44
2	A	300	NDP	O3X-P2B-O2X	2.26	115.76	107.44
2	B	300	NDP	O2N-PN-O3	2.34	115.31	105.27
2	A	300	NDP	O2N-PN-O3	2.37	115.42	105.27
2	A	300	NDP	N6A-C6A-N1A	2.76	123.15	118.52
2	B	300	NDP	N6A-C6A-N1A	2.80	123.21	118.52
2	A	300	NDP	O4D-C1D-N1N	2.90	113.63	108.09
2	B	300	NDP	O4D-C1D-N1N	2.94	113.69	108.09
2	A	300	NDP	C5N-C4N-C3N	3.40	121.22	112.41
2	B	300	NDP	C5N-C4N-C3N	3.43	121.28	112.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	NDP	15	0
2	B	300	NDP	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	234/270 (86%)	-0.07	8 (3%) 49 58	21, 30, 53, 79	0
1	B	243/270 (90%)	0.17	15 (6%) 24 32	24, 35, 65, 97	0
All	All	477/540 (88%)	0.05	23 (4%) 34 43	21, 33, 61, 97	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	6.9
1	B	209	LEU	6.2
1	B	206	ILE	5.9
1	B	205	ARG	5.0
1	B	196	PHE	5.0
1	B	192	MET	4.8
1	B	215	ILE	4.2
1	B	207	ALA	3.4
1	A	209	LEU	3.2
1	A	187	PHE	3.1
1	B	187	PHE	3.1
1	A	87	VAL	2.8
1	B	194	ARG	2.8
1	A	1	MET	2.7
1	A	189	GLU	2.7
1	B	197	LEU	2.6
1	A	192	MET	2.4
1	A	215	ILE	2.3
1	B	2	GLU	2.2
1	B	191	ASP	2.2
1	B	75	ARG	2.0
1	B	193	THR	2.0
1	A	210	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NDP	A	300	48/48	0.40	0.71	11.45	20,27,30,33	0
2	NDP	B	300	48/48	0.56	0.56	6.64	20,27,30,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.