



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:02 PM GMT

PDB ID : 4B9K
Title : pVHL-ELOB-ELOC complex_(2S,4R)-1-(3-amino-2-methylbenzoyl)-4-hydroxy-N-(4-(4-methylthiazol-5-yl)benzyl)pyrrolidine-2-carboxamide bound
Authors : Buckley, D.L.; Gustafson, J.L.; VanMolle, I.; Roth, A.G.; SeopTae, H.; Gareiss, P.C.; Jorgensen, W.L.; Ciulli, A.; Crews, C.M.
Deposited on : 2012-09-05
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	 96%
1	D	104	 94%
1	G	104	 96%
1	J	104	 93% 6%
2	B	97	 78% 8% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	97	
2	K	97	
3	C	171	
3	F	171	
3	I	171	
3	L	171	
4	E	97	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ACT	H	1114	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 11711 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	103	Total	As	C	N	O	S	0	1	0
			811	3	516	137	150	5			
1	D	101	Total	As	C	N	O	S	0	2	0
			803	3	509	137	150	4			
1	G	103	Total	As	C	N	O	S	0	1	0
			827	2	523	140	157	5			
1	J	103	Total	As	C	N	O	S	0	0	0
			810	2	515	136	152	5			

- Molecule 2 is a protein called TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	88	Total	C	N	O	S	0	1	0
			698	452	113	127	6			
2	H	89	Total	C	N	O	S	0	1	0
			693	449	110	128	6			
2	K	91	Total	C	N	O	S	0	0	0
			714	461	112	134	7			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	MET	-	EXPRESSION TAG	UNP Q15369
H	16	MET	-	EXPRESSION TAG	UNP Q15369
K	16	MET	-	EXPRESSION TAG	UNP Q15369

- Molecule 3 is a protein called VON HIPPEL-LINDAU DISEASE TUMOR SUPPRESSOR.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	138	Total	As	C	N	O	S	0	2	0
			1119	1	717	198	201	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	F	143	Total	As	C	N	O	S	0	2	0
			1157	1	738	206	210	2			
3	I	147	Total	As	C	N	O	S	0	1	0
			1172	1	751	206	212	2			
3	L	146	Total	As	C	N	O	S	0	0	0
			1164	1	742	212	207	2			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	43	SER	-	EXPRESSION TAG	UNP P40337
C	44	MET	-	EXPRESSION TAG	UNP P40337
C	45	SER	-	EXPRESSION TAG	UNP P40337
C	46	GLU	-	EXPRESSION TAG	UNP P40337
C	47	ASN	-	EXPRESSION TAG	UNP P40337
C	48	LEU	-	EXPRESSION TAG	UNP P40337
C	49	TYR	-	EXPRESSION TAG	UNP P40337
C	50	PHE	-	EXPRESSION TAG	UNP P40337
C	51	GLN	-	EXPRESSION TAG	UNP P40337
C	52	GLY	-	EXPRESSION TAG	UNP P40337
C	53	SER	-	EXPRESSION TAG	UNP P40337
F	43	SER	-	EXPRESSION TAG	UNP P40337
F	44	MET	-	EXPRESSION TAG	UNP P40337
F	45	SER	-	EXPRESSION TAG	UNP P40337
F	46	GLU	-	EXPRESSION TAG	UNP P40337
F	47	ASN	-	EXPRESSION TAG	UNP P40337
F	48	LEU	-	EXPRESSION TAG	UNP P40337
F	49	TYR	-	EXPRESSION TAG	UNP P40337
F	50	PHE	-	EXPRESSION TAG	UNP P40337
F	51	GLN	-	EXPRESSION TAG	UNP P40337
F	52	GLY	-	EXPRESSION TAG	UNP P40337
F	53	SER	-	EXPRESSION TAG	UNP P40337
I	43	SER	-	EXPRESSION TAG	UNP P40337
I	44	MET	-	EXPRESSION TAG	UNP P40337
I	45	SER	-	EXPRESSION TAG	UNP P40337
I	46	GLU	-	EXPRESSION TAG	UNP P40337
I	47	ASN	-	EXPRESSION TAG	UNP P40337
I	48	LEU	-	EXPRESSION TAG	UNP P40337
I	49	TYR	-	EXPRESSION TAG	UNP P40337
I	50	PHE	-	EXPRESSION TAG	UNP P40337
I	51	GLN	-	EXPRESSION TAG	UNP P40337
I	52	GLY	-	EXPRESSION TAG	UNP P40337

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	53	SER	-	EXPRESSION TAG	UNP P40337
L	43	SER	-	EXPRESSION TAG	UNP P40337
L	44	MET	-	EXPRESSION TAG	UNP P40337
L	45	SER	-	EXPRESSION TAG	UNP P40337
L	46	GLU	-	EXPRESSION TAG	UNP P40337
L	47	ASN	-	EXPRESSION TAG	UNP P40337
L	48	LEU	-	EXPRESSION TAG	UNP P40337
L	49	TYR	-	EXPRESSION TAG	UNP P40337
L	50	PHE	-	EXPRESSION TAG	UNP P40337
L	51	GLN	-	EXPRESSION TAG	UNP P40337
L	52	GLY	-	EXPRESSION TAG	UNP P40337
L	53	SER	-	EXPRESSION TAG	UNP P40337


- Molecule 4 is a protein called TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	E	87	Total	As	C	N	O	S	0	0	0
			690	1	446	109	128	6			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	16	MET	-	EXPRESSION TAG	UNP Q15369

- Molecule 5 is (2S,4R)-1-(3-AMINO-2-METHYLBENZOYL)-4-HYDROXY-N-(4-(4-METHYLTHIAZOL-5-YL)BENZYL)PYRROLIDINE-2-CARBOXAMIDE (three-letter code: TG0) (formula: C₂₄H₂₆N₄O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	S	0	0
			32	24	4	3	1		
5	F	1	Total	C	N	O	S	0	0
			32	24	4	3	1		
5	I	1	Total	C	N	O	S	0	0
			32	24	4	3	1		
5	L	1	Total	C	N	O	S	0	0
			32	24	4	3	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	F	1	Total C O 4 2 2	0	0
6	F	1	Total C O 4 2 2	0	0
6	H	1	Total C O 4 2 2	0	0
6	H	1	Total C O 4 2 2	0	0
6	I	1	Total C O 4 2 2	0	0
6	K	1	Total C O 4 2 2	0	0
6	L	1	Total C O 4 2 2	0	0
6	L	1	Total C O 4 2 2	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	96	Total O 96 96	0	0
7	B	41	Total O 41 41	0	0
7	C	85	Total O 85 85	0	0
7	D	58	Total O 58 58	0	0
7	E	33	Total O 33 33	0	0
7	F	82	Total O 82 82	0	0
7	G	82	Total O 82 82	0	0
7	H	60	Total O 60 60	0	0
7	I	93	Total O 93 93	0	0
7	J	105	Total O 105 105	0	0
7	K	58	Total O 58 58	0	0
7	L	100	Total O 100 100	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2

Chain A:  96%



- Molecule 1: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2

Chain D:  94%




- Molecule 1: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2

Chain G:  96%





- Molecule 1: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 2

Chain J:  93% 6%





- Molecule 2: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1

Chain B:  78% 8% 9%





- Molecule 2: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1


Chain H:  81% 9% 8%




- Molecule 2: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1




- Molecule 3: VON HIPPEL-LINDAU DISEASE TUMOR SUPPRESSOR




- Molecule 3: VON HIPPEL-LINDAU DISEASE TUMOR SUPPRESSOR




- Molecule 3: VON HIPPEL-LINDAU DISEASE TUMOR SUPPRESSOR



- Molecule 3: VON HIPPEL-LINDAU DISEASE TUMOR SUPPRESSOR



- Molecule 4: TRANSCRIPTION ELONGATION FACTOR B POLYPEPTIDE 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.86Å 92.86Å 364.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.06 – 2.00 46.06 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.06-2.00) 100.0 (46.06-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.00Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.189 , 0.220 0.199 , 0.233	Depositor DCC
R_{free} test set	5443 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.452	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	14 of 108859 reflections (0.013%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11711	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.64 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.4981e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, TG0, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/797	0.67	0/1073
1	D	0.44	0/788	0.69	1/1063 (0.1%)
1	G	0.47	0/823	0.69	0/1110
1	J	0.50	0/806	0.67	0/1087
2	B	0.49	0/712	0.69	0/961
2	H	0.53	0/707	0.63	0/956
2	K	0.54	0/729	0.71	0/984
3	C	0.46	0/1138	0.68	1/1555 (0.1%)
3	F	0.46	0/1177	0.67	0/1609
3	I	0.47	0/1192	0.65	0/1629
3	L	0.48	0/1184	0.70	0/1618
4	E	0.48	0/694	0.64	0/938
All	All	0.48	0/10747	0.67	2/14583 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	103	PRO	N-CA-C	5.39	126.12	112.10
1	D	100	PRO	N-CA-CB	5.06	109.37	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	811	0	792	1	0
1	D	803	0	766	0	0
1	G	827	0	819	1	0
1	J	810	0	802	5	0
2	B	698	0	683	6	0
2	H	693	0	685	4	0
2	K	714	0	706	10	0
3	C	1119	0	1081	8	0
3	F	1157	0	1115	5	0
3	I	1172	0	1131	6	0
3	L	1164	0	1136	8	0
4	E	690	0	677	10	0
5	C	32	0	26	0	0
5	F	32	0	26	0	0
5	I	32	0	26	0	0
5	L	32	0	26	0	0
6	F	8	0	6	1	0
6	H	8	0	6	0	0
6	I	4	0	3	0	0
6	K	4	0	3	0	0
6	L	8	0	6	1	0
7	A	96	0	0	0	0
7	B	41	0	0	0	0
7	C	85	0	0	0	0
7	D	58	0	0	0	0
7	E	33	0	0	0	0
7	F	82	0	0	0	0
7	G	82	0	0	0	0
7	H	60	0	0	1	0
7	I	93	0	0	2	0
7	J	105	0	0	1	0
7	K	58	0	0	0	0
7	L	100	0	0	0	0
All	All	11711	0	10521	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:77:PHE:HE1	4:E:112:CAS:AS	2.35	0.69
4:E:57:THR:HA	4:E:59:GLU:H	1.61	0.65
4:E:77:PHE:CE1	4:E:112:CAS:AS	3.11	0.64
3:L:91:PHE:H	6:L:1210:ACT:H2	1.65	0.60
3:C:120[B]:ARG:NH1	3:C:197:ASP:OD2	2.28	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/104 (93%)	94 (97%)	3 (3%)	0	100	100
1	D	96/104 (92%)	92 (96%)	3 (3%)	1 (1%)	19	11
1	G	100/104 (96%)	97 (97%)	3 (3%)	0	100	100
1	J	99/104 (95%)	96 (97%)	3 (3%)	0	100	100
2	B	85/97 (88%)	80 (94%)	2 (2%)	3 (4%)	4	1
2	H	86/97 (89%)	84 (98%)	2 (2%)	0	100	100
2	K	87/97 (90%)	83 (95%)	2 (2%)	2 (2%)	8	3
3	C	135/171 (79%)	130 (96%)	4 (3%)	1 (1%)	26	19
3	F	142/171 (83%)	138 (97%)	4 (3%)	0	100	100
3	I	145/171 (85%)	140 (97%)	4 (3%)	1 (1%)	26	19
3	L	143/171 (84%)	134 (94%)	6 (4%)	3 (2%)	9	3
4	E	83/97 (86%)	82 (99%)	1 (1%)	0	100	100
All	All	1298/1488 (87%)	1250 (96%)	37 (3%)	11 (1%)	21	15

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	142	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	180	ILE
2	B	89	GLU
3	I	206	ILE
2	K	88	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/90 (92%)	82 (99%)	1 (1%)	78	81
1	D	80/90 (89%)	79 (99%)	1 (1%)	76	79
1	G	89/90 (99%)	88 (99%)	1 (1%)	80	83
1	J	86/90 (96%)	86 (100%)	0	100	100
2	B	74/86 (86%)	72 (97%)	2 (3%)	52	52
2	H	76/86 (88%)	70 (92%)	6 (8%)	15	9
2	K	79/86 (92%)	76 (96%)	3 (4%)	40	36
3	C	121/156 (78%)	120 (99%)	1 (1%)	86	89
3	F	125/156 (80%)	123 (98%)	2 (2%)	70	73
3	I	125/156 (80%)	123 (98%)	2 (2%)	70	73
3	L	126/156 (81%)	124 (98%)	2 (2%)	70	73
4	E	75/85 (88%)	72 (96%)	3 (4%)	38	33
All	All	1139/1327 (86%)	1115 (98%)	24 (2%)	63	63

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	3	VAL
2	H	61	ASN
3	L	63	LEU
2	H	38	THR
2	H	42	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	10	HIS
3	I	110	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

15 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	60	1	5,8,9	0.74	0	2,9,11	2.74	2 (100%)
1	CAS	A	89[A]	1	5,8,9	1.13	0	2,9,11	1.50	1 (50%)
1	CAS	A	89[B]	1	5,8,9	1.23	0	2,9,11	2.18	2 (100%)
3	CAS	C	77	3	5,8,9	0.99	0	2,9,11	1.57	1 (50%)
1	CAS	D	60	1	5,8,9	0.54	0	2,9,11	2.22	1 (50%)
1	CAS	D	89[A]	1	5,8,9	1.20	1 (20%)	2,9,11	1.57	1 (50%)
1	CAS	D	89[B]	1	5,8,9	0.97	0	2,9,11	2.24	2 (100%)
4	CAS	E	112	4	4,9,9	0.95	0	1,11,11	0.32	0
3	CAS	F	77	3	5,8,9	1.18	1 (20%)	2,9,11	1.61	1 (50%)
1	CAS	G	60	1	5,8,9	0.93	0	2,9,11	2.15	2 (100%)
1	CAS	G	89	1	5,8,9	1.01	0	2,9,11	1.83	1 (50%)
3	CAS	I	77	3	5,8,9	0.64	0	2,9,11	1.78	1 (50%)
1	CAS	J	60	1	5,8,9	0.70	0	2,9,11	2.24	2 (100%)
1	CAS	J	89	1	5,8,9	1.04	0	2,9,11	1.55	1 (50%)
3	CAS	L	77	3	5,8,9	0.68	0	2,9,11	1.67	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	60	1	-	0/0/7/9	0/0/0/0
1	CAS	A	89[A]	1	-	0/0/7/9	0/0/0/0
1	CAS	A	89[B]	1	-	0/0/7/9	0/0/0/0
3	CAS	C	77	3	-	0/0/7/9	0/0/0/0
1	CAS	D	60	1	-	0/0/7/9	0/0/0/0
1	CAS	D	89[A]	1	-	0/0/7/9	0/0/0/0
1	CAS	D	89[B]	1	-	0/0/7/9	0/0/0/0
4	CAS	E	112	4	-	0/2/9/9	0/0/0/0
3	CAS	F	77	3	-	0/0/7/9	0/0/0/0
1	CAS	G	60	1	-	0/0/7/9	0/0/0/0
1	CAS	G	89	1	-	0/0/7/9	0/0/0/0
3	CAS	I	77	3	-	0/0/7/9	0/0/0/0
1	CAS	J	60	1	-	0/0/7/9	0/0/0/0
1	CAS	J	89	1	-	0/0/7/9	0/0/0/0
3	CAS	L	77	3	-	0/0/7/9	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	89[A]	CAS	AS-CE1	2.13	2.01	1.96
3	F	77	CAS	AS-CE2	2.46	2.02	1.96

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	60	CAS	CA-CB-SG	-3.04	103.34	114.16
3	I	77	CAS	O-C-CA	-2.51	118.95	125.49
1	D	60	CAS	O-C-CA	-2.51	118.96	125.49
1	A	60	CAS	O-C-CA	-2.41	119.22	125.49
3	L	77	CAS	O-C-CA	-2.36	119.35	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	112	CAS	4	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	TG0	C	1203	-	31,35,35	1.90	6 (19%)	36,50,50	1.39	3 (8%)
6	ACT	F	1205	-	1,3,3	4.24	1 (100%)	0,3,3	0.00	-
5	TG0	F	1206	-	31,35,35	1.95	6 (19%)	36,50,50	1.45	6 (16%)
6	ACT	F	1207	-	1,3,3	4.36	1 (100%)	0,3,3	0.00	-
6	ACT	H	1113	-	1,3,3	4.58	1 (100%)	0,3,3	0.00	-
6	ACT	H	1114	-	1,3,3	3.91	1 (100%)	0,3,3	0.00	-
6	ACT	I	1209	-	1,3,3	4.66	1 (100%)	0,3,3	0.00	-
5	TG0	I	1210	-	31,35,35	2.04	6 (19%)	36,50,50	1.35	4 (11%)
6	ACT	K	1113	-	1,3,3	3.96	1 (100%)	0,3,3	0.00	-
6	ACT	L	1208	-	1,3,3	5.07	1 (100%)	0,3,3	0.00	-
5	TG0	L	1209	-	31,35,35	1.83	5 (16%)	36,50,50	1.46	6 (16%)
6	ACT	L	1210	-	1,3,3	2.18	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TG0	C	1203	-	-	0/21/33/33	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ACT	F	1205	-	-	0/0/0/0	0/0/0/0
5	TG0	F	1206	-	-	0/21/33/33	0/4/4/4
6	ACT	F	1207	-	-	0/0/0/0	0/0/0/0
6	ACT	H	1113	-	-	0/0/0/0	0/0/0/0
6	ACT	H	1114	-	-	0/0/0/0	0/0/0/0
6	ACT	I	1209	-	-	0/0/0/0	0/0/0/0
5	TG0	I	1210	-	-	0/21/33/33	0/4/4/4
6	ACT	K	1113	-	-	0/0/0/0	0/0/0/0
6	ACT	L	1208	-	-	0/0/0/0	0/0/0/0
5	TG0	L	1209	-	-	0/21/33/33	0/4/4/4
6	ACT	L	1210	-	-	0/0/0/0	0/0/0/0

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1203	TG0	C25-C26	-2.23	1.47	1.52
5	L	1209	TG0	C25-C08	-2.15	1.49	1.53
5	I	1210	TG0	C25-C26	-2.08	1.48	1.52
5	C	1203	TG0	C02-N01	2.06	1.44	1.38
5	F	1206	TG0	C13-C12	2.15	1.43	1.38

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1203	TG0	C11-N10-C09	-3.94	116.92	122.34
5	F	1206	TG0	C11-N10-C09	-3.12	118.05	122.34
5	L	1209	TG0	C12-C11-N10	-2.81	106.46	112.88
5	F	1206	TG0	C12-C11-N10	-2.37	107.48	112.88
5	I	1210	TG0	C28-N07-C08	-2.32	108.07	111.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	F	1205	ACT	1	0
6	L	1210	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	101/104 (97%)	-0.26	0 100 100	27, 38, 65, 85	0
1	D	99/104 (95%)	0.08	1 (1%) 84 84	30, 50, 86, 112	0
1	G	101/104 (97%)	-0.19	0 100 100	28, 42, 62, 73	0
1	J	101/104 (97%)	-0.28	0 100 100	24, 33, 56, 70	0
2	B	88/97 (90%)	-0.05	2 (2%) 64 64	26, 40, 72, 86	0
2	H	89/97 (91%)	-0.05	2 (2%) 65 66	26, 37, 62, 82	0
2	K	91/97 (93%)	0.16	4 (4%) 38 39	24, 36, 68, 83	0
3	C	137/171 (80%)	-0.12	1 (0%) 89 89	27, 38, 61, 107	0
3	F	142/171 (83%)	-0.05	1 (0%) 89 89	26, 39, 65, 117	0
3	I	146/171 (85%)	0.01	3 (2%) 67 67	28, 41, 61, 129	0
3	L	145/171 (84%)	0.03	4 (2%) 56 57	23, 37, 74, 121	0
4	E	86/97 (88%)	0.08	4 (4%) 35 37	27, 45, 71, 85	0
All	All	1326/1488 (89%)	-0.05	22 (1%) 73 73	23, 40, 71, 129	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	K	48	GLY	6.3
2	K	49	PRO	5.1
3	F	201	LEU	4.4
1	D	103	MET	3.7
3	L	143	ASP	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CAS	G	89	9/10	0.66	0.20	-	32,39,59,60	3
4	CAS	E	112	10/10	0.90	0.20	-	45,55,58,63	3
3	CAS	L	77	9/10	0.94	0.13	-	26,29,43,45	3
1	CAS	A	89[A]	9/10	0.92	0.15	-	43,46,62,63	9
3	CAS	F	77	9/10	0.96	0.10	-	29,32,45,47	3
1	CAS	G	60	9/10	0.95	0.12	-	38,40,47,49	3
1	CAS	D	60	9/10	0.96	0.10	-	36,38,49,50	3
3	CAS	C	77	9/10	0.92	0.15	-	33,38,51,54	3
1	CAS	A	89[B]	9/10	0.92	0.15	-	44,46,60,62	9
1	CAS	J	60	9/10	0.92	0.13	-	34,38,48,49	3
1	CAS	A	60	9/10	0.94	0.12	-	31,34,42,42	3
3	CAS	I	77	9/10	0.97	0.11	-	31,35,49,49	3
1	CAS	D	89[A]	9/10	0.92	0.14	-	60,62,76,76	9
1	CAS	D	89[B]	9/10	0.92	0.14	-	60,62,74,78	9
1	CAS	J	89	9/10	0.96	0.13	-	32,34,44,45	3

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	ACT	H	1114	4/4	0.78	0.30	4.34	64,64,64,65	0
6	ACT	K	1113	4/4	0.91	0.13	1.93	52,54,55,61	0
6	ACT	I	1209	4/4	0.86	0.14	1.19	53,65,66,67	0
6	ACT	F	1207	4/4	0.90	0.16	0.68	75,76,76,76	0
6	ACT	H	1113	4/4	0.84	0.13	0.52	61,61,63,64	0
5	TG0	L	1209	32/32	0.97	0.11	0.15	23,28,31,31	0
5	TG0	I	1210	32/32	0.95	0.12	0.08	31,35,40,42	0
5	TG0	C	1203	32/32	0.96	0.11	-0.09	23,32,42,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	TG0	F	1206	32/32	0.97	0.10	-0.81	24,31,35,36	0
6	ACT	L	1208	4/4	0.41	0.35	-	59,60,61,62	0
6	ACT	L	1210	4/4	0.74	0.28	-	43,50,55,59	0
6	ACT	F	1205	4/4	0.67	0.20	-	66,69,69,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.