



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 02:19 PM BST

PDB ID : 1BA9
Title : THE SOLUTION STRUCTURE OF REDUCED MONOMERIC SUPEROXIDE DISMUTASE, NMR, 36 STRUCTURES
Authors : Banci, L.; Benedetto, M.; Bertini, I.; Del Conte, R.; Piccioli, M.; Viezzoli, M.S.
Deposited on : 1998-04-24

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	153	 72% 13% 15%

2 Ensemble composition and analysis

This entry contains 36 models. Model 30 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:21, A:28-A:49, A:62-A:106, A:110-A:153 (130)	0.34	30

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 7, 8, 9, 11, 12, 14, 24, 27, 28, 30, 31, 32, 33, 34, 35, 36
2	4, 10, 22, 25
3	13, 15, 21, 29
4	17, 18
Single-model clusters	16; 19; 20; 23; 26

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2194 atoms, of which 1080 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called SUPEROXIDE DISMUTASE.

Mol	Chain	Residues	Atoms						Trace
1	A	153	Total	C	H	N	O	S	0
			2192	678	1080	204	228	2	

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	ALA	CYS	ENGINEERED	UNP P00441
A	50	GLU	PHE	ENGINEERED	UNP P00441
A	51	GLU	GLY	ENGINEERED	UNP P00441
A	111	SER	CYS	ENGINEERED	UNP P00441
A	133	GLN	GLU	ENGINEERED	UNP P00441

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

- Molecule 3 is COPPER (I) ION (three-letter code: CU1) (formula: Cu).


Mol	Chain	Residues	Atoms	
3	A	1	Total	Cu
			1	1

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: SUPEROXIDE DISMUTASE




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: SUPEROXIDE DISMUTASE



4.2.2 Score per residue for model 2

- Molecule 1: SUPEROXIDE DISMUTASE






4.2.3 Score per residue for model 3

- Molecule 1: SUPEROXIDE DISMUTASE

Chain A: 69% 14% 15%



4.2.4 Score per residue for model 4

- Molecule 1: SUPEROXIDE DISMUTASE

Chain A: 67% 17% 15%



4.2.5 Score per residue for model 5


- Molecule 1: SUPEROXIDE DISMUTASE

Chain A: 65% 19% 15%




4.2.6 Score per residue for model 6

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.7 Score per residue for model 7

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.8 Score per residue for model 8

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.9 Score per residue for model 9

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.14 Score per residue for model 14

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.15 Score per residue for model 15

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.16 Score per residue for model 16

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.17 Score per residue for model 17

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.18 Score per residue for model 18

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.19 Score per residue for model 19

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.20 Score per residue for model 20

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.21 Score per residue for model 21

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.22 Score per residue for model 22

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.23 Score per residue for model 23

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.24 Score per residue for model 24

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.25 Score per residue for model 25

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.26 Score per residue for model 26

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.27 Score per residue for model 27

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.28 Score per residue for model 28

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.29 Score per residue for model 29

• Molecule 1: SUPEROXIDE DISMUTASE




4.2.30 Score per residue for model 30 (medoid)

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.31 Score per residue for model 31

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.32 Score per residue for model 32

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.33 Score per residue for model 33

- Molecule 1: SUPEROXIDE DISMUTASE




4.2.34 Score per residue for model 34


- Molecule 1: SUPEROXIDE DISMUTASE



4.2.35 Score per residue for model 35

- Molecule 1: SUPEROXIDE DISMUTASE


Chain A:  73% 10% 15%



4.2.36 Score per residue for model 36

- Molecule 1: SUPEROXIDE DISMUTASE

Chain A:  73% 10% 15%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *TORTION ANGLES DYNAMIC (DYANA)*.

Of the 36 calculated structures, 36 were deposited, based on the following criterion: *TARGET FUNCTION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	4.0
XEASY (ETH	structure solution	(ETH
ZURICH)	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4202
Number of chemical shift lists	1
Total number of shifts	1695
Number of shifts mapped to atoms	1695
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.50±0.00	0±0/976 (0.0±0.0%)	0.94±0.02	0±1/1314 (0.0±0.0%)
All	All	0.50	0/35136 (0.0%)	0.94	15/47304 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.2	0.8±0.7
All	All	1	28

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	79	ARG	NE-CZ-NH2	-6.83	116.89	120.30	17	1
1	A	124	ASP	CB-CG-OD2	-6.08	112.82	118.30	5	8
1	A	79	ARG	NE-CZ-NH1	5.51	123.06	120.30	17	1
1	A	69	ARG	NE-CZ-NH2	5.28	122.94	120.30	10	1
1	A	115	ARG	NE-CZ-NH2	-5.27	117.66	120.30	24	2
1	A	8	LEU	CB-CA-C	5.15	119.99	110.20	29	1
1	A	115	ARG	NE-CZ-NH1	5.03	122.81	120.30	24	1

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	62	PRO	CA	1

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	80	HIS	Sidechain,Peptide	11
1	A	71	HIS	Sidechain	8
1	A	115	ARG	Sidechain	4
1	A	45	PHE	Sidechain	2
1	A	64	PHE	Sidechain	1
1	A	79	ARG	Sidechain	1
1	A	62	PRO	Peptide	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	959	952	951	3±2
All	All	34596	34272	34236	111

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:35:ILE:HG21	1:A:38:LEU:HD23	0.71	1.61	1	12
1:A:38:LEU:HD12	1:A:43:HIS:CE1	0.63	2.27	2	14
1:A:14:VAL:HG23	1:A:38:LEU:HD22	0.62	1.70	2	1
1:A:117:LEU:CD1	1:A:149:ILE:HD11	0.62	2.24	22	1
1:A:117:LEU:HD13	1:A:149:ILE:HD11	0.60	1.74	22	1
1:A:42:LEU:HG	1:A:123:ALA:HB2	0.57	1.76	4	3
1:A:35:ILE:HG22	1:A:95:ALA:H	0.56	1.60	17	6
1:A:8:LEU:HD11	1:A:117:LEU:HD23	0.53	1.80	24	1
1:A:112:ILE:HG12	1:A:149:ILE:HG21	0.51	1.82	9	1
1:A:14:VAL:HG21	1:A:144:LEU:HB3	0.51	1.82	2	1
1:A:81:VAL:HG13	1:A:104:ILE:HG22	0.50	1.82	16	1
1:A:131:ASN:ND2	1:A:134:SER:H	0.49	2.06	23	4
1:A:45:PHE:CE1	1:A:117:LEU:HD12	0.49	2.42	16	1
1:A:115:ARG:HG2	1:A:149:ILE:HD12	0.48	1.86	13	1
1:A:115:ARG:HB2	1:A:149:ILE:HD12	0.47	1.85	11	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:70:LYS:NZ	1:A:78:GLU:OE2	0.47	2.47	6	1
1:A:46:HIS:CE1	1:A:63:HIS:CE1	0.46	3.03	19	2
1:A:115:ARG:CG	1:A:149:ILE:HD12	0.46	2.40	13	1
1:A:38:LEU:HD23	1:A:89:ALA:HB2	0.46	1.87	16	2
1:A:4:ALA:HB3	1:A:20:PHE:HB2	0.46	1.88	29	1
1:A:35:ILE:HG21	1:A:38:LEU:HD22	0.46	1.86	11	1
1:A:35:ILE:CG2	1:A:38:LEU:HD23	0.45	2.41	2	2
1:A:35:ILE:O	1:A:94:VAL:HG13	0.45	2.11	16	1
1:A:45:PHE:CZ	1:A:117:LEU:HD22	0.45	2.46	2	1
1:A:46:HIS:O	1:A:117:LEU:HD23	0.45	2.12	2	1
1:A:117:LEU:HD13	1:A:118:VAL:N	0.45	2.26	28	1
1:A:20:PHE:CE1	1:A:31:VAL:HG22	0.45	2.47	27	1
1:A:117:LEU:HD12	1:A:149:ILE:HD11	0.45	1.87	3	1
1:A:6:ALA:HB3	1:A:18:ILE:HB	0.44	1.89	10	2
1:A:91:LYS:NZ	1:A:92:ASP:OD1	0.44	2.50	14	1
1:A:73:GLY:HA3	1:A:126:LEU:HD22	0.43	1.90	9	1
1:A:84:LEU:HD22	1:A:84:LEU:N	0.43	2.28	23	1
1:A:38:LEU:HD23	1:A:43:HIS:CG	0.43	2.48	13	1
1:A:76:ASP:OD1	1:A:128:LYS:NZ	0.43	2.51	17	1
1:A:36:LYS:NZ	1:A:92:ASP:OD2	0.43	2.52	34	7
1:A:43:HIS:CE1	1:A:121:GLU:O	0.42	2.72	20	2
1:A:3:LYS:NZ	1:A:21:GLU:OE2	0.42	2.52	23	1
1:A:69:ARG:NH2	1:A:77:GLU:OE2	0.42	2.53	16	1
1:A:115:ARG:O	1:A:149:ILE:HD12	0.42	2.14	17	1
1:A:30:LYS:NZ	1:A:100:GLU:OE2	0.42	2.46	27	2
1:A:35:ILE:HG21	1:A:38:LEU:HB2	0.42	1.92	23	1
1:A:65:ASN:HD21	1:A:80:HIS:CD2	0.42	2.33	34	7
1:A:65:ASN:N	1:A:66:PRO:CD	0.42	2.83	28	1
1:A:17:ILE:N	1:A:17:ILE:HD12	0.42	2.30	10	1
1:A:46:HIS:CE1	1:A:120:HIS:CD2	0.41	3.08	18	3
1:A:125:ASP:OD2	1:A:128:LYS:NZ	0.41	2.51	26	1
1:A:4:ALA:HB2	1:A:113:ILE:CD1	0.41	2.45	5	1
1:A:38:LEU:HD21	1:A:43:HIS:CG	0.41	2.50	23	1
1:A:76:ASP:OD2	1:A:128:LYS:NZ	0.41	2.53	18	1
1:A:8:LEU:CD1	1:A:117:LEU:HD23	0.41	2.46	12	1
1:A:46:HIS:NE2	1:A:124:ASP:OD2	0.41	2.54	21	1
1:A:38:LEU:HD12	1:A:43:HIS:NE2	0.41	2.31	10	1
1:A:45:PHE:CE1	1:A:117:LEU:HD22	0.41	2.51	2	1
1:A:4:ALA:HB2	1:A:113:ILE:HD11	0.40	1.93	17	1
1:A:119:VAL:CG1	1:A:145:ALA:HB3	0.40	2.46	16	1
1:A:117:LEU:HD12	1:A:149:ILE:CD1	0.40	2.47	19	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/153 (84%)	109±3 (84±2%)	16±3 (12±2%)	5±2 (4±1%)	7	36
All	All	4644/5508 (84%)	3909 (84%)	568 (12%)	167 (4%)	7	36

All 26 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	85	GLY	29
1	A	111	SER	23
1	A	14	VAL	18
1	A	79	ARG	18
1	A	132	GLU	10
1	A	66	PRO	10
1	A	133	GLN	9
1	A	80	HIS	8
1	A	120	HIS	7
1	A	49	GLU	7
1	A	3	LYS	4
1	A	81	VAL	3
1	A	110	HIS	3
1	A	68	SER	2
1	A	87	VAL	2
1	A	37	GLY	2
1	A	38	LEU	2
1	A	67	LEU	2
1	A	8	LEU	1
1	A	125	ASP	1
1	A	65	ASN	1
1	A	134	SER	1
1	A	9	LYS	1
1	A	104	ILE	1
1	A	105	SER	1
1	A	62	PRO	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/118 (86%)	85±4 (83±4%)	17±4 (17±4%)	6	43
All	All	3672/4248 (86%)	3065 (83%)	607 (17%)	6	43

All 71 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	35	ILE	36
1	A	133	GLN	36
1	A	131	ASN	33
1	A	92	ASP	30
1	A	45	PHE	28
1	A	144	LEU	27
1	A	91	LYS	25
1	A	46	HIS	23
1	A	64	PHE	23
1	A	143	ARG	22
1	A	3	LYS	14
1	A	49	GLU	14
1	A	128	LYS	14
1	A	31	VAL	14
1	A	117	LEU	13
1	A	146	CYS	13
1	A	112	ILE	11
1	A	67	LEU	10
1	A	86	ASN	9
1	A	124	ASP	9
1	A	90	ASP	8
1	A	83	ASP	8
1	A	81	VAL	8
1	A	29	VAL	8
1	A	79	ARG	8
1	A	121	GLU	8
1	A	9	LYS	7
1	A	75	LYS	7
1	A	40	GLU	7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	69	ARG	7
1	A	19	ASN	7
1	A	106	LEU	6
1	A	77	GLU	6
1	A	32	TRP	5
1	A	21	GLU	5
1	A	104	ILE	5
1	A	36	LYS	5
1	A	11	ASP	4
1	A	101	ASP	4
1	A	134	SER	4
1	A	115	ARG	4
1	A	15	GLN	4
1	A	139	ASN	4
1	A	137	THR	4
1	A	126	LEU	4
1	A	119	VAL	4
1	A	153	GLN	4
1	A	98	SER	4
1	A	78	GLU	3
1	A	125	ASP	3
1	A	105	SER	3
1	A	42	LEU	3
1	A	30	LYS	3
1	A	100	GLU	3
1	A	102	SER	2
1	A	38	LEU	2
1	A	135	THR	2
1	A	47	VAL	2
1	A	122	LYS	2
1	A	110	HIS	2
1	A	111	SER	2
1	A	103	VAL	2
1	A	116	THR	2
1	A	68	SER	1
1	A	96	ASP	1
1	A	132	GLU	1
1	A	70	LYS	1
1	A	63	HIS	1
1	A	149	ILE	1
1	A	17	ILE	1
1	A	142	SER	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 88% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 4202

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1695
Number of shifts mapped to atoms	1695
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	8

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	151	0.01 ± 0.11	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	126	0.20 ± 0.19	None needed (< 0.5 ppm)
$^{13}\text{C}'$	150	0.28 ± 0.16	None needed (< 0.5 ppm)
^{15}N	139	-0.11 ± 0.53	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1324 atoms were assigned a chemical shift out of a possible 1491. 21 out of 23 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	638/640 (100%)	254/255 (100%)	260/260 (100%)	124/125 (99%)
Sidechain	639/749 (85%)	391/434 (90%)	245/285 (86%)	3/30 (10%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	47/102 (46%)	35/53 (66%)	0/33 (0%)	12/16 (75%)
Overall	1324/1491 (89%)	680/742 (92%)	505/578 (87%)	139/171 (81%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 1493 atoms were assigned a chemical shift out of a possible 1702. 21 out of 23 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	731/755 (97%)	291/301 (97%)	301/306 (98%)	139/148 (94%)
Sidechain	715/845 (85%)	440/489 (90%)	269/322 (84%)	6/34 (18%)
Aromatic	47/102 (46%)	35/53 (66%)	0/33 (0%)	12/16 (75%)
Overall	1493/1702 (88%)	766/843 (91%)	570/661 (86%)	157/198 (79%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	62	PRO	CD	34.00	55.31 – 45.41	-16.5
1	A	71	HIS	NE2	17.00	255.67 – 109.17	-11.3
1	A	17	ILE	CG1	8.00	36.54 – 18.94	-11.2
1	A	19	ASN	HB2	0.32	4.36 – 1.26	-8.0
1	A	35	ILE	CG2	27.30	24.63 – 10.43	6.9
1	A	106	LEU	CD1	14.50	32.77 – 16.57	-6.3
1	A	43	HIS	C	164.70	185.27 – 165.27	-5.3
1	A	35	ILE	CG1	18.90	36.54 – 18.94	-5.0

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

