



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:33 PM GMT

PDB ID : 1BEJ  
Title : INTERACTION BETWEEN PROXIMAL AND DISTALS REGIONS OF  
CYTOCHROME C PEROXIDASE  
Authors : Miller, M.A.; Kraut, J.  
Deposited on : 1998-05-16  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	291	



In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	S	0	0	0
			2315	1478	385	446	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ILE	THR	MUTATION	UNP P00431
A	152	GLY	ASP	MUTATION	UNP P00431
A	191	GLY	TRP	MUTATION	UNP P00431
A	272	ASP	ASN	MUTATION	UNP P00431

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	195	Total 195	O 195	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: CYTOCHROME C PEROXIDASE



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.96Å 74.11Å 45.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40	Depositor
% Data completeness (in resolution range)	97.0 (20.00-2.40)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
Refinement program	TNT	Depositor
R, $R_{free}$	0.164 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2553	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.03	17/2379 (0.7%)	1.66	57/3223 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	2

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	118	GLU	CD-OE1	5.95	1.32	1.25
1	A	167	GLU	CD-OE1	5.71	1.31	1.25
1	A	76	GLU	CD-OE1	5.69	1.31	1.25
1	A	291	GLU	CD-OE1	5.62	1.31	1.25
1	A	214	GLU	CD-OE1	5.58	1.31	1.25
1	A	135	GLU	CD-OE1	5.51	1.31	1.25
1	A	188	GLU	CD-OE1	5.42	1.31	1.25
1	A	201	GLU	CD-OE2	5.39	1.31	1.25
1	A	290	GLU	CD-OE1	5.36	1.31	1.25
1	A	209	GLU	CD-OE2	5.27	1.31	1.25
1	A	291	GLU	CD-OE2	-5.23	1.19	1.25
1	A	11	GLU	CD-OE1	5.19	1.31	1.25
1	A	98	GLU	CD-OE1	5.18	1.31	1.25
1	A	221	GLU	CD-OE1	5.08	1.31	1.25
1	A	290	GLU	CD-OE2	-5.07	1.20	1.25
1	A	167	GLU	CD-OE2	-5.05	1.20	1.25
1	A	32	GLU	CD-OE2	5.03	1.31	1.25

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	127	ARG	NE-CZ-NH1	9.86	125.23	120.30
1	A	165	ASP	CB-CG-OD2	-9.57	109.68	118.30
1	A	61	ASP	CB-CG-OD2	-8.94	110.26	118.30
1	A	14	ARG	NE-CZ-NH2	-8.57	116.02	120.30
1	A	146	ASP	CB-CG-OD1	-8.55	110.61	118.30
1	A	18	ASP	CB-CG-OD2	8.39	125.85	118.30
1	A	254	ASP	CB-CG-OD2	-8.22	110.90	118.30
1	A	150	ASP	CB-CG-OD1	8.10	125.59	118.30
1	A	58	ASP	CB-CG-OD2	-8.09	111.02	118.30
1	A	165	ASP	CB-CG-OD1	7.92	125.43	118.30
1	A	106	ASP	CB-CG-OD1	7.80	125.33	118.30
1	A	18	ASP	CB-CG-OD1	-7.78	111.30	118.30
1	A	150	ASP	CB-CG-OD2	-7.76	111.32	118.30
1	A	272	ASP	CB-CG-OD2	-7.64	111.43	118.30
1	A	38	ASN	CA-CB-CG	-7.55	96.79	113.40
1	A	106	ASP	CB-CG-OD2	-7.54	111.51	118.30
1	A	132	ASP	CB-CG-OD2	-7.50	111.55	118.30
1	A	272	ASP	CB-CG-OD1	7.45	125.01	118.30
1	A	61	ASP	CB-CG-OD1	7.16	124.75	118.30
1	A	256	ASP	CB-CG-OD2	-7.06	111.95	118.30
1	A	244	TYR	CB-CG-CD1	7.02	125.21	121.00
1	A	241	ASP	CB-CG-OD1	-7.01	111.99	118.30
1	A	136	ASP	CB-CG-OD1	-6.85	112.14	118.30
1	A	254	ASP	CB-CG-OD1	6.74	124.37	118.30
1	A	58	ASP	CB-CG-OD1	6.72	124.34	118.30
1	A	196	ASN	N-CA-CB	6.70	122.66	110.60
1	A	256	ASP	CB-CG-OD1	6.47	124.13	118.30
1	A	132	ASP	CB-CG-OD1	6.47	124.12	118.30
1	A	252	ALA	N-CA-CB	6.47	119.15	110.10
1	A	48	ARG	NE-CZ-NH1	6.46	123.53	120.30
1	A	235	ASP	CB-CG-OD2	-6.43	112.51	118.30
1	A	38	ASN	N-CA-CB	6.22	121.80	110.60
1	A	210	ASP	CB-CG-OD1	-6.20	112.72	118.30
1	A	127	ARG	CD-NE-CZ	6.18	132.25	123.60
1	A	67	TYR	CB-CG-CD2	-6.18	117.29	121.00
1	A	34	ASP	CB-CG-OD1	6.15	123.83	118.30
1	A	146	ASP	CB-CG-OD2	5.90	123.61	118.30
1	A	187	TYR	N-CA-CB	5.88	121.18	110.60
1	A	244	TYR	CB-CG-CD2	-5.84	117.49	121.00
1	A	185	SER	N-CA-CB	-5.80	101.81	110.50
1	A	33	ASP	CB-CG-OD1	-5.78	113.10	118.30
1	A	48	ARG	CD-NE-CZ	5.76	131.67	123.60

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	THR	CA-CB-CG2	-5.75	104.34	112.40
1	A	229	TYR	CB-CG-CD2	-5.75	117.55	121.00
1	A	160	ARG	NE-CZ-NH2	-5.74	117.43	120.30
1	A	136	ASP	CB-CG-OD2	5.71	123.44	118.30
1	A	148	ASP	CB-CG-OD1	5.53	123.27	118.30
1	A	210	ASP	CB-CG-OD2	5.40	123.16	118.30
1	A	229	TYR	CB-CG-CD1	5.37	124.22	121.00
1	A	160	ARG	NE-CZ-NH1	5.29	122.95	120.30
1	A	70	THR	CA-CB-CG2	-5.29	104.99	112.40
1	A	265	ALA	CB-CA-C	5.24	117.96	110.10
1	A	34	ASP	CB-CG-OD2	-5.23	113.59	118.30
1	A	44	PRO	N-CA-CB	5.19	109.53	103.30
1	A	79	ASP	CB-CG-OD1	5.19	122.97	118.30
1	A	162	ASN	CB-CA-C	-5.07	100.27	110.40
1	A	140	ASP	CB-CG-OD2	-5.03	113.77	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	38	ASN	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162	ASN	Sidechain
1	A	38	ASN	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2315	0	2167	36	0
2	A	43	0	30	5	0
3	A	195	0	0	4	0
All	All	2553	0	2197	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 9.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:GLU:HB3	1:A:99:PHE:CD1	2.25	0.71
1:A:35:GLU:HG2	1:A:39:TYR:CE1	2.26	0.70
1:A:149:LYS:HG2	1:A:153:TYR:CG	2.29	0.68
1:A:149:LYS:HG2	1:A:153:TYR:CD1	2.30	0.66
1:A:59:LYS:HE3	1:A:159:GLN:O	1.97	0.65
1:A:213:LEU:HD11	1:A:221:GLU:HB3	1.78	0.65
1:A:257:LYS:HE3	1:A:261:ASP:OD2	1.99	0.63
1:A:245:LEU:HD22	1:A:245:LEU:O	2.01	0.61
1:A:75:LYS:NZ	1:A:138:THR:O	2.32	0.57
1:A:206:LEU:HD13	1:A:231:MET:SD	2.47	0.55
1:A:245:LEU:O	1:A:249:LYS:HG3	2.09	0.53
1:A:165:ASP:HA	3:A:465:HOH:O	2.08	0.53
1:A:113:VAL:HG13	1:A:124:ILE:HB	1.91	0.52
1:A:178:GLY:HA3	2:A:296:HEM:O2D	2.09	0.52
1:A:194:ALA:HB1	1:A:197:VAL:HG13	1.90	0.52
1:A:84:GLY:N	1:A:86:GLN:OE1	2.42	0.52
2:A:296:HEM:HBB2	2:A:296:HEM:CMB	2.41	0.50
1:A:122:PRO:HA	3:A:645:HOH:O	2.12	0.49
1:A:53:ILE:HG22	1:A:71:TYR:HB2	1.95	0.49
2:A:296:HEM:HMC1	2:A:296:HEM:HBC2	1.95	0.49
1:A:52:HIS:HE1	1:A:81:SER:O	1.95	0.49
1:A:148:ASP:HA	1:A:233:PRO:HG2	1.95	0.48
2:A:296:HEM:HMC1	2:A:296:HEM:CBC	2.43	0.48
1:A:73:PHE:CE2	1:A:135:GLU:HA	2.49	0.48
1:A:119:MET:O	1:A:120:GLN:HB2	2.13	0.47
1:A:195:ASN:OD1	1:A:195:ASN:N	2.40	0.47
1:A:272:ASP:HA	3:A:604:HOH:O	2.15	0.47
1:A:163:MET:HA	1:A:167:GLU:OE1	2.14	0.46
1:A:98:GLU:HB3	1:A:99:PHE:CE1	2.51	0.46
1:A:147:ALA:O	1:A:234:THR:HG23	2.17	0.45
1:A:184:ASN:HB2	2:A:296:HEM:O1A	2.17	0.45
1:A:75:LYS:HD3	1:A:140:ASP:HA	1.98	0.44
1:A:241:ASP:HA	1:A:242:PRO:HD3	1.88	0.44
1:A:103:SER:HB2	1:A:132:ASP:OD1	2.18	0.43
1:A:63:THR:HA	3:A:706:HOH:O	2.19	0.43
1:A:72:ARG:NH2	1:A:73:PHE:CZ	2.87	0.43
1:A:35:GLU:HG2	1:A:39:TYR:HE1	1.78	0.42
1:A:35:GLU:CG	1:A:39:TYR:CE1	3.00	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:PHE:CD2	1:A:135:GLU:HA	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/291 (99%)	278 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/248 (97%)	226 (94%)	14 (6%)	25	39

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	TRP
1	A	74	LYS
1	A	75	LYS
1	A	86	GLN
1	A	166	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	197	VAL
1	A	206	LEU
1	A	212	LYS
1	A	225	SER
1	A	237	SER
1	A	245	LEU
1	A	253	ASN
1	A	278	LYS
1	A	281	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	HIS
1	A	24	ASN
1	A	87	ASN
1	A	292	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	296	1,3	30,50,50	2.60	6 (20%)	24,82,82	2.77	14 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	296	1,3	-	0/10/54/54	0/0/8/8

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	296	HEM	C3B-C4B	-9.39	1.43	1.51
2	A	296	HEM	C2D-C3D	-6.86	1.33	1.54
2	A	296	HEM	C3D-C4D	-4.79	1.45	1.51
2	A	296	HEM	C2C-C1C	-3.87	1.45	1.52
2	A	296	HEM	C2B-C1B	-2.02	1.45	1.51
2	A	296	HEM	C1C-NC	2.28	1.38	1.36

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	296	HEM	C3B-CAB-CBB	-4.27	117.90	124.46
2	A	296	HEM	CAA-C2A-C1A	-4.13	122.52	127.01
2	A	296	HEM	C3B-C4B-CHC	-3.83	117.78	123.16
2	A	296	HEM	C4B-CHC-C1C	-3.59	119.82	125.82
2	A	296	HEM	C3B-C4B-NB	-2.27	107.30	111.63
2	A	296	HEM	CHD-C1D-ND	2.18	129.76	124.52
2	A	296	HEM	CMC-C2C-C3C	2.42	122.58	116.53
2	A	296	HEM	CAA-CBA-CGA	2.63	117.57	112.75
2	A	296	HEM	C2C-C1C-CHC	2.79	127.92	123.68
2	A	296	HEM	CMD-C2D-C3D	2.97	127.48	114.35
2	A	296	HEM	CAD-C3D-C4D	3.62	125.25	112.47
2	A	296	HEM	C2D-C3D-C4D	3.73	107.83	101.50
2	A	296	HEM	CMB-C2B-C3B	4.47	127.68	116.53
2	A	296	HEM	CAD-C3D-C2D	4.76	126.92	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	296	HEM	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.