



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:05 PM GMT

PDB ID : 4BHN
Title : Crystal Structures of Ask1-inhibitor Complexes
Authors : Singh, O.; Shillings, A.; Craggs, P.; Wall, I.; Rowland, P.; Skarzynski, T.; Hobbs, C.I.; Hardwick, P.; Tanner, R.; Blunt, M.; Witty, D.R.; Smith, K.J.
Deposited on : 2013-04-04
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>4%</div> <div>71% 5% • 23%</div> </div>
1	B	334	<div> <div>10%</div> <div>70% 7% 23%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1941	-	-	-	X
3	GOL	B	1939	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4290 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	0	0
			2016	1301	328	378	9			
1	B	257	Total	C	N	O	S	0	1	0
			1992	1282	322	379	9			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	644	MET	-	EXPRESSION TAG	UNP Q99683
A	645	ASP	-	EXPRESSION TAG	UNP Q99683
A	646	TYR	-	EXPRESSION TAG	UNP Q99683
A	647	LYS	-	EXPRESSION TAG	UNP Q99683
A	648	ASP	-	EXPRESSION TAG	UNP Q99683
A	649	ASP	-	EXPRESSION TAG	UNP Q99683
A	650	ASP	-	EXPRESSION TAG	UNP Q99683
A	651	ASP	-	EXPRESSION TAG	UNP Q99683
A	652	LYS	-	EXPRESSION TAG	UNP Q99683
A	653	GLU	-	EXPRESSION TAG	UNP Q99683
A	654	ASN	-	EXPRESSION TAG	UNP Q99683
A	655	LEU	-	EXPRESSION TAG	UNP Q99683
A	656	TYR	-	EXPRESSION TAG	UNP Q99683
A	657	PHE	-	EXPRESSION TAG	UNP Q99683
A	658	GLN	-	EXPRESSION TAG	UNP Q99683
A	659	GLY	-	EXPRESSION TAG	UNP Q99683
A	838	GLU	THR	ENGINEERED MUTATION	UNP Q99683
B	644	MET	-	EXPRESSION TAG	UNP Q99683
B	645	ASP	-	EXPRESSION TAG	UNP Q99683
B	646	TYR	-	EXPRESSION TAG	UNP Q99683
B	647	LYS	-	EXPRESSION TAG	UNP Q99683
B	648	ASP	-	EXPRESSION TAG	UNP Q99683
B	649	ASP	-	EXPRESSION TAG	UNP Q99683
B	650	ASP	-	EXPRESSION TAG	UNP Q99683

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	651	ASP	-	EXPRESSION TAG	UNP Q99683
B	652	LYS	-	EXPRESSION TAG	UNP Q99683
B	653	GLU	-	EXPRESSION TAG	UNP Q99683
B	654	ASN	-	EXPRESSION TAG	UNP Q99683
B	655	LEU	-	EXPRESSION TAG	UNP Q99683
B	656	TYR	-	EXPRESSION TAG	UNP Q99683
B	657	PHE	-	EXPRESSION TAG	UNP Q99683
B	658	GLN	-	EXPRESSION TAG	UNP Q99683
B	659	GLY	-	EXPRESSION TAG	UNP Q99683
B	838	GLU	THR	ENGINEERED MUTATION	UNP Q99683

- Molecule 2 is 4-TERT-BUTYL-N-[6-(1H-PYRAZOL-4-YL)IMIDAZO[1,2-A]PYRIDIN-2-YL]BENZAMIDE (three-letter code: BH9) (formula: C₂₁H₂₁N₅O).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	130	Total	O	0	0
			130	130		
4	B	86	Total	O	0	0
			86	86		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5



- Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.49 Å 77.49 Å 423.83 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	31.20 – 2.30 31.20 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.6 (31.20-2.30) 95.1 (31.20-2.30)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.31 Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.196 , 0.235 0.206 , 0.253	Depositor DCC
R_{free} test set	1351 reflections (4.26%)	DCC
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 64.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 33093 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4290	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BH9, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/2061	0.70	0/2780
1	B	0.49	0/2040	0.68	0/2760
All	All	0.50	0/4101	0.69	0/5540

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2016	0	1983	11	0
1	B	1992	0	1925	6	0
2	A	27	0	21	0	0
2	B	27	0	21	0	0
3	A	6	0	8	1	0
3	B	6	0	8	1	0
4	A	130	0	0	2	0
4	B	86	0	0	0	0
All	All	4290	0	3966	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:815:SER:HB3	3:B:1939:GOL:H31	1.71	0.72
1:A:710:GLU:HG2	1:A:751:LYS:HG2	1.82	0.61
1:B:869:CYS:HB3	1:B:880:PRO:HG3	1.93	0.51
1:A:872:ILE:HG21	1:A:901:PRO:HB2	1.93	0.50
1:A:728:LEU:HD11	1:A:799:ILE:HG12	1.94	0.49
1:A:698:ARG:HD3	3:A:1941:GOL:C3	2.42	0.49
1:A:738:VAL:HG11	4:A:2001:HOH:O	2.13	0.48
1:A:844:GLN:HG2	4:A:2087:HOH:O	2.13	0.48
1:A:849:GLU:HG2	1:A:923:PRO:HD3	1.96	0.47
1:A:726:ILE:O	1:A:730:LYS:HB2	2.15	0.47
1:B:817:VAL:HG12	1:B:819:LYS:HE2	1.95	0.46
1:A:825:THR:HB	1:A:839:PHE:CE2	2.52	0.45
1:B:692:GLY:HA2	1:B:711:ILE:HD13	2.00	0.43
1:A:849:GLU:H	1:A:849:GLU:HG3	1.20	0.42
1:A:738:VAL:HG12	1:A:821:SER:HA	2.02	0.40
1:B:907:MET:O	1:B:912:LYS:HE2	2.21	0.40
1:B:890:ALA:O	1:B:894:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/334 (75%)	241 (96%)	8 (3%)	1 (0%)	39	48
1	B	252/334 (75%)	241 (96%)	10 (4%)	1 (0%)	39	48
All	All	502/668 (75%)	482 (96%)	18 (4%)	2 (0%)	39	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	822	ASP
1	B	822	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/289 (73%)	208 (98%)	4 (2%)	65	81
1	B	207/289 (72%)	195 (94%)	12 (6%)	25	33
All	All	419/578 (72%)	403 (96%)	16 (4%)	40	54

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	690	THR
1	A	844	GLN
1	A	849	GLU
1	A	883	GLU
1	B	670	LEU
1	B	676	GLU
1	B	679	GLU
1	B	709	LYS
1	B	719	SER
1	B	722	LEU
1	B	745	SER
1	B	752	ILE
1	B	819	LYS
1	B	843	LEU
1	B	905	GLU
1	B	920	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	756	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BH9	A	1	-	24,30,30	2.45	3 (12%)	29,44,44	1.22	4 (13%)
3	GOL	A	1941	-	5,5,5	0.36	0	5,5,5	0.54	0
2	BH9	B	1	-	24,30,30	2.43	3 (12%)	29,44,44	1.32	2 (6%)
3	GOL	B	1939	-	5,5,5	0.48	0	5,5,5	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BH9	A	1	-	-	0/16/18/18	0/4/4/4
3	GOL	A	1941	-	-	0/4/4/4	0/0/0/0
2	BH9	B	1	-	-	0/16/18/18	0/4/4/4
3	GOL	B	1939	-	-	0/4/4/4	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	BH9	C2-N1	-3.60	1.26	1.33
2	B	1	BH9	C2-N1	-3.49	1.26	1.33
2	B	1	BH9	C9-N14	3.48	1.36	1.33
2	A	1	BH9	C9-N14	3.59	1.36	1.33
2	A	1	BH9	C4-N5	10.25	1.54	1.33
2	B	1	BH9	C4-N5	10.34	1.54	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	BH9	C13-N15-C16	-3.38	119.28	128.09
2	B	1	BH9	C11-C10-C9	-2.89	116.22	119.72
2	A	1	BH9	C11-C10-C9	-2.46	116.74	119.72
2	A	1	BH9	C13-N15-C16	-2.44	121.71	128.09
2	A	1	BH9	C4-C14-C6	-2.17	125.08	127.84
2	A	1	BH9	C17-C16-N15	-2.15	111.75	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1941	GOL	1	0
3	B	1939	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/334 (76%)	0.04	15 (5%) 26 34	26, 42, 77, 111	0
1	B	257/334 (76%)	0.52	34 (13%) 4 7	33, 51, 80, 105	0
All	All	513/668 (76%)	0.28	49 (9%) 10 15	26, 46, 79, 111	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	721	PRO	6.6
1	B	839	PHE	6.6
1	B	851	ILE	6.1
1	B	840	THR	5.7
1	B	855	PRO	5.1
1	B	852	ASP	4.7
1	A	670	LEU	4.7
1	B	867	LEU	4.6
1	B	887	PRO	4.6
1	B	714	ARG	4.6
1	B	804	ILE	4.0
1	B	854	GLY	3.9
1	B	712	PRO	3.6
1	B	895	GLY	3.6
1	B	886	GLU	3.5
1	A	731	HIS	3.4
1	B	830	ALA	3.4
1	B	884	LEU	3.3
1	A	723	HIS	3.2
1	A	940	VAL	3.2
1	B	809	VAL	3.2
1	A	804	ILE	3.0
1	B	892	PHE	2.9
1	B	787	ILE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	853	LYS	2.8
1	B	857	GLY	2.8
1	B	868	GLY	2.7
1	B	871	ILE	2.7
1	B	723	HIS	2.7
1	A	867	LEU	2.7
1	B	866	SER	2.7
1	A	931	ASP	2.6
1	A	829	LEU	2.6
1	B	721	PRO	2.5
1	A	787	ILE	2.5
1	B	934	VAL	2.5
1	A	939	LYS	2.5
1	B	869	CYS	2.4
1	B	670	LEU	2.4
1	A	691	TYR	2.4
1	B	691	TYR	2.4
1	B	896	MET	2.4
1	A	882	TYR	2.3
1	B	870	THR	2.3
1	B	820	ILE	2.3
1	A	828	ARG	2.3
1	B	889	ALA	2.2
1	B	856	ARG	2.2
1	A	934	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	B	1939	6/6	0.40	0.72	42.71	165,166,166,167	0
3	GOL	A	1941	6/6	0.86	0.33	5.08	56,58,60,60	0
2	BH9	B	1	27/27	0.96	0.13	-0.32	30,34,39,42	0
2	BH9	A	1	27/27	0.96	0.12	-0.36	23,33,36,36	0

6.5 Other polymers ⓘ

There are no such residues in this entry.