Full wwPDB X-ray Structure Validation Report (i)

Jan 31, 2016 – 06:35 PM GMT

PDB ID	:	1BIZ
Title	:	HIV-1 INTEGRASE CORE DOMAIN
Authors	:	Goldgur, Y.; Dyda, F.; Hickman, A.B.; Jenkins, T.M.; Craigie, R.; Davies, D.R.
Deposited on	:	1998-06-21
Resolution	:	1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	166	72% 14% • • 11%
1	B	166	73% 12% • 14%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 INTEGRASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	147	1124	716	195	209	4	0	0	0
			Total	C	N	O	S			
1	B	143	1088	692	191	201	4	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	SER	GLU	CONFLICT	UNP Q76353
A	49	HIS	ALA	CONFLICT	UNP Q76353
A	56	SER	CYS	ENGINEERED	UNP Q76353
A	185	LYS	PHE	ENGINEERED	UNP Q76353
B	48	SER	GLU	CONFLICT	UNP Q76353
B	49	HIS	ALA	CONFLICT	UNP Q76353
B	56	SER	CYS	CONFLICT	UNP Q76353
B	185	LYS	PHE	ENGINEERED	UNP Q76353

- Molecule 2 is CACODYLATE ION (three-letter code: CAC) (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total As 1 1	0	0
2	A	1	Total As 1 1	0	0
2	B	1	Total As 1 1	0	0
2	B	1	Total As 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	95	Total O 95 95	0	0
3	B	117	Total O 117 117	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HIV-1 INTEGRASE

Chain A:



- Molecule 1: HIV-1 INTEGRASE

Chain B:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	50.84Å 71.34Å 91.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.95	Depositor
% Data completeness (in resolution range)	93.8 ((Not available)-1.95)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.212 , 0.253	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2428	wwPDB-VP
Average B, all atoms (Å^2)	32.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.36	0/1145	0.67	2/1548 (0.1%)
1	B	0.35	0/1108	0.56	0/1497
All	All	0.36	0/2253	0.62	2/3045 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	56	SER	O-C-N	5.68	131.79	122.70
1	A	56	SER	CA-C-N	-5.00	106.19	117.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1124	0	1126	23	0
1	B	1088	0	1085	13	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	95	0	0	2	0
3	B	117	0	0	4	0
All	All	2428	0	2211	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:68:LEU:HG	1:B:159:LYS:HE3	1.69	0.72
1:A:84:ILE:HD11	1:A:154:MET:HG2	1.75	0.66
1:A:79:VAL:HG22	1:A:151:ILE:HD11	1.77	0.66
1:A:187:ARG:HH11	1:A:187:ARG:HB2	1.63	0.62
1:A:112:THR:HB	1:A:138:GLU:HG2	1.81	0.61
1:A:112:THR:CB	1:A:138:GLU:HG2	2.31	0.59
1:A:77:VAL:HG22	1:A:84:ILE:CD1	2.35	0.55
1:B:65:CYS:SG	1:B:74:LEU:HD13	2.47	0.53
1:A:57:SER:HB3	1:A:60:ILE:HG13	1.90	0.53
1:A:77:VAL:HG22	1:A:84:ILE:HD12	1.89	0.53
1:A:112:THR:OG1	1:A:138:GLU:HG2	2.07	0.53
1:B:157:GLU:HG2	3:B:1522:HOH:O	2.08	0.53
1:A:89:ILE:HB	1:A:90:PRO:HD2	1.92	0.51
1:A:85:GLU:OE2	1:A:107:ARG:HD2	2.12	0.50
3:A:1059:HOH:O	1:B:87:GLU:HG3	2.12	0.50
1:B:181:PHE:O	1:B:185:LYS:HD3	2.12	0.49
1:A:60:ILE:HG23	1:A:138:GLU:HG3	1.95	0.48
1:B:185:LYS:NZ	3:B:1505:HOH:O	2.37	0.48
1:A:81:SER:HB2	1:A:199:ARG:HB3	1.96	0.46
1:B:117:ASN:HB3	1:B:120:ASN:ND2	2.30	0.46
1:A:173:LYS:HB3	1:B:99:TYR:CE1	2.53	0.44
1:A:76:ALA:HB2	1:A:100:PHE:HE2	1.83	0.44
1:B:169:ALA:HB1	1:B:174:THR:HB	2.00	0.43
1:B:173:LYS:HG3	3:B:1060:HOH:O	2.19	0.43
1:B:58:PRO:HB2	3:B:1086:HOH:O	2.19	0.43
1:A:173:LYS:HG3	3:A:1459:HOH:O	2.18	0.42
1:A:187:ARG:HB2	1:A:187:ARG:NH1	2.30	0.42
1:A:112:THR:HA	1:A:136:LYS:O	2.19	0.42
1:B:206:THR:O	1:B:210:THR:HG23	2.20	0.41
1:A:107:ARG:HG3	1:A:108:TRP:CD1	2.55	0.41
1:A:62:GLN:HG2	1:A:151:ILE:HG21	2.02	0.40
1:A:199:ARG:HA	1:A:199:ARG:HH11	1.87	0.40
1:B:63:LEU:HD13	1:B:76:ALA:HB2	2.04	0.40
1:A:200:ILE:O	1:A:204:ILE:HG12	2.21	0.40
1:A:102:LEU:HD12	1:A:102:LEU:HA	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/166 (86%)	139 (97%)	3 (2%)	1 (1%)	26	14
1	B	139/166 (84%)	133 (96%)	5 (4%)	1 (1%)	26	14
All	All	282/332 (85%)	272 (96%)	8 (3%)	2 (1%)	26	14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	SER
1	B	189	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/135 (87%)	112 (96%)	5 (4%)	35	20
1	B	111/135 (82%)	107 (96%)	4 (4%)	42	28
All	All	228/270 (84%)	219 (96%)	9 (4%)	39	24

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	SER
1	A	57	SER
1	A	123	SER
1	A	154	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	199	ARG
1	B	68	LEU
1	B	117	ASN
1	B	154	MET
1	B	168	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN
1	A	168	GLN
1	A	171	HIS
1	B	117	ASN
1	B	137	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates (i)

EDS was not executed - this section will therefore be empty.

6.4 Ligands (i)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers (i)

EDS was not executed - this section will therefore be empty.