



# Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 03:05 PM GMT

PDB ID : 4BIC  
Title : Crystal Structures of Ask1-inhibitor Complexes  
Authors : Singh, O.; Shillings, A.; Craggs, P.; Wall, I.; Rowland, P.; Skarzynski, T.; Hobbs, C.I.; Hardwick, P.; Tanner, R.; Blunt, M.; Witty, D.R.; Smith, K.J.  
Deposited on : 2013-04-10  
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2700 (2.64-2.60)
Clashscore	102246	3065 (2.64-2.60)
Ramachandran outliers	100387	3015 (2.64-2.60)
Sidechain outliers	100360	3015 (2.64-2.60)
RSRZ outliers	91569	2706 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 4162 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MITOGEN-ACTIVATED PROTEIN KINASE KINASE KINASE 5.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	260	Total	C 2021	N 330	O 380	S 9	0	0
1	B	259	Total	C 1979	N 1271	O 323	S 376	9	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	838	GLU	THR	ENGINEERED MUTATION	UNP Q99683
B	838	GLU	THR	ENGINEERED MUTATION	UNP Q99683

- Molecule 2 is N-(2-AMINOETHYL)-5-{1H-PYRROLO[2,3-B]PYRIDIN-3-YL}THIOPHEN E-2-SULFONAMIDE (three-letter code: IE4) (formula: C<sub>13</sub>H<sub>14</sub>N<sub>4</sub>O<sub>2</sub>S<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			21	13	4	2	2		

Mol	Chain	Residues	Total	C	N	O	S	ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			21	13	4	2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

Mol	Chain	Residues	Total	C	O	ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	74	Total	O	0	0
			74	74		


  

Mol	Chain	Residues	Total	O	ZeroOcc	AltConf
4	B	34	Total	O	0	0
			34	34		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE KINASE 5



GLU  
ASP  
THR  
SER

- Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE KINASE 5



S841  
SER  
THR  
GLU  
GLU  
GLU  
PHE  
T840  
LYS  
G841  
GLY  
ASP  
CYS  
T842  
GLU  
L843  
SER  
PRO  
LYS  
Y845  
LEU  
SER  
A847  
ALA  
LEU  
SER  
ALA  
GLY  
SER  
ASN  
GLU  
TRP  
LEU  
ARG  
SER  
ILE  
SER  
LEU  
PRO  
VAL  
PRO  
VAL  
LEU  
VAL  
GLU  
ASP  
THR  
SER

## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.95Å 77.95Å 418.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.84 – 2.62 38.98 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.84-2.62) 98.9 (38.98-2.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	6.50 (at 2.61Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
$R$ , $R_{free}$	0.207 , 0.265 0.266 , 0.307	Depositor DCC
$R_{free}$ test set	983 reflections (4.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	55.0	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 58.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Outliers	0 of 24247 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4162	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IE4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.53	0/2064	0.70	0/2786
1	B	0.48	0/2020	0.68	0/2732
All	All	0.50	0/4084	0.69	0/5518

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2021	0	1980	8	3
1	B	1979	0	1915	9	3
2	A	42	0	28	3	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	74	0	0	2	1
4	B	34	0	0	0	1
All	All	4162	0	3939	18	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2:IE4:H18	1:B:757:VAL:H	1.31	0.79
1:A:844:GLN:HG2	4:A:2058:HOH:O	1.84	0.78
1:A:757:VAL:H	2:A:1:IE4:H18	1.33	0.73
1:A:895:GLY:O	1:A:898:LYS:NZ	2.27	0.67
1:B:869:CYS:HB3	1:B:880:PRO:HG3	1.80	0.63
1:B:706:ILE:HD12	1:B:753:PHE:HD2	1.70	0.56
1:B:890:ALA:O	1:B:894:VAL:HG23	2.09	0.53
1:A:869:CYS:HB3	1:A:880:PRO:HG3	1.91	0.52
1:B:725:GLU:HG2	1:B:824:GLY:HA2	1.93	0.49
1:A:797:ASN:HB2	1:A:799:ILE:HD12	1.96	0.48
1:B:901:PRO:HD2	1:B:919:PHE:CZ	2.50	0.45
1:B:920:GLU:HB3	1:B:926:ARG:HG3	1.98	0.45
1:A:710:GLU:HG2	1:A:751:LYS:HG2	1.99	0.44
1:A:728:LEU:HD11	1:A:799:ILE:HG12	2.00	0.43
1:B:679:GLU:CD	1:B:679:GLU:H	2.22	0.43
2:A:2:IE4:H21	2:A:2:IE4:S13	2.59	0.42
1:A:676:GLU:HG2	4:A:2023:HOH:O	2.20	0.42
1:B:844:GLN:HB3	1:B:880:PRO:HG2	2.03	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:852:ASP:O	1:B:893:LYS:NZ[8_665]	2.08	0.12
1:A:747:ASN:O	4:B:2004:HOH:O[1_655]	2.13	0.07
1:A:680:ASN:OD1	1:B:747:ASN:ND2[1_655]	2.17	0.03
1:B:680:ASN:O	4:A:2003:HOH:O[1_455]	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	254/318 (80%)	240 (94%)	12 (5%)	2 (1%)	24 44
1	B	253/318 (80%)	242 (96%)	10 (4%)	1 (0%)	39 63
All	All	507/636 (80%)	482 (95%)	22 (4%)	3 (1%)	30 54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	822	ASP
1	A	747	ASN
1	B	822	ASP

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/274 (77%)	204 (97%)	6 (3%)	50 76
1	B	203/274 (74%)	194 (96%)	9 (4%)	35 62
All	All	413/548 (75%)	398 (96%)	15 (4%)	42 70

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	674	ASP
1	A	679	GLU
1	A	825	THR
1	A	844	GLN
1	A	878	LYS
1	A	883	GLU
1	B	670	LEU
1	B	671	LEU
1	B	729	HIS
1	B	733	LYS
1	B	745	SER
1	B	822	ASP
1	B	825	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	851	ILE
1	B	916	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	930	ASN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	IE4	A	1	-	19,23,23	2.54	3 (15%)	17,33,33	2.61	7 (41%)
3	GOL	A	1941	-	5,5,5	0.98	1 (20%)	5,5,5	2.18	2 (40%)
2	IE4	A	2	-	19,23,23	2.44	3 (15%)	17,33,33	2.31	7 (41%)
3	GOL	B	1942	-	5,5,5	0.81	0	5,5,5	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IE4	A	1	-	-	0/5/15/15	0/3/3/3
3	GOL	A	1941	-	-	0/4/4/4	0/0/0/0
2	IE4	A	2	-	-	0/5/15/15	0/3/3/3
3	GOL	B	1942	-	-	0/4/4/4	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	IE4	C17-N18	-2.54	1.33	1.37
3	A	1941	GOL	C1-C2	-2.08	1.44	1.52
2	A	2	IE4	C17-N18	-2.02	1.33	1.37
2	A	2	IE4	O08-S06	2.57	1.46	1.43
2	A	1	IE4	O08-S06	2.92	1.46	1.43
2	A	2	IE4	S06-N05	9.36	1.73	1.61
2	A	1	IE4	S06-N05	9.73	1.74	1.61

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	IE4	O07-S06-C09	-5.97	97.79	107.63
2	A	2	IE4	O07-S06-C09	-5.59	98.42	107.63
2	A	1	IE4	C20-C21-C22	-4.30	114.79	120.88
2	A	1	IE4	C19-N18-C17	-4.03	111.24	116.44
2	A	1	IE4	C09-S06-N05	-3.39	103.38	107.92
2	A	2	IE4	C19-N18-C17	-3.14	112.38	116.44
2	A	2	IE4	C20-C21-C22	-2.89	116.80	120.88
3	A	1941	GOL	C3-C2-C1	-2.88	99.84	111.12
2	A	2	IE4	C14-C15-N16	-2.09	104.87	108.91
2	A	2	IE4	C09-S06-N05	-2.07	105.16	107.92
2	A	1	IE4	C20-C19-N18	2.34	127.62	123.94
3	A	1941	GOL	O2-C2-C3	2.76	121.30	108.65
2	A	1	IE4	O08-S06-O07	3.13	123.70	119.54
2	A	2	IE4	O07-S06-N05	3.25	112.25	107.03
2	A	1	IE4	O07-S06-N05	3.91	113.31	107.03
2	A	2	IE4	O08-S06-O07	3.93	124.76	119.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	IE4	1	0
2	A	2	IE4	2	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	260/318 (81%)	0.55	17 (6%) 22 16	32, 52, 95, 130	0
1	B	259/318 (81%)	1.15	44 (16%) 2 1	40, 69, 109, 132	0
All	All	519/636 (81%)	0.85	61 (11%) 6 4	32, 59, 104, 132	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	940	VAL	7.7
1	B	855	PRO	7.6
1	B	858	TYR	6.8
1	B	854	GLY	6.0
1	B	851	ILE	5.9
1	B	896	MET	5.5
1	A	721	PRO	5.1
1	B	852	ASP	4.9
1	B	750	ILE	4.7
1	B	714	ARG	4.3
1	B	748	GLY	4.2
1	B	712	PRO	4.1
1	A	750	ILE	4.0
1	B	940	VAL	3.9
1	B	889	ALA	3.8
1	B	749	PHE	3.8
1	A	670	LEU	3.8
1	A	722	LEU	3.8
1	B	884	LEU	3.5
1	A	720	GLN	3.5
1	A	719	SER	3.4
1	B	937	PHE	3.4
1	B	802	ARG	3.4
1	B	909	ALA	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	881	PHE	3.4
1	B	856	ARG	3.3
1	B	840	THR	3.3
1	B	843	LEU	3.3
1	B	744	PHE	3.2
1	B	895	GLY	3.2
1	B	916	LEU	3.1
1	B	713	GLU	3.1
1	B	921	PRO	2.9
1	B	857	GLY	2.9
1	B	941	SER	2.8
1	A	731	HIS	2.8
1	B	764	ALA	2.8
1	A	897	PHE	2.8
1	B	848	PRO	2.8
1	B	747	ASN	2.7
1	A	939	LYS	2.6
1	B	925	LYS	2.6
1	A	712	PRO	2.6
1	A	671	LEU	2.6
1	B	841	GLY	2.6
1	A	848	PRO	2.6
1	B	671	LEU	2.5
1	B	719	SER	2.5
1	A	723	HIS	2.5
1	B	824	GLY	2.4
1	A	829	LEU	2.4
1	B	927	ALA	2.4
1	B	846	MET	2.3
1	B	862	ALA	2.3
1	B	680	ASN	2.2
1	B	934	VAL	2.2
1	A	679	GLU	2.1
1	A	724	GLU	2.1
1	B	823	PHE	2.1
1	B	745	SER	2.1
1	B	892	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	1942	6/6	0.86	0.28	1.34	54,57,58,58	0
2	IE4	A	2	21/21	0.78	0.25	1.04	41,54,79,79	0
3	GOL	A	1941	6/6	0.90	0.24	0.85	53,53,54,55	0
2	IE4	A	1	21/21	0.93	0.17	-0.58	31,40,68,71	0

### 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.