



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 23, 2017 – 11:40 PM EST

PDB ID : 5BOZ  
Title : Ricin A chain bound to camelid nanobody (VHH9)(E1)  
Authors : Rudolph, M.J.; Mantis, N.  
Deposited on : 2015-05-27  
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	<div> <div>56%</div> <div>39%</div> <div>..</div> </div>
1	B	261	<div> <div>49%</div> <div>45%</div> <div>5%</div> </div>
1	C	261	<div> <div>53%</div> <div>42%</div> <div>..</div> </div>
1	D	261	<div> <div>2%</div> <div>52%</div> <div>42%</div> <div>..</div> </div>
1	E	261	<div> <div>50%</div> <div>44%</div> <div>5%</div> </div>
1	F	261	<div> <div>51%</div> <div>44%</div> <div>.</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	127	
2	H	127	
2	I	127	
2	J	127	
2	K	127	
2	L	127	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	301	-	-	X	-
3	SO4	B	301	-	-	X	-
3	SO4	B	304	-	-	X	-
3	SO4	C	301	-	-	X	-
3	SO4	E	301	-	-	-	X
3	SO4	F	301	-	-	X	-
3	SO4	G	201	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 18136 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ricin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			2046	1296	362	383	5			
1	B	260	Total	C	N	O	S	0	0	0
			2052	1299	363	385	5			
1	C	259	Total	C	N	O	S	0	0	0
			2046	1296	362	383	5			
1	D	259	Total	C	N	O	S	0	0	0
			2046	1296	362	383	5			
1	E	259	Total	C	N	O	S	0	2	0
			2068	1308	370	385	5			
1	F	260	Total	C	N	O	S	0	0	0
			2052	1299	363	385	5			

- Molecule 2 is a protein called VHH single chain antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	123	Total	C	N	O	S	0	1	0
			961	601	173	182	5			
2	H	125	Total	C	N	O	S	0	1	0
			973	607	176	185	5			
2	I	118	Total	C	N	O	S	0	0	0
			911	572	160	174	5			
2	J	119	Total	C	N	O	S	0	0	0
			918	576	161	176	5			
2	K	125	Total	C	N	O	S	0	1	0
			973	607	176	185	5			
2	L	123	Total	C	N	O	S	0	1	0
			961	601	173	182	5			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	I	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	J	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Cl	0	0
			1	1		
4	K	1	Total	Cl	0	0
			1	1		
4	E	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	3	Total	O	0	0
			3	3		
5	B	1	Total	O	0	0
			1	1		
5	H	1	Total	O	0	0
			1	1		
5	C	4	Total	O	0	0
			4	4		
5	I	1	Total	O	0	0
			1	1		
5	D	1	Total	O	0	0
			1	1		
5	E	5	Total	O	0	0
			5	5		
5	K	6	Total	O	0	0
			6	6		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	F	2	Total	O	0	0
			2	2		
5	L	2	Total	O	0	0
			2	2		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Ricin




#### • Molecule 1: Ricin




#### • Molecule 1: Ricin









• Molecule 1: Ricin







- Molecule 2: VHH single chain antibody




- Molecule 2: VHH single chain antibody




- Molecule 2: VHH single chain antibody




- Molecule 2: VHH single chain antibody



- Molecule 2: VHH single chain antibody



• Molecule 2: VHH single chain antibody



## 4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	239.61Å 242.88Å 355.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.68 – 3.10 49.68 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.68-3.10) 99.8 (49.68-3.10)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 3.12Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.215 , 0.277 0.218 , 0.285	Depositor DCC
$R_{free}$ test set	4664 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	64.8	Xtriage
Anisotropy	0.664	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 98.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.000 for -k,-h,-l	Xtriage
Reported twinning fraction	0.017 for -k,-h,-l	Depositor
Outliers	12 of 93146 reflections (0.013%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	18136	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 64.97 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.6203e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.67	0/2091	0.83	1/2847 (0.0%)
1	B	0.74	1/2097 (0.0%)	0.87	2/2855 (0.1%)
1	C	0.70	2/2091 (0.1%)	0.83	1/2847 (0.0%)
1	D	0.64	0/2091	0.81	1/2847 (0.0%)
1	E	0.91	2/2113 (0.1%)	0.99	3/2875 (0.1%)
1	F	0.80	0/2097	0.89	2/2855 (0.1%)
2	G	0.68	0/977	0.87	0/1317
2	H	0.74	0/990	0.93	0/1336
2	I	0.61	0/927	0.83	0/1250
2	J	0.61	0/934	0.80	0/1260
2	K	1.16	1/990 (0.1%)	1.26	6/1336 (0.4%)
2	L	0.90	0/977	1.09	4/1317 (0.3%)
All	All	0.77	6/18375 (0.0%)	0.91	20/24942 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
1	D	0	2
1	E	0	3
1	F	0	3
2	H	0	1
2	K	0	1
2	L	0	2
All	All	0	18

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	64	ASN	CG-ND2	-8.30	1.12	1.32
1	E	64	ASN	CG-OD1	-7.53	1.07	1.24
1	C	64	ASN	CG-ND2	-7.48	1.14	1.32
1	C	64	ASN	CG-OD1	-7.10	1.08	1.24
1	B	177	GLU	CG-CD	5.76	1.60	1.51

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	26	GLY	N-CA-C	-10.52	86.79	113.10
2	L	45	ARG	NE-CZ-NH2	-9.31	115.65	120.30
2	K	109	ARG	NE-CZ-NH1	8.04	124.32	120.30
2	K	45	ARG	NE-CZ-NH2	-7.78	116.41	120.30
1	E	107	LEU	CA-CB-CG	-6.86	99.53	115.30

There are no chirality outliers.

5 of 18 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	36	ALA	Peptide
1	A	5	GLN	Peptide
1	B	36	ALA	Peptide
1	B	5	GLN	Peptide
2	H	25	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2046	0	2014	98	1
1	B	2052	0	2019	125	0
1	C	2046	0	2014	108	0
1	D	2046	0	2014	102	1
1	E	2068	0	2038	143	0
1	F	2052	0	2019	111	1
2	G	961	0	930	73	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	973	0	940	81	1
2	I	911	0	876	69	0
2	J	918	0	883	71	1
2	K	973	0	940	66	0
2	L	961	0	930	53	0
3	A	10	0	0	3	0
3	B	25	0	0	7	0
3	C	15	0	0	3	0
3	D	5	0	0	0	0
3	E	15	0	0	1	0
3	F	10	0	0	4	0
3	G	5	0	0	0	0
3	I	5	0	0	0	0
3	J	5	0	0	0	0
3	L	5	0	0	0	0
4	D	1	0	0	1	0
4	E	1	0	0	1	0
4	K	1	0	0	0	0
5	B	1	0	0	0	0
5	C	4	0	0	1	0
5	D	1	0	0	0	0
5	E	5	0	0	1	0
5	F	2	0	0	1	0
5	G	3	0	0	0	0
5	H	1	0	0	0	0
5	I	1	0	0	0	0
5	K	6	0	0	0	0
5	L	2	0	0	0	0
All	All	18136	0	17617	1058	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 1058 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:189:ARG:HH21	1:E:193[A]:ARG:NH2	1.20	1.37
1:E:189:ARG:NH2	1:E:193[A]:ARG:HH22	1.45	1.15
1:E:189:ARG:NH2	1:E:193[A]:ARG:NH2	2.00	1.07
2:H:72[A]:ARG:HH11	2:H:72[A]:ARG:HG2	1.28	0.99
2:G:88:PRO:HA	2:G:124:VAL:CG2	1.93	0.99

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:24:ALA:O	2:G:27:ARG:NH2[8_544]	1.85	0.35
2:H:3:GLN:NE2	2:H:3:GLN:NE2[8_544]	1.96	0.24
2:J:85:ASN:ND2	1:F:239:LYS:NZ[15_545]	2.17	0.03
1:A:244:ASP:OD1	1:D:244:ASP:OD1[13_455]	2.18	0.02

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/261 (98%)	244 (95%)	13 (5%)	0	100	100
1	B	258/261 (99%)	244 (95%)	14 (5%)	0	100	100
1	C	257/261 (98%)	243 (95%)	14 (5%)	0	100	100
1	D	257/261 (98%)	244 (95%)	13 (5%)	0	100	100
1	E	259/261 (99%)	248 (96%)	11 (4%)	0	100	100
1	F	258/261 (99%)	244 (95%)	14 (5%)	0	100	100
2	G	118/127 (93%)	105 (89%)	13 (11%)	0	100	100
2	H	122/127 (96%)	108 (88%)	14 (12%)	0	100	100
2	I	112/127 (88%)	100 (89%)	12 (11%)	0	100	100
2	J	113/127 (89%)	99 (88%)	14 (12%)	0	100	100
2	K	122/127 (96%)	112 (92%)	10 (8%)	0	100	100
2	L	118/127 (93%)	106 (90%)	12 (10%)	0	100	100
All	All	2251/2328 (97%)	2097 (93%)	154 (7%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/220 (99%)	200 (92%)	18 (8%)	14	46
1	B	219/220 (100%)	200 (91%)	19 (9%)	13	44
1	C	218/220 (99%)	201 (92%)	17 (8%)	16	49
1	D	218/220 (99%)	199 (91%)	19 (9%)	13	44
1	E	220/220 (100%)	204 (93%)	16 (7%)	17	52
1	F	219/220 (100%)	202 (92%)	17 (8%)	16	49
2	G	100/103 (97%)	85 (85%)	15 (15%)	3	15
2	H	101/103 (98%)	86 (85%)	15 (15%)	4	15
2	I	95/103 (92%)	81 (85%)	14 (15%)	4	16
2	J	96/103 (93%)	82 (85%)	14 (15%)	4	16
2	K	101/103 (98%)	82 (81%)	19 (19%)	2	8
2	L	100/103 (97%)	84 (84%)	16 (16%)	3	13
All	All	1905/1938 (98%)	1706 (90%)	199 (10%)	9	32

5 of 199 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	53	TRP
1	D	116	THR
2	L	28	THR
2	I	67	ARG
1	D	20	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	223	GLN
1	F	223	GLN
1	E	223	GLN
1	B	223	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	160	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 23 ligands modelled in this entry, 3 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	301	-	4,4,4	0.29	0	6,6,6	0.23	0
3	SO4	A	302	-	4,4,4	0.22	0	6,6,6	0.29	0
3	SO4	B	301	-	4,4,4	0.47	0	6,6,6	1.03	0
3	SO4	B	302	-	4,4,4	0.20	0	6,6,6	0.29	0
3	SO4	B	303	-	4,4,4	0.18	0	6,6,6	0.25	0
3	SO4	B	304	-	4,4,4	0.14	0	6,6,6	0.14	0
3	SO4	B	305	-	4,4,4	0.23	0	6,6,6	0.54	0
3	SO4	C	301	-	4,4,4	0.32	0	6,6,6	1.04	1 (16%)
3	SO4	C	302	-	4,4,4	0.12	0	6,6,6	0.38	0
3	SO4	C	303	-	4,4,4	0.19	0	6,6,6	0.42	0
3	SO4	D	301	-	4,4,4	0.12	0	6,6,6	0.43	0
3	SO4	E	301	-	4,4,4	0.36	0	6,6,6	0.48	0
3	SO4	E	302	-	4,4,4	0.29	0	6,6,6	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	E	303	-	4,4,4	0.47	0	6,6,6	0.77	0
3	SO4	F	301	-	4,4,4	0.60	0	6,6,6	0.31	0
3	SO4	F	302	-	4,4,4	0.14	0	6,6,6	0.38	0
3	SO4	G	201	-	4,4,4	0.24	0	6,6,6	0.50	0
3	SO4	I	201	-	4,4,4	0.08	0	6,6,6	0.38	0
3	SO4	J	201	-	4,4,4	0.12	0	6,6,6	0.24	0
3	SO4	L	201	-	4,4,4	0.20	0	6,6,6	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	301	-	-	0/0/0/0	0/0/0/0
3	SO4	A	302	-	-	0/0/0/0	0/0/0/0
3	SO4	B	301	-	-	0/0/0/0	0/0/0/0
3	SO4	B	302	-	-	0/0/0/0	0/0/0/0
3	SO4	B	303	-	-	0/0/0/0	0/0/0/0
3	SO4	B	304	-	-	0/0/0/0	0/0/0/0
3	SO4	B	305	-	-	0/0/0/0	0/0/0/0
3	SO4	C	301	-	-	0/0/0/0	0/0/0/0
3	SO4	C	302	-	-	0/0/0/0	0/0/0/0
3	SO4	C	303	-	-	0/0/0/0	0/0/0/0
3	SO4	D	301	-	-	0/0/0/0	0/0/0/0
3	SO4	E	301	-	-	0/0/0/0	0/0/0/0
3	SO4	E	302	-	-	0/0/0/0	0/0/0/0
3	SO4	E	303	-	-	0/0/0/0	0/0/0/0
3	SO4	F	301	-	-	0/0/0/0	0/0/0/0
3	SO4	F	302	-	-	0/0/0/0	0/0/0/0
3	SO4	G	201	-	-	0/0/0/0	0/0/0/0
3	SO4	I	201	-	-	0/0/0/0	0/0/0/0
3	SO4	J	201	-	-	0/0/0/0	0/0/0/0
3	SO4	L	201	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	301	SO4	O4-S-O3	2.10	117.54	109.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	SO4	3	0
3	B	301	SO4	2	0
3	B	304	SO4	4	0
3	B	305	SO4	1	0
3	C	301	SO4	2	0
3	C	303	SO4	1	0
3	E	303	SO4	1	0
3	F	301	SO4	3	0
3	F	302	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/261 (99%)	-0.15	2 (0%) 87 75	62, 93, 122, 145	0
1	B	260/261 (99%)	-0.21	2 (0%) 87 75	48, 80, 111, 147	0
1	C	259/261 (99%)	-0.17	0 100 100	60, 90, 119, 140	0
1	D	259/261 (99%)	-0.02	5 (1%) 70 48	74, 101, 128, 154	0
1	E	259/261 (99%)	-0.23	0 100 100	33, 66, 102, 134	0
1	F	260/261 (99%)	-0.28	0 100 100	45, 77, 116, 130	0
2	G	123/127 (96%)	0.38	4 (3%) 50 26	78, 108, 137, 179	0
2	H	125/127 (98%)	0.34	3 (2%) 62 39	60, 93, 120, 171	0
2	I	118/127 (92%)	0.34	9 (7%) 17 6	77, 104, 135, 156	0
2	J	119/127 (93%)	0.76	11 (9%) 11 4	96, 121, 144, 171	0
2	K	125/127 (98%)	-0.17	0 100 100	26, 58, 98, 111	0
2	L	123/127 (96%)	-0.02	1 (0%) 87 75	52, 84, 120, 139	0
All	All	2289/2328 (98%)	-0.03	37 (1%) 74 55	26, 89, 128, 179	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	85	ASN	3.8
2	I	67	ARG	3.3
2	G	65	LYS	3.1
2	I	84	ASN	3.1
2	G	28	THR	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	E	301	5/5	0.95	0.27	5.24	16,52,139,160	0
3	SO4	G	201	5/5	0.79	0.29	3.91	57,87,160,182	0
3	SO4	L	201	5/5	0.95	0.21	1.49	87,97,114,115	0
3	SO4	I	201	5/5	0.93	0.21	1.35	48,90,150,162	0
3	SO4	E	302	5/5	0.94	0.25	0.93	28,70,183,217	0
3	SO4	B	303	5/5	0.90	0.22	0.68	30,74,172,189	0
3	SO4	J	201	5/5	0.87	0.22	0.45	60,119,163,174	0
3	SO4	B	302	5/5	0.84	0.23	0.32	54,130,184,193	0
3	SO4	A	301	5/5	0.97	0.16	-0.84	40,47,104,133	0
3	SO4	B	301	5/5	0.98	0.10	-2.36	40,44,58,115	0
3	SO4	C	301	5/5	0.96	0.13	-2.86	12,42,80,124	0
3	SO4	B	305	5/5	0.90	0.14	-	84,119,120,156	0
4	CL	E	304	1/1	0.98	0.33	-	30,30,30,30	0
3	SO4	C	302	5/5	0.93	0.15	-	98,104,154,156	0
4	CL	D	302	1/1	0.96	0.16	-	46,46,46,46	0
3	SO4	E	303	5/5	0.94	0.12	-	38,66,138,203	0
3	SO4	A	302	5/5	0.90	0.20	-	58,95,185,235	0
3	SO4	B	304	5/5	0.80	0.32	-	149,162,186,206	0
3	SO4	F	301	5/5	0.97	0.11	-	11,25,74,146	0
3	SO4	D	301	5/5	0.76	0.23	-	89,122,180,197	0
3	SO4	C	303	5/5	0.74	0.34	-	109,132,166,186	0
4	CL	K	201	1/1	0.93	0.20	-	42,42,42,42	0
3	SO4	F	302	5/5	0.95	0.14	-	54,97,156,163	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.