



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:12 PM GMT

PDB ID : 4BQ3  
Title : Structural analysis of an exo-beta-agarase  
Authors : Pluvinaige, B.; Hehemann, J.H.; Boraston, A.B.  
Deposited on : 2013-05-29  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	750	<div> <div>88%</div> <div>9% ..</div> </div>
1	B	750	<div> <div>89%</div> <div>10% ..</div> </div>
1	C	750	<div> <div>85%</div> <div>13% ..</div> </div>
1	D	750	<div> <div>90%</div> <div>9% .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1795	-	-	-	X
3	GOL	A	1797	-	-	X	X
3	GOL	A	1798	-	-	X	-
3	GOL	B	1797	-	-	-	X
3	GOL	B	1799	-	-	X	-
3	GOL	B	1800	-	-	-	X
3	GOL	B	1801	-	-	X	X
3	GOL	B	1802	-	-	-	X
3	GOL	C	1794	-	-	-	X
3	GOL	C	1796	-	-	-	X
3	GOL	C	1797	-	-	X	X
3	GOL	D	1795	-	-	-	X
3	GOL	D	1796	-	-	-	X
3	GOL	D	1798	-	-	-	X
3	GOL	D	1799	-	-	X	X
3	GOL	D	1800	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 26251 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called B-AGARASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	740	Total	C	N	O	S	0	5	0
			5920	3778	998	1120	24			
1	B	741	Total	C	N	O	S	0	6	0
			5940	3792	995	1128	25			
1	C	740	Total	C	N	O	S	0	4	0
			5899	3769	993	1114	23			
1	D	741	Total	C	N	O	S	0	3	0
			5886	3758	989	1116	23			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	GLY	-	EXPRESSION TAG	UNP Q21HC5
A	45	SER	-	EXPRESSION TAG	UNP Q21HC5
A	46	HIS	-	EXPRESSION TAG	UNP Q21HC5
B	44	GLY	-	EXPRESSION TAG	UNP Q21HC5
B	45	SER	-	EXPRESSION TAG	UNP Q21HC5
B	46	HIS	-	EXPRESSION TAG	UNP Q21HC5
C	44	GLY	-	EXPRESSION TAG	UNP Q21HC5
C	45	SER	-	EXPRESSION TAG	UNP Q21HC5
C	46	HIS	-	EXPRESSION TAG	UNP Q21HC5
D	44	GLY	-	EXPRESSION TAG	UNP Q21HC5
D	45	SER	-	EXPRESSION TAG	UNP Q21HC5
D	46	HIS	-	EXPRESSION TAG	UNP Q21HC5

- Molecule 2 is NEOAGAROBIOSE (three-letter code: 47N) (formula: C<sub>12</sub>H<sub>20</sub>O<sub>10</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			22	12	10		
2	B	1	Total	C	O	0	0
			22	12	10		
2	C	1	Total	C	O	0	0
			22	12	10		
2	D	1	Total	C	O	0	0
			22	12	10		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 6	C 3	O 3	0	0
3	A	1	Total 6	C 3	O 3	0	0
3	A	1	Total 6	C 3	O 3	0	0
3	A	1	Total 6	C 3	O 3	0	0
3	A	1	Total 6	C 3	O 3	0	0
3	A	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	C	1	Total 6	C 3	O 3	0	0
3	D	1	Total 6	C 3	O 3	0	0
3	D	1	Total 6	C 3	O 3	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	645	Total	O	0	0
			645	645		
5	B	667	Total	O	0	0
			667	667		
5	C	512	Total	O	0	0
			512	512		
5	D	528	Total	O	0	0
			528	528		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: B-AGARASE




#### • Molecule 1: B-AGARASE





#### • Molecule 1: B-AGARASE







• Molecule 1: B-AGARASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	170.59Å 170.59Å 116.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.98 – 2.10 38.98 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.98-2.10) 99.8 (38.98-2.10)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.02 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.159 , 0.209 0.160 , 0.209	Depositor DCC
$R_{free}$ test set	9757 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.0	EDS
Estimated twinning fraction	0.021 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 194282 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	26251	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, 47N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.75	0/6092	0.84	11/8275 (0.1%)
1	B	0.76	0/6112	0.83	11/8302 (0.1%)
1	C	0.70	0/6071	0.77	6/8252 (0.1%)
1	D	0.68	0/6058	0.75	2/8233 (0.0%)
All	All	0.73	0/24333	0.80	30/33062 (0.1%)

There are no bond length outliers.

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	471	ARG	NE-CZ-NH2	-11.54	114.53	120.30
1	B	471	ARG	NE-CZ-NH2	-11.01	114.80	120.30
1	A	471	ARG	NE-CZ-NH1	10.31	125.45	120.30
1	B	471	ARG	NE-CZ-NH1	10.07	125.34	120.30
1	A	611	ARG	NE-CZ-NH2	-8.04	116.28	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5920	0	5630	61	0
1	B	5940	0	5653	51	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	5899	0	5606	78	0
1	D	5886	0	5586	50	0
2	A	22	0	20	0	0
2	B	22	0	20	0	0
2	C	22	0	20	0	0
2	D	22	0	20	0	0
3	A	36	0	48	11	0
3	B	54	0	71	12	0
3	C	30	0	39	13	0
3	D	42	0	56	10	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	645	0	0	14	0
5	B	667	0	0	14	0
5	C	512	0	0	10	0
5	D	528	0	0	14	0
All	All	26251	0	22769	253	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 253 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1798:GOL:H11	5:A:2072:HOH:O	1.35	1.25
1:B:576:LYS:HE3	5:B:2528:HOH:O	1.39	1.20
1:B:614:TYR:O	1:B:617:LEU:O	1.62	1.13
1:C:630:HIS:CD2	3:C:1797:GOL:H32	1.87	1.10
3:D:1799:GOL:H2	5:D:2460:HOH:O	1.56	1.05

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	741/750 (99%)	718 (97%)	23 (3%)	0	100	100
1	B	743/750 (99%)	725 (98%)	17 (2%)	1 (0%)	56	58
1	C	740/750 (99%)	718 (97%)	22 (3%)	0	100	100
1	D	740/750 (99%)	720 (97%)	20 (3%)	0	100	100
All	All	2964/3000 (99%)	2881 (97%)	82 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	618	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	623/630 (99%)	616 (99%)	7 (1%)	80	85
1	B	628/630 (100%)	625 (100%)	3 (0%)	92	95
1	C	619/630 (98%)	612 (99%)	7 (1%)	80	85
1	D	618/630 (98%)	612 (99%)	6 (1%)	82	87
All	All	2488/2520 (99%)	2465 (99%)	23 (1%)	84	89

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	54	ASP
1	C	80	SER
1	D	745	MET
1	C	59	ASN
1	C	129	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	716	GLN
1	C	336	ASN
1	D	697	HIS
1	C	194	GLN
1	C	409	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 35 ligands modelled in this entry, 4 are monoatomic - leaving 31 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	47N	A	1792	-	24,24,24	1.15	2 (8%)	35,36,36	1.45	7 (20%)
3	GOL	A	1793	-	5,5,5	0.64	0	5,5,5	0.35	0
3	GOL	A	1794	-	5,5,5	0.60	0	5,5,5	1.76	1 (20%)
3	GOL	A	1795	-	5,5,5	0.52	0	5,5,5	0.86	0
3	GOL	A	1796	-	5,5,5	0.84	0	5,5,5	1.28	0
3	GOL	A	1797	-	5,5,5	0.99	0	5,5,5	1.44	1 (20%)
3	GOL	A	1798	-	5,5,5	0.81	0	5,5,5	1.89	2 (40%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	47N	B	1793	-	24,24,24	1.13	3 (12%)	35,36,36	1.61	8 (22%)
3	GOL	B	1794	-	5,5,5	0.63	0	5,5,5	0.27	0
3	GOL	B	1795	-	5,5,5	0.40	0	5,5,5	1.81	1 (20%)
3	GOL	B	1796	-	5,5,5	0.97	0	5,5,5	1.10	0
3	GOL	B	1797	-	5,5,5	0.99	0	5,5,5	2.17	2 (40%)
3	GOL	B	1798	-	5,5,5	0.38	0	5,5,5	0.77	0
3	GOL	B	1799	-	5,5,5	0.76	0	5,5,5	1.29	0
3	GOL	B	1800	-	5,5,5	0.28	0	5,5,5	0.54	0
3	GOL	B	1801	-	5,5,5	1.65	1 (20%)	5,5,5	2.59	1 (20%)
3	GOL	B	1802	-	5,5,5	0.46	0	5,5,5	0.66	0
2	47N	C	1792	-	24,24,24	0.87	1 (4%)	35,36,36	1.82	7 (20%)
3	GOL	C	1793	-	5,5,5	0.58	0	5,5,5	0.39	0
3	GOL	C	1794	-	5,5,5	0.88	0	5,5,5	0.41	0
3	GOL	C	1795	-	5,5,5	0.82	0	5,5,5	0.70	0
3	GOL	C	1796	-	5,5,5	0.24	0	5,5,5	0.73	0
3	GOL	C	1797	-	5,5,5	1.18	1 (20%)	5,5,5	0.92	0
2	47N	D	1793	-	24,24,24	1.23	3 (12%)	35,36,36	1.14	3 (8%)
3	GOL	D	1794	-	5,5,5	0.42	0	5,5,5	0.61	0
3	GOL	D	1795	-	5,5,5	0.62	0	5,5,5	1.64	1 (20%)
3	GOL	D	1796	-	5,5,5	0.55	0	5,5,5	0.78	0
3	GOL	D	1797	-	5,5,5	0.51	0	5,5,5	1.09	1 (20%)
3	GOL	D	1798	-	5,5,5	0.28	0	5,5,5	0.53	0
3	GOL	D	1799	-	5,5,5	0.95	1 (20%)	5,5,5	0.80	0
3	GOL	D	1800	-	5,5,5	0.27	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	47N	A	1792	-	-	0/6/51/51	0/1/3/3
3	GOL	A	1793	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1794	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1795	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1796	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1797	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1798	-	-	0/4/4/4	0/0/0/0
2	47N	B	1793	-	-	0/6/51/51	0/1/3/3
3	GOL	B	1794	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1795	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1796	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1797	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1798	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1799	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1800	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1801	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1802	-	-	0/4/4/4	0/0/0/0
2	47N	C	1792	-	-	0/6/51/51	0/1/3/3
3	GOL	C	1793	-	-	0/4/4/4	0/0/0/0
3	GOL	C	1794	-	-	0/4/4/4	0/0/0/0
3	GOL	C	1795	-	-	0/4/4/4	0/0/0/0
3	GOL	C	1796	-	-	0/4/4/4	0/0/0/0
3	GOL	C	1797	-	-	0/4/4/4	0/0/0/0
2	47N	D	1793	-	-	0/6/51/51	0/1/3/3
3	GOL	D	1794	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1795	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1796	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1797	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1798	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1799	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1800	-	-	0/4/4/4	0/0/0/0

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1801	GOL	O2-C2	-3.03	1.34	1.43
3	C	1797	GOL	O2-C2	-2.50	1.36	1.43
3	D	1799	GOL	O2-C2	-2.06	1.37	1.43
2	B	1793	47N	OAJ-CAE	2.10	1.49	1.44
2	B	1793	47N	OAH-CAC	2.35	1.49	1.43

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1801	GOL	O2-C2-C3	-5.25	84.58	108.65
2	C	1792	47N	C6-C5-C4	-4.17	102.72	113.02
2	C	1792	47N	O5-C1-C2	-3.99	103.43	109.80
2	C	1792	47N	C2-C3-C4	-3.95	105.68	110.89
3	A	1794	GOL	C3-C2-C1	-3.65	96.80	111.12

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 46 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1797	GOL	5	0
3	A	1798	GOL	6	0
3	B	1797	GOL	1	0
3	B	1799	GOL	4	0
3	B	1801	GOL	7	0
3	B	1802	GOL	1	0
3	C	1794	GOL	1	0
3	C	1795	GOL	1	0
3	C	1797	GOL	11	0
3	D	1795	GOL	1	0
3	D	1796	GOL	1	0
3	D	1799	GOL	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	740/750 (98%)	-0.50	4 (0%) 91 93	8, 16, 35, 57	0
1	B	741/750 (98%)	-0.48	5 (0%) 89 91	7, 16, 33, 54	0
1	C	740/750 (98%)	-0.34	10 (1%) 78 82	11, 20, 43, 58	0
1	D	741/750 (98%)	-0.40	8 (1%) 82 86	11, 20, 39, 56	0
All	All	2962/3000 (98%)	-0.43	27 (0%) 85 88	7, 18, 38, 58	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	355	ALA	3.2
1	C	791	GLY	3.0
1	C	584	ALA	3.0
1	D	175	LEU	2.8
1	D	110	PHE	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	B	1797	6/6	0.90	0.20	18.92	28,32,35,35	0
3	GOL	A	1797	6/6	0.88	0.18	11.25	10,20,24,29	0
3	GOL	B	1801	6/6	0.92	0.19	9.54	15,21,24,33	0
3	GOL	B	1800	6/6	0.84	0.21	8.45	39,43,44,46	0
3	GOL	D	1800	6/6	0.83	0.16	7.37	41,45,46,47	0
3	GOL	D	1799	6/6	0.93	0.21	7.16	24,31,33,37	0
3	GOL	C	1796	6/6	0.91	0.17	5.63	29,38,39,44	0
3	GOL	C	1794	6/6	0.84	0.25	4.96	26,30,35,35	0
3	GOL	A	1795	6/6	0.81	0.16	4.80	42,43,46,47	0
3	GOL	B	1802	6/6	0.82	0.16	4.13	37,41,45,53	0
3	GOL	D	1798	6/6	0.83	0.21	4.12	41,45,46,47	0
3	GOL	C	1797	6/6	0.93	0.15	3.21	21,23,26,32	0
3	GOL	D	1796	6/6	0.97	0.13	2.62	19,23,24,24	0
3	GOL	D	1795	6/6	0.93	0.16	2.13	22,25,27,28	0
3	GOL	A	1796	6/6	0.96	0.14	2.00	18,22,23,24	0
3	GOL	D	1797	6/6	0.85	0.14	1.90	34,36,38,40	0
4	CA	C	1798	1/1	0.95	0.18	1.80	55,55,55,55	0
3	GOL	A	1798	6/6	0.89	0.21	1.71	20,28,30,34	0
3	GOL	C	1795	6/6	0.94	0.13	1.68	19,22,24,25	0
3	GOL	B	1798	6/6	0.88	0.12	1.43	24,32,32,34	0
3	GOL	B	1799	6/6	0.81	0.17	1.41	29,32,35,35	0
3	GOL	B	1794	6/6	0.98	0.19	1.39	12,14,14,16	0
3	GOL	A	1793	6/6	0.98	0.18	1.22	12,13,14,14	0
3	GOL	B	1795	6/6	0.96	0.16	1.14	18,22,23,24	0
2	47N	A	1792	22/22	0.97	0.14	0.92	12,16,19,28	0
3	GOL	C	1793	6/6	0.98	0.16	0.83	12,12,13,15	0
3	GOL	A	1794	6/6	0.95	0.14	0.76	18,21,22,23	0
3	GOL	B	1796	6/6	0.96	0.10	0.54	14,16,17,17	0
3	GOL	D	1794	6/6	0.97	0.14	0.47	15,16,16,17	0
2	47N	C	1792	22/22	0.98	0.14	0.18	13,17,22,28	0
2	47N	D	1793	22/22	0.98	0.11	0.04	15,19,21,24	0
2	47N	B	1793	22/22	0.97	0.12	-0.21	11,13,17,23	0
4	CA	A	1799	1/1	0.98	0.07	-1.23	34,34,34,34	0
4	CA	D	1801	1/1	0.97	0.06	-1.60	44,44,44,44	0
4	CA	B	1803	1/1	0.99	0.05	-1.82	27,27,27,27	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.