



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:10 PM GMT

PDB ID : 4BQ8
Title : Crystal structure of the RGMB-NEO1 complex form 3
Authors : Bell, C.H.; Healey, E.; van Erp, S.; Bishop, B.; Tang, C.; Gilbert, R.J.C.; Aricescu, A.R.; Pasterkamp, R.J.; Siebold, C.
Deposited on : 2013-05-30
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	 79% 12% • 8%
2	B	122	 9% • 89%
3	C	251	 2% 55% 9% 36%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEOGENIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	0	0
			1540	981	262	291	6			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	880	GLU	-	EXPRESSION TAG	UNP P97798
A	881	THR	-	EXPRESSION TAG	UNP P97798
A	882	GLY	-	EXPRESSION TAG	UNP P97798
A	1084	GLY	-	EXPRESSION TAG	UNP P97798
A	1085	THR	-	EXPRESSION TAG	UNP P97798
A	1086	LYS	-	EXPRESSION TAG	UNP P97798
A	1087	HIS	-	EXPRESSION TAG	UNP P97798
A	1088	HIS	-	EXPRESSION TAG	UNP P97798
A	1089	HIS	-	EXPRESSION TAG	UNP P97798
A	1090	HIS	-	EXPRESSION TAG	UNP P97798
A	1091	HIS	-	EXPRESSION TAG	UNP P97798
A	1092	HIS	-	EXPRESSION TAG	UNP P97798

- Molecule 2 is a protein called RGM DOMAIN FAMILY MEMBER B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	S	0	0	0
			111	74	15	20	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	47	GLU	-	EXPRESSION TAG	UNP Q6NW40
B	48	THR	-	EXPRESSION TAG	UNP Q6NW40
B	49	GLY	-	EXPRESSION TAG	UNP Q6NW40


- Molecule 3 is a protein called RGM DOMAIN FAMILY MEMBER B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	161	Total	C	N	O	S	0	0	0
			1244	780	217	240	7			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	411	GLY	-	EXPRESSION TAG	UNP Q6NW40
C	412	THR	-	EXPRESSION TAG	UNP Q6NW40
C	413	LYS	-	EXPRESSION TAG	UNP Q6NW40
C	414	HIS	-	EXPRESSION TAG	UNP Q6NW40
C	415	HIS	-	EXPRESSION TAG	UNP Q6NW40
C	416	HIS	-	EXPRESSION TAG	UNP Q6NW40
C	417	HIS	-	EXPRESSION TAG	UNP Q6NW40
C	418	HIS	-	EXPRESSION TAG	UNP Q6NW40
C	419	HIS	-	EXPRESSION TAG	UNP Q6NW40
C	225	GLY	GLU	CONFLICT	UNP Q6NW40

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	12	Total 12	O 12	0	0
5	B	1	Total 1	O 1	0	0
5	C	2	Total 2	O 2	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	116.92Å 116.92Å 91.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 82.67 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.80) 99.8 (82.67-2.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.82Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.183 , 0.199 0.188 , 0.208	Depositor DCC
R_{free} test set	813 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	71.3	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 16184 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2924	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/1580	0.73	0/2157
2	B	0.50	0/114	0.67	0/151
3	C	0.52	0/1266	0.80	0/1717
All	All	0.51	0/2960	0.76	0/4025

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1540	0	1529	20	0
2	B	111	0	93	1	0
3	C	1244	0	1219	13	0
4	A	14	0	13	0	0
5	A	12	0	0	0	0
5	B	1	0	0	0	0
5	C	2	0	0	0	0
All	All	2924	0	2854	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:304:GLU:H	3:C:307:GLN:HE21	1.29	0.80
1:A:1005:ILE:HG12	1:A:1049:GLN:HG2	1.75	0.68
1:A:885:MET:HG3	1:A:968:ARG:HB3	1.74	0.68
1:A:993:THR:CG2	3:C:186:ALA:H	2.09	0.65
3:C:259:ARG:HH21	3:C:274:ARG:HD3	1.60	0.64
1:A:993:THR:HG21	3:C:185:GLY:HA3	1.81	0.62
1:A:993:THR:HG21	3:C:186:ALA:H	1.64	0.62
3:C:304:GLU:H	3:C:307:GLN:NE2	1.99	0.61
1:A:992:VAL:HG22	1:A:1063:ILE:HD12	1.86	0.58
1:A:960:SER:HB3	1:A:975:THR:HG22	1.88	0.56
2:B:161:LEU:HD13	3:C:174:PHE:HB3	1.91	0.53
3:C:234:ALA:HB1	3:C:239:LEU:HD22	1.90	0.53
3:C:234:ALA:HB1	3:C:239:LEU:CD2	2.38	0.52
3:C:215:LYS:HD3	3:C:233:GLN:HE21	1.77	0.49
1:A:991:ASP:O	1:A:993:THR:CG2	2.61	0.49
1:A:931:PRO:HD2	1:A:934:THR:CG2	2.45	0.46
1:A:993:THR:HG21	3:C:186:ALA:N	2.31	0.46
1:A:1014:ALA:HB1	1:A:1018:ILE:HD11	2.00	0.44
1:A:994:VAL:HG13	1:A:1004:ILE:HD13	2.01	0.43
1:A:991:ASP:O	1:A:993:THR:HG22	2.18	0.42
1:A:931:PRO:HD2	1:A:934:THR:HG21	2.02	0.42
1:A:948:VAL:HG12	1:A:951:LEU:HD21	2.03	0.41
1:A:989:PRO:HG2	1:A:1063:ILE:HG22	2.02	0.41
1:A:1005:ILE:HD13	3:C:309:LEU:HD22	2.03	0.40
1:A:993:THR:HG23	3:C:186:ALA:H	1.85	0.40
1:A:1024:TYR:HA	1:A:1037:VAL:O	2.21	0.40
1:A:994:VAL:HG22	1:A:1006:VAL:HG13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/213 (90%)	183 (96%)	7 (4%)	1 (0%)	34	69
2	B	10/122 (8%)	9 (90%)	1 (10%)	0	100	100
3	C	155/251 (62%)	144 (93%)	10 (6%)	1 (1%)	30	65
All	All	356/586 (61%)	336 (94%)	18 (5%)	2 (1%)	30	65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	919	SER
3	C	208	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/190 (91%)	167 (96%)	6 (4%)	43	77
2	B	12/103 (12%)	10 (83%)	2 (17%)	3	8
3	C	137/215 (64%)	126 (92%)	11 (8%)	15	40
All	All	322/508 (63%)	303 (94%)	19 (6%)	24	57

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	885	MET
1	A	943	THR
1	A	993	THR
1	A	996	SER
1	A	1022	ILE
1	A	1041	VAL
2	B	140	ASN
2	B	168	ASP
3	C	177	ASN
3	C	189	LEU
3	C	204	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	224	HIS
3	C	235	VAL
3	C	236	THR
3	C	247	THR
3	C	257	SER
3	C	260	ILE
3	C	304	GLU
3	C	306	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	233	GLN
3	C	307	GLN
3	C	326	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	2086	1	14,14,15	0.27	0	15,19,21	0.83	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	2086	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2086	NAG	C1-O5-C5	3.01	116.07	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/213 (91%)	0.23	1 (0%) 91 88	46, 66, 133, 160	0
2	B	14/122 (11%)	0.53	0 100 100	56, 80, 114, 119	0
3	C	161/251 (64%)	0.59	4 (2%) 61 48	49, 70, 111, 135	0
All	All	370/586 (63%)	0.40	5 (1%) 78 69	46, 69, 120, 160	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	324	ASP	3.3
3	C	224	HIS	2.4
3	C	333	LEU	2.4
3	C	329	VAL	2.2
1	A	969	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NAG	A	2086	14/15	0.92	0.16	-0.37	113,116,119,119	0

6.5 Other polymers [i](#)

There are no such residues in this entry.