



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 09:53 PM GMT

PDB ID : 5BR6  
Title : Crystal structure of hemagglutinin of A/Taiwan/2/2013 (H6N1) in complex with LSTc  
Authors : Ni, F.; Kondrashkina, E.; Wang, Q.  
Deposited on : 2015-05-29  
Resolution : 2.43 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1003 (2.46-2.42)
Clashscore	102246	1071 (2.46-2.42)
Ramachandran outliers	100387	1065 (2.46-2.42)
Sidechain outliers	100360	1065 (2.46-2.42)
RSRZ outliers	91569	1005 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	<div> <div></div> <div>84%16%</div> </div>
1	C	326	<div> <div></div> <div>87%13%</div> </div>
2	B	171	<div> <div></div> <div>89%11%</div> </div>
2	D	171	<div> <div></div> <div>91%9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	403	-	-	-	X
3	NAG	B	301	-	-	-	X
3	NAG	C	403	-	-	-	X
3	NAG	D	301	-	-	-	X
6	FUC	B	302	X	-	-	-
6	FUC	D	302	X	-	-	-

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 9064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HEMAGGLUTININ HA1 CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	0	0
			2577	1634	439	491	13			
1	C	326	Total	C	N	O	S	0	0	0
			2577	1634	439	491	13			

- Molecule 2 is a protein called HEMAGGLUTININ HA2 CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	171	Total	C	N	O	S	0	0	0
			1381	860	242	272	7			
2	D	171	Total	C	N	O	S	0	0	0
			1381	860	242	272	7			

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is O-SIALIC ACID (three-letter code: SIA) (formula: C<sub>11</sub>H<sub>19</sub>NO<sub>9</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			20	11	1	8		
4	C	1	Total	C	N	O	0	0
			20	11	1	8		

- Molecule 5 is BETA-D-GALACTOSE (three-letter code: GAL) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	C	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula:  $C_6H_{12}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			10	6	4		
6	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			11	6	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 8 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			11	6	5		
8	B	1	Total	C	O	0	0
			11	6	5		
8	D	1	Total	C	O	0	0
			11	6	5		
8	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	227	Total	O	0	0
			227	227		
9	B	164	Total	O	0	0
			164	164		
9	C	234	Total	O	0	0
			234	234		
9	D	179	Total	O	0	0
			179	179		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HEMAGGLUTININ HA1 CHAIN




- Molecule 1: HEMAGGLUTININ HA1 CHAIN



- Molecule 2: HEMAGGLUTININ HA2 CHAIN



- Molecule 2: HEMAGGLUTININ HA2 CHAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.67Å 113.67Å 163.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.68 – 2.43 37.68 – 2.43	Depositor EDS
% Data completeness (in resolution range)	100.0 (37.68-2.43) 100.0 (37.68-2.43)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.37 (at 2.42Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.203 , 0.225 0.212 , 0.232	Depositor DCC
$R_{free}$ test set	4463 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.2	Xtriage
Anisotropy	0.390	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 26.8	EDS
Estimated twinning fraction	0.458 for -h,-k,l 0.044 for h,-h-k,-l 0.045 for -k,-h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 88848 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9064	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, SIA, GAL, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.23	0/2638	0.40	0/3589
1	C	0.25	1/2638 (0.0%)	0.40	0/3589
2	B	0.22	0/1409	0.36	0/1897
2	D	0.22	0/1409	0.37	0/1897
All	All	0.23	1/8094 (0.0%)	0.39	0/10972

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	132	THR	C-N	-5.82	1.20	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	133	ASN	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2577	0	2527	55	0
1	C	2577	0	2526	44	0
2	B	1381	0	1294	24	0
2	D	1381	0	1294	19	0
3	A	70	0	64	2	0
3	B	28	0	23	1	0
3	C	70	0	64	6	0
3	D	28	0	23	0	0
4	A	20	0	17	0	0
4	C	20	0	17	2	0
5	A	11	0	9	1	0
5	C	11	0	9	2	0
6	B	10	0	10	0	0
6	D	10	0	10	0	0
7	B	11	0	8	0	0
7	D	11	0	8	0	0
8	B	22	0	20	1	0
8	D	22	0	20	0	0
9	A	227	0	0	37	1
9	B	164	0	0	17	3
9	C	234	0	0	28	1
9	D	179	0	0	15	2
All	All	9064	0	7943	147	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 147 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:66:VAL:HG23	9:D:402:HOH:O	1.24	1.33
1:A:123:ALA:HA	9:A:501:HOH:O	1.47	1.12
3:C:407:NAG:H81	9:C:511:HOH:O	1.50	1.12
1:A:124:GLY:N	9:A:501:HOH:O	1.83	1.10
1:A:132:THR:HG23	9:A:506:HOH:O	1.53	1.07

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:B:543:HOH:O	9:B:559:HOH:O[3_675]	1.89	0.31
9:A:614:HOH:O	9:B:442:HOH:O[3_675]	1.95	0.25
9:D:425:HOH:O	9:D:517:HOH:O[3_775]	2.00	0.20
9:C:694:HOH:O	9:D:562:HOH:O[3_775]	2.04	0.16
9:B:538:HOH:O	9:B:561:HOH:O[2_765]	2.06	0.14

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/326 (99%)	315 (97%)	9 (3%)	0	100	100
1	C	324/326 (99%)	313 (97%)	11 (3%)	0	100	100
2	B	169/171 (99%)	166 (98%)	3 (2%)	0	100	100
2	D	169/171 (99%)	166 (98%)	3 (2%)	0	100	100
All	All	986/994 (99%)	960 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/287 (100%)	287 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	287/287 (100%)	287 (100%)	0	100	100
2	B	146/146 (100%)	146 (100%)	0	100	100
2	D	146/146 (100%)	146 (100%)	0	100	100
All	All	866/866 (100%)	866 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	133	ASN
1	C	133	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	401	1	14,14,15	0.89	2 (14%)	15,19,21	1.24	1 (6%)
3	NAG	A	402	1	14,14,15	0.50	0	15,19,21	0.55	0
3	NAG	A	403	1	14,14,15	0.19	0	15,19,21	0.85	1 (6%)
3	NAG	A	404	1	14,14,15	0.18	0	15,19,21	0.34	0
4	SIA	A	405	5	17,20,21	1.08	2 (11%)	18,28,31	1.27	1 (5%)
5	GAL	A	406	3,4	11,11,12	0.67	0	15,15,17	1.00	0
3	NAG	A	407	5	14,14,15	0.20	0	15,19,21	0.34	0
3	NAG	B	301	3,2,6	14,14,15	0.85	1 (7%)	15,19,21	0.95	1 (6%)
6	FUC	B	302	3	10,10,11	0.58	0	13,14,16	0.93	0
3	NAG	B	303	3,7	14,14,15	0.29	0	15,19,21	0.33	0
7	BMA	B	304	8,3	11,11,12	0.95	0	15,15,17	1.40	2 (13%)
8	MAN	B	305	7	11,11,12	0.79	0	15,15,17	1.35	3 (20%)
8	MAN	B	306	7	11,11,12	1.13	2 (18%)	15,15,17	1.71	5 (33%)
3	NAG	C	401	1	14,14,15	0.88	1 (7%)	15,19,21	1.09	1 (6%)
3	NAG	C	402	1	14,14,15	0.72	1 (7%)	15,19,21	0.72	1 (6%)
3	NAG	C	403	1	14,14,15	1.08	1 (7%)	15,19,21	1.27	2 (13%)
3	NAG	C	404	1	14,14,15	0.21	0	15,19,21	0.32	0
4	SIA	C	405	5	17,20,21	1.09	2 (11%)	18,28,31	1.31	1 (5%)
5	GAL	C	406	3,4	11,11,12	0.67	0	15,15,17	0.96	0
3	NAG	C	407	5	14,14,15	0.21	0	15,19,21	0.32	0
3	NAG	D	301	3,2,6	14,14,15	0.29	0	15,19,21	0.86	1 (6%)
6	FUC	D	302	3	10,10,11	0.70	0	13,14,16	1.27	2 (15%)
3	NAG	D	303	3,7	14,14,15	0.29	0	15,19,21	0.55	1 (6%)
7	BMA	D	304	8,3	11,11,12	0.96	1 (9%)	15,15,17	1.25	3 (20%)
8	MAN	D	305	7	11,11,12	0.73	0	15,15,17	1.03	2 (13%)
8	MAN	D	306	7	11,11,12	0.73	0	15,15,17	1.27	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1	-	0/6/23/26	0/1/1/1
3	NAG	A	402	1	-	0/6/23/26	0/1/1/1
3	NAG	A	403	1	-	0/6/23/26	0/1/1/1
3	NAG	A	404	1	-	0/6/23/26	0/1/1/1
4	SIA	A	405	5	-	0/14/34/38	0/1/1/1
5	GAL	A	406	3,4	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	407	5	-	0/6/23/26	0/1/1/1
3	NAG	B	301	3,2,6	-	0/6/23/26	0/1/1/1
6	FUC	B	302	3	1/1/4/5	0/0/17/20	0/1/1/1
3	NAG	B	303	3,7	-	0/6/23/26	0/1/1/1
7	BMA	B	304	8,3	-	0/2/19/22	0/1/1/1
8	MAN	B	305	7	-	0/2/19/22	1/1/1/1
8	MAN	B	306	7	-	0/2/19/22	0/1/1/1
3	NAG	C	401	1	-	0/6/23/26	0/1/1/1
3	NAG	C	402	1	-	0/6/23/26	0/1/1/1
3	NAG	C	403	1	-	0/6/23/26	0/1/1/1
3	NAG	C	404	1	-	0/6/23/26	0/1/1/1
4	SIA	C	405	5	-	0/14/34/38	0/1/1/1
5	GAL	C	406	3,4	-	0/2/19/22	0/1/1/1
3	NAG	C	407	5	-	0/6/23/26	0/1/1/1
3	NAG	D	301	3,2,6	-	0/6/23/26	0/1/1/1
6	FUC	D	302	3	1/1/4/5	0/0/17/20	0/1/1/1
3	NAG	D	303	3,7	-	0/6/23/26	0/1/1/1
7	BMA	D	304	8,3	-	0/2/19/22	0/1/1/1
8	MAN	D	305	7	-	0/2/19/22	1/1/1/1
8	MAN	D	306	7	-	0/2/19/22	1/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	NAG	O5-C1	-3.00	1.38	1.43
3	C	402	NAG	O5-C1	-2.42	1.39	1.43
4	C	405	SIA	O6-C6	-2.14	1.40	1.43
4	A	405	SIA	O6-C6	-2.08	1.40	1.43
7	D	304	BMA	O3-C3	2.08	1.47	1.43

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	405	SIA	C7-C6-C5	-3.19	109.66	114.06
4	A	405	SIA	C7-C6-C5	-3.12	109.76	114.06
8	D	306	MAN	O2-C2-C3	-2.31	105.54	110.19
8	B	305	MAN	O2-C2-C3	-2.30	105.55	110.19
8	D	305	MAN	O2-C2-C3	-2.29	105.58	110.19

All (2) chirality outliers are listed below:



Mol	Chain	Res	Type	Atom
6	D	302	FUC	C1
6	B	302	FUC	C1

There are no torsion outliers.

All (3) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	305	MAN	C1-C2-C3-C4-C5-O5
8	D	305	MAN	C1-C2-C3-C4-C5-O5
8	D	306	MAN	C1-C2-C3-C4-C5-O5

9 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	NAG	2	0
5	A	406	GAL	1	0
3	B	301	NAG	1	0
8	B	306	MAN	1	0
3	C	401	NAG	1	0
3	C	402	NAG	1	0
4	C	405	SIA	2	0
5	C	406	GAL	2	0
3	C	407	NAG	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	326/326 (100%)	-0.38	2 (0%) 90 91	17, 39, 64, 104	0
1	C	326/326 (100%)	-0.37	2 (0%) 90 91	17, 39, 63, 99	0
2	B	171/171 (100%)	-0.38	0 100 100	15, 26, 47, 77	0
2	D	171/171 (100%)	-0.37	0 100 100	14, 27, 46, 74	0
All	All	994/994 (100%)	-0.38	4 (0%) 93 94	14, 33, 61, 104	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	155	ASP	4.9
1	A	156	SER	3.4
1	A	155	ASP	2.8
1	C	156	SER	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	C	403	14/15	0.87	0.14	2.91	51,54,57,57	0
3	NAG	B	301	14/15	0.84	0.18	2.61	37,41,44,45	0
3	NAG	A	403	14/15	0.84	0.17	2.24	50,54,56,57	0
3	NAG	D	301	14/15	0.81	0.17	2.02	40,42,44,45	0
4	SIA	C	405	20/21	0.93	0.15	-0.10	50,59,70,71	0
4	SIA	A	405	20/21	0.93	0.14	-0.22	51,60,71,73	0
8	MAN	B	306	11/12	0.62	0.40	-	52,53,54,54	0
7	BMA	D	304	11/12	0.70	0.35	-	51,51,52,52	0
8	MAN	D	306	11/12	0.57	0.35	-	51,51,52,52	0
5	GAL	C	406	11/12	0.77	0.27	-	68,92,101,101	0
3	NAG	D	303	14/15	0.82	0.18	-	46,47,50,53	0
8	MAN	D	305	11/12	0.70	0.38	-	51,52,52,52	0
3	NAG	C	401	14/15	0.78	0.29	-	39,42,43,43	0
3	NAG	B	303	14/15	0.81	0.16	-	46,49,51,52	0
3	NAG	A	401	14/15	0.72	0.29	-	40,45,45,45	0
6	FUC	D	302	10/11	0.86	0.15	-	45,45,46,47	0
3	NAG	A	404	14/15	0.78	0.35	-	50,52,52,53	0
3	NAG	A	407	14/15	0.85	0.36	-	74,101,108,111	0
3	NAG	C	407	14/15	0.87	0.33	-	75,100,111,112	0
8	MAN	B	305	11/12	0.68	0.39	-	53,54,54,54	0
5	GAL	A	406	11/12	0.76	0.26	-	67,91,100,102	0
3	NAG	A	402	14/15	0.66	0.27	-	47,48,48,51	0
6	FUC	B	302	10/11	0.83	0.16	-	44,45,45,46	0
3	NAG	C	402	14/15	0.72	0.24	-	46,47,48,48	0
7	BMA	B	304	11/12	0.68	0.31	-	52,53,54,54	0
3	NAG	C	404	14/15	0.79	0.28	-	47,51,51,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.