



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:11 PM GMT

PDB ID : 4BSK
Title : Crystal structure of VEGF-C in complex with VEGFR-3 domains D1-2
Authors : Leppanen, V.M.; Tvorogov, D.; Kisko, K.; Protá, A.E.; Jeltsch, M.; Anisimov, A.; Markovic-Mueller, S.; Stutfeld, E.; Goldie, K.N.; Ballmer-Hofer, K.; Alitalo, K.
Deposited on : 2013-06-10
Resolution : 4.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1039 (4.80-3.60)
Clashscore	102246	1140 (4.80-3.60)
Ramachandran outliers	100387	1083 (4.80-3.60)
Sidechain outliers	100360	1067 (4.80-3.60)
RSRZ outliers	91569	1042 (4.80-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div>3%</div> <div>62%</div> <div>17%</div> <div>20%</div> </div>
2	C	121	<div> <div>%</div> <div>74%</div> <div>6%</div> <div>19%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	400	X	-	-	-
3	NAG	C	600	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1876 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	172	Total	C	N	O	S	0	0	0
			1117	705	187	220	5			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	230	ALA	-	CLONING ARTIFACT	UNP P35916
A	231	ASP	-	CLONING ARTIFACT	UNP P35916
A	232	PRO	-	CLONING ARTIFACT	UNP P35916
A	233	ILE	-	CLONING ARTIFACT	UNP P35916
A	234	GLU	-	CLONING ARTIFACT	UNP P35916
A	235	GLY	-	CLONING ARTIFACT	UNP P35916
A	236	ARG	-	CLONING ARTIFACT	UNP P35916

- Molecule 2 is a protein called VASCULAR ENDOTHELIAL GROWTH FACTOR C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	98	Total	C	N	O	S	0	0	0
			675	414	116	137	8			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	101	ASP	-	EXPRESSION TAG	UNP P49767
C	102	PRO	-	EXPRESSION TAG	UNP P49767
C	137	ALA	CYS	ENGINEERED MUTATION	UNP P49767
C	216	HIS	-	EXPRESSION TAG	UNP P49767
C	217	HIS	-	EXPRESSION TAG	UNP P49767
C	218	HIS	-	EXPRESSION TAG	UNP P49767
C	219	HIS	-	EXPRESSION TAG	UNP P49767
C	220	HIS	-	EXPRESSION TAG	UNP P49767

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	221	HIS	-	EXPRESSION TAG	UNP P49767

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	2	Total	C	N	O	0	0
			28	16	2	10		

• Molecule 1: VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR 3



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	166.66Å 166.66Å 166.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.92 – 4.20 19.92 – 4.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.92-4.20) 100.0 (19.92-4.20)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 4.21Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.334 , 0.372 0.331 , 0.366	Depositor DCC
R_{free} test set	568 reflections (9.98%)	DCC
Wilson B-factor (Å ²)	241.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	4.31 , 5180.2	EDS
Estimated twinning fraction	0.053 for -l,-k,-h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 5721 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	1876	wwPDB-VP
Average B, all atoms (Å ²)	135.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/1141	0.48	0/1583
2	C	0.27	0/688	0.52	0/941
All	All	0.26	0/1829	0.50	0/2524

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1117	0	880	23	0
2	C	675	0	573	7	0
3	A	42	0	39	1	0
3	C	14	0	13	0	0
4	C	28	0	25	1	0
All	All	1876	0	1530	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:THR:OG1	1:A:102:HIS:HA	1.95	0.66
1:A:139:PHE:HE1	1:A:158:CYS:HB2	1.63	0.63
2:C:162:CYS:H	2:C:206:HIS:HD2	1.49	0.59
1:A:105:ASP:O	1:A:109:TYR:OH	2.23	0.57
1:A:171:SER:HB3	1:A:204:LEU:HD23	1.87	0.56
1:A:194:SER:HB2	1:A:197:LEU:HD23	1.87	0.55
2:C:162:CYS:H	2:C:206:HIS:CD2	2.23	0.55
1:A:75:SER:O	1:A:90:ARG:N	2.40	0.54
1:A:141:ASN:HD22	1:A:142:LYS:H	1.56	0.54
1:A:185:TRP:NE1	1:A:187:ASP:OD1	2.42	0.53
1:A:33:ASN:O	3:A:400:NAG:O6	2.24	0.51
1:A:155:TRP:HH2	1:A:186:ASP:HB3	1.76	0.50
1:A:36:GLU:HB2	1:A:39:HIS:CE1	2.47	0.50
1:A:62:PRO:HD2	1:A:109:TYR:CE2	2.49	0.47
1:A:101:VAL:HG12	1:A:132:VAL:HG11	1.99	0.45
2:C:175:ASN:O	2:C:175:ASN:ND2	2.51	0.44
1:A:169:LEU:HD21	1:A:204:LEU:HD13	1.98	0.44
1:A:193:VAL:HB	1:A:198:LEU:HD23	1.99	0.44
1:A:156:VAL:HG12	1:A:191:MET:O	2.18	0.44
1:A:106:THR:HG23	1:A:132:VAL:H	1.82	0.43
1:A:103:ALA:C	1:A:105:ASP:H	2.22	0.43
4:C:500:NAG:H62	4:C:501:NAG:O5	2.18	0.43
2:C:149:ASN:HB2	2:C:190:VAL:CG2	2.49	0.42
1:A:172:GLN:HB3	1:A:203:TYR:HE1	1.84	0.42
1:A:154:MET:CB	1:A:198:LEU:HD21	2.49	0.41
2:C:189:THR:HG22	2:C:190:VAL:H	1.85	0.41
1:A:141:ASN:OD1	1:A:218:ASN:N	2.44	0.41
1:A:169:LEU:HD11	1:A:204:LEU:HB3	2.03	0.41
1:A:141:ASN:ND2	1:A:142:LYS:H	2.18	0.40
2:C:189:THR:HG22	2:C:190:VAL:N	2.37	0.40
2:C:146:VAL:HG21	2:C:152:PHE:HE2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/214 (77%)	143 (87%)	19 (12%)	2 (1%)	16	62
2	C	96/121 (79%)	92 (96%)	4 (4%)	0	100	100
All	All	260/335 (78%)	235 (90%)	23 (9%)	2 (1%)	24	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	PRO
1	A	104	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/187 (49%)	88 (96%)	4 (4%)	35	72
2	C	69/109 (63%)	68 (99%)	1 (1%)	74	89
All	All	161/296 (54%)	156 (97%)	5 (3%)	47	78

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	139	PHE
1	A	140	ILE
1	A	141	ASN
1	A	152	ASP
2	C	175	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	206	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	500	2,4	14,14,15	0.49	0	15,19,21	1.13	1 (6%)
4	NAG	C	501	4	14,14,15	0.53	0	15,19,21	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	500	2,4	-	0/6/23/26	0/1/1/1
4	NAG	C	501	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	500	NAG	C1-O5-C5	3.75	117.00	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	500	NAG	1	0
4	C	501	NAG	1	0

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	A	400	1	14,14,15	0.51	0	15,19,21	0.71	0
3	NAG	A	500	1	14,14,15	0.50	0	15,19,21	2.27	2 (13%)
3	NAG	A	600	1	14,14,15	0.50	0	15,19,21	1.06	1 (6%)
3	NAG	C	600	2	14,14,15	0.61	0	15,19,21	1.32	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	400	1	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	A	500	1	-	0/6/23/26	0/1/1/1
3	NAG	A	600	1	-	0/6/23/26	0/1/1/1
3	NAG	C	600	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	NAG	C4-C3-C2	-2.31	107.63	111.23
3	A	600	NAG	C1-O5-C5	2.76	115.75	112.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	600	NAG	C1-O5-C5	4.54	118.00	112.25
3	A	500	NAG	C1-O5-C5	7.93	122.31	112.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	400	NAG	C1

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	400	NAG	C8-C7-N2-C2
3	A	400	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	400	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/214 (80%)	-0.23	6 (3%) 48 37	57, 130, 231, 288	0
2	C	98/121 (80%)	-0.47	1 (1%) 84 77	74, 122, 236, 289	0
All	All	270/335 (80%)	-0.32	7 (2%) 59 48	57, 128, 233, 289	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	70	THR	4.4
1	A	68	PRO	3.5
1	A	95	VAL	3.4
2	C	164	GLY	2.5
1	A	53	GLY	2.2
1	A	60	ALA	2.1
1	A	136	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAG	C	500	14/15	0.93	0.24	0.09	144,144,144,144	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NAG	C	501	14/15	0.82	0.25	-	199,199,199,199	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	C	600	14/15	0.70	0.50	3.04	167,167,167,167	0
3	NAG	A	400	14/15	0.82	0.23	-0.61	215,215,215,215	0
3	NAG	A	600	14/15	0.85	0.14	-	147,147,147,147	0
3	NAG	A	500	14/15	0.72	0.39	-	222,222,222,222	0

6.5 Other polymers [i](#)

There are no such residues in this entry.