



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 11:28 PM GMT

PDB ID : 5BSF
Title : Crystal structure of Medicago truncatula (delta)1-Pyrroline-5-Carboxylate Reductase (MtP5CR) in complex with NAD+
Authors : Ruszkowski, M.; Nocek, B.; Forlani, G.; Dauter, Z.
Deposited on : 2015-06-02
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>.</div> </div> </div>
1	B	277	<div> <div>3%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>..</div> </div> </div>
1	C	277	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>..</div> </div> </div>
1	D	277	<div> <div>7%</div> <div> <div></div> <div>87%</div> <div>11%</div> <div>.</div> </div> </div>
1	E	277	<div> <div>3%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>..</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	277	
1	G	277	
1	H	277	
1	I	277	
1	J	277	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MPO	B	302	-	-	-	X
3	MPO	C	302	-	-	-	X
3	MPO	J	302	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 22411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyrroline-5-carboxylate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	4	0
			2017	1272	350	388	7			
1	B	272	Total	C	N	O	S	0	5	0
			2028	1279	352	390	7			
1	C	270	Total	C	N	O	S	0	5	0
			2003	1263	348	385	7			
1	D	271	Total	C	N	O	S	0	5	0
			2019	1270	351	392	6			
1	E	272	Total	C	N	O	S	0	7	0
			2034	1284	353	391	6			
1	F	272	Total	C	N	O	S	0	5	0
			2026	1277	354	389	6			
1	G	268	Total	C	N	O	S	4	4	0
			1987	1250	347	383	7			
1	H	272	Total	C	N	O	S	0	5	0
			2024	1278	352	388	6			
1	I	272	Total	C	N	O	S	0	2	0
			2004	1266	349	383	6			
1	J	272	Total	C	N	O	S	0	4	0
			2019	1274	351	388	6			

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP G7KRM5
A	-1	ASN	-	expression tag	UNP G7KRM5
A	0	ALA	-	expression tag	UNP G7KRM5
B	-2	SER	-	expression tag	UNP G7KRM5
B	-1	ASN	-	expression tag	UNP G7KRM5
B	0	ALA	-	expression tag	UNP G7KRM5
C	-2	SER	-	expression tag	UNP G7KRM5
C	-1	ASN	-	expression tag	UNP G7KRM5
C	0	ALA	-	expression tag	UNP G7KRM5

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	SER	-	expression tag	UNP G7KRM5
D	-1	ASN	-	expression tag	UNP G7KRM5
D	0	ALA	-	expression tag	UNP G7KRM5
E	-2	SER	-	expression tag	UNP G7KRM5
E	-1	ASN	-	expression tag	UNP G7KRM5
E	0	ALA	-	expression tag	UNP G7KRM5
F	-2	SER	-	expression tag	UNP G7KRM5
F	-1	ASN	-	expression tag	UNP G7KRM5
F	0	ALA	-	expression tag	UNP G7KRM5
G	-2	SER	-	expression tag	UNP G7KRM5
G	-1	ASN	-	expression tag	UNP G7KRM5
G	0	ALA	-	expression tag	UNP G7KRM5
H	-2	SER	-	expression tag	UNP G7KRM5
H	-1	ASN	-	expression tag	UNP G7KRM5
H	0	ALA	-	expression tag	UNP G7KRM5
I	-2	SER	-	expression tag	UNP G7KRM5
I	-1	ASN	-	expression tag	UNP G7KRM5
I	0	ALA	-	expression tag	UNP G7KRM5
J	-2	SER	-	expression tag	UNP G7KRM5
J	-1	ASN	-	expression tag	UNP G7KRM5
J	0	ALA	-	expression tag	UNP G7KRM5

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	E	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	F	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	G	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	H	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	I	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	J	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (three-letter code: MPO) (formula: C₇H₁₅NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	C	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	D	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	E	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	F	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	G	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	H	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	I	1	Total 13	C 7	N 1	O 4	S 1	0	0
3	J	1	Total 13	C 7	N 1	O 4	S 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	1	Total 1	Cl 1	0	0
4	J	1	Total 1	Cl 1	0	0
4	D	1	Total 1	Cl 1	0	0
4	E	1	Total 1	Cl 1	0	0
4	H	1	Total 1	Cl 1	0	0
4	B	1	Total 1	Cl 1	0	0
4	I	1	Total 1	Cl 1	0	0
4	C	1	Total 1	Cl 1	0	0
4	A	1	Total 1	Cl 1	0	0
4	F	1	Total 1	Cl 1	0	0



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	199	Total 199	O 199	0	0
5	B	195	Total 195	O 195	0	0
5	C	171	Total 171	O 171	0	0
5	D	120	Total 120	O 120	0	0
5	E	184	Total 184	O 184	0	0
5	F	198	Total 198	O 198	0	0
5	G	112	Total 112	O 112	0	0
5	H	156	Total 156	O 156	0	0
5	I	183	Total 183	O 183	0	0
5	J	151	Total 152	O 152	0	1



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Pyrroline-5-carboxylate reductase





- Molecule 1: Pyrroline-5-carboxylate reductase



- Molecule 1: Pyrroline-5-carboxylate reductase



- Molecule 1: Pyrroline-5-carboxylate reductase




- Molecule 1: Pyrroline-5-carboxylate reductase









- Molecule 1: Pyrroline-5-carboxylate reductase




- Molecule 1: Pyrroline-5-carboxylate reductase





- Molecule 1: Pyrroline-5-carboxylate reductase



- Molecule 1: Pyrroline-5-carboxylate reductase



- Molecule 1: Pyrroline-5-carboxylate reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	87.54Å 100.78Å 101.61Å 67.50° 85.86° 89.78°	Depositor
Resolution (Å)	39.13 – 1.85 39.13 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.5 (39.13-1.85) 85.8 (39.13-1.85)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.85Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.155 , 0.177 0.169 , 0.189	Depositor DCC
R_{free} test set	2131 reflections (0.81%)	DCC
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.4	EDS
Estimated twinning fraction	0.008 for -h,l,k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 266309 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	22411	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MPO, NAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.02	1/2046 (0.0%)	0.88	1/2768 (0.0%)
1	B	1.01	4/2057 (0.2%)	0.90	7/2782 (0.3%)
1	C	0.94	1/2035 (0.0%)	0.86	2/2752 (0.1%)
1	D	0.91	3/2047 (0.1%)	0.87	5/2766 (0.2%)
1	E	1.00	3/2066 (0.1%)	0.92	2/2796 (0.1%)
1	F	1.03	1/2058 (0.0%)	0.89	3/2784 (0.1%)
1	G	0.90	1/2015 (0.0%)	0.87	4/2724 (0.1%)
1	H	1.00	4/2056 (0.2%)	0.88	2/2780 (0.1%)
1	I	0.94	1/2033 (0.0%)	0.89	4/2749 (0.1%)
1	J	0.97	1/2048 (0.0%)	0.89	5/2770 (0.2%)
All	All	0.97	20/20461 (0.1%)	0.89	35/27671 (0.1%)

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	78	SER	CB-OG	5.95	1.50	1.42
1	B	272	GLU	CD-OE1	5.68	1.31	1.25
1	A	155	SER	CB-OG	-5.63	1.34	1.42
1	H	272	GLU	CD-OE1	5.53	1.31	1.25
1	B	48	PRO	N-CD	5.50	1.55	1.47

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	249	GLU	OE1-CD-OE2	5.97	130.47	123.30
1	J	6	ILE	CA-CB-CG1	-5.93	99.74	111.00
1	A	249	GLU	OE1-CD-OE2	5.87	130.34	123.30
1	I	205	ASP	CB-CG-OD1	5.85	123.56	118.30
1	D	249	GLU	OE1-CD-OE2	5.77	130.22	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2017	0	2080	12	0
1	B	2028	0	2092	14	0
1	C	2003	0	2066	14	0
1	D	2019	0	2071	29	0
1	E	2034	0	2100	20	0
1	F	2026	0	2092	14	0
1	G	1987	0	2046	28	0
1	H	2024	0	2098	16	0
1	I	2004	0	2077	17	0
1	J	2019	0	2086	22	0
2	A	44	0	24	1	0
2	B	44	0	24	0	0
2	C	44	0	24	1	0
2	D	44	0	24	1	0
2	E	44	0	24	2	0
2	F	44	0	23	2	0
2	G	44	0	25	1	0
2	H	44	0	25	1	0
2	I	44	0	24	1	0
2	J	44	0	22	2	0
3	A	13	0	15	0	0
3	B	13	0	15	0	0
3	C	13	0	15	0	0
3	D	13	0	15	0	0
3	E	13	0	15	0	0
3	F	13	0	15	0	0
3	G	13	0	15	0	0
3	H	13	0	15	0	0
3	I	13	0	15	0	0
3	J	13	0	15	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
5	A	199	0	0	3	0
5	B	195	0	0	3	0
5	C	171	0	0	3	0
5	D	120	0	0	3	0
5	E	184	0	0	5	0
5	F	198	0	0	5	0
5	G	112	0	0	8	0
5	H	156	0	0	5	0
5	I	183	0	0	7	0
5	J	152	0	0	3	0
All	All	22411	0	21197	177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:110:MET:HE2	1:G:143:GLY:CA	1.68	1.23
1:B:168[B]:ASP:OD1	5:B:402:HOH:O	1.56	1.18
1:D:168[A]:ASP:OD1	5:D:404:HOH:O	1.67	1.12
1:J:168[A]:ASP:OD1	5:J:401:HOH:O	1.70	1.09
1:G:110:MET:CE	1:G:143:GLY:CA	2.32	1.07

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/277 (99%)	271 (99%)	3 (1%)	0	100	100
1	B	275/277 (99%)	272 (99%)	3 (1%)	0	100	100
1	C	273/277 (99%)	270 (99%)	3 (1%)	0	100	100
1	D	273/277 (99%)	269 (98%)	4 (2%)	0	100	100
1	E	277/277 (100%)	274 (99%)	3 (1%)	0	100	100
1	F	275/277 (99%)	272 (99%)	3 (1%)	0	100	100
1	G	270/277 (98%)	266 (98%)	4 (2%)	0	100	100
1	H	275/277 (99%)	271 (98%)	4 (2%)	0	100	100
1	I	272/277 (98%)	268 (98%)	4 (2%)	0	100	100
1	J	274/277 (99%)	271 (99%)	3 (1%)	0	100	100
All	All	2738/2770 (99%)	2704 (99%)	34 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/212 (100%)	207 (98%)	5 (2%)	57	39
1	B	213/212 (100%)	208 (98%)	5 (2%)	58	41
1	C	210/212 (99%)	205 (98%)	5 (2%)	57	39
1	D	212/212 (100%)	210 (99%)	2 (1%)	84	79
1	E	214/212 (101%)	210 (98%)	4 (2%)	65	49
1	F	213/212 (100%)	208 (98%)	5 (2%)	58	41
1	G	208/212 (98%)	201 (97%)	7 (3%)	44	24
1	H	213/212 (100%)	206 (97%)	7 (3%)	45	25
1	I	210/212 (99%)	206 (98%)	4 (2%)	65	49

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	212/212 (100%)	209 (99%)	3 (1%)	74	63
All	All	2117/2120 (100%)	2070 (98%)	47 (2%)	58	43

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	46	SER
1	G	9	ASP
1	I	233	LYS
1	F	78	SER
1	G	12	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	114	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 30 ligands modelled in this entry, 10 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	A	301	-	42,48,48	1.74	12 (28%)	46,73,73	1.80	10 (21%)
3	MPO	A	302	-	13,13,13	2.01	3 (23%)	16,17,17	2.63	5 (31%)
2	NAD	B	301	-	42,48,48	1.73	10 (23%)	46,73,73	1.82	10 (21%)
3	MPO	B	302	-	13,13,13	1.44	2 (15%)	16,17,17	2.87	5 (31%)
2	NAD	C	301	-	42,48,48	1.70	11 (26%)	46,73,73	1.87	13 (28%)
3	MPO	C	302	-	13,13,13	2.06	3 (23%)	16,17,17	3.27	8 (50%)
2	NAD	D	301	-	42,48,48	1.76	9 (21%)	46,73,73	1.87	10 (21%)
3	MPO	D	302	-	13,13,13	1.68	3 (23%)	16,17,17	2.62	6 (37%)
2	NAD	E	301	-	42,48,48	1.79	9 (21%)	46,73,73	1.77	12 (26%)
3	MPO	E	302	-	13,13,13	1.74	3 (23%)	16,17,17	1.59	5 (31%)
2	NAD	F	301	-	42,48,48	1.93	14 (33%)	46,73,73	1.76	10 (21%)
3	MPO	F	302	-	13,13,13	1.71	3 (23%)	16,17,17	2.45	3 (18%)
2	NAD	G	301	-	42,48,48	1.80	12 (28%)	46,73,73	1.82	12 (26%)
3	MPO	G	302	-	13,13,13	1.64	3 (23%)	16,17,17	2.97	8 (50%)
2	NAD	H	301	-	42,48,48	1.88	11 (26%)	46,73,73	1.75	11 (23%)
3	MPO	H	302	-	13,13,13	1.97	3 (23%)	16,17,17	1.84	5 (31%)
2	NAD	I	301	-	42,48,48	1.64	8 (19%)	46,73,73	1.78	11 (23%)
3	MPO	I	302	-	13,13,13	1.53	2 (15%)	16,17,17	2.79	5 (31%)
2	NAD	J	301	-	42,48,48	1.91	12 (28%)	46,73,73	1.82	11 (23%)
3	MPO	J	302	-	13,13,13	1.71	3 (23%)	16,17,17	2.07	5 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	301	-	-	0/22/62/62	0/5/5/5
3	MPO	A	302	-	-	0/7/15/15	0/1/1/1
2	NAD	B	301	-	-	0/22/62/62	0/5/5/5
3	MPO	B	302	-	-	0/7/15/15	0/1/1/1
2	NAD	C	301	-	-	0/22/62/62	0/5/5/5
3	MPO	C	302	-	-	0/7/15/15	0/1/1/1
2	NAD	D	301	-	-	0/22/62/62	0/5/5/5
3	MPO	D	302	-	-	0/7/15/15	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	E	301	-	-	0/22/62/62	0/5/5/5
3	MPO	E	302	-	-	0/7/15/15	0/1/1/1
2	NAD	F	301	-	-	0/22/62/62	0/5/5/5
3	MPO	F	302	-	-	0/7/15/15	0/1/1/1
2	NAD	G	301	-	-	0/22/62/62	0/5/5/5
3	MPO	G	302	-	-	0/7/15/15	0/1/1/1
2	NAD	H	301	-	-	0/22/62/62	0/5/5/5
3	MPO	H	302	-	-	0/7/15/15	0/1/1/1
2	NAD	I	301	-	-	0/22/62/62	0/5/5/5
3	MPO	I	302	-	-	0/7/15/15	0/1/1/1
2	NAD	J	301	-	-	0/22/62/62	0/5/5/5
3	MPO	J	302	-	-	0/7/15/15	0/1/1/1

The worst 5 of 136 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	301	NAD	O3D-C3D	-5.46	1.30	1.43
2	E	301	NAD	O3D-C3D	-4.94	1.31	1.43
2	I	301	NAD	O3D-C3D	-4.36	1.32	1.43
2	B	301	NAD	O3D-C3D	-4.32	1.32	1.43
2	F	301	NAD	O2B-C2B	-3.83	1.33	1.43

The worst 5 of 165 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	MPO	O1-S1-C1	-8.18	101.09	106.87
3	A	302	MPO	O1-S1-C1	-6.96	101.95	106.87
2	I	301	NAD	N3A-C2A-N1A	-6.90	123.45	128.87
2	C	301	NAD	N3A-C2A-N1A	-6.55	123.73	128.87
3	I	302	MPO	O2-S1-O1	-6.54	95.49	113.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	NAD	1	0
2	C	301	NAD	1	0
2	D	301	NAD	1	0
2	E	301	NAD	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	NAD	2	0
2	G	301	NAD	1	0
2	H	301	NAD	1	0
2	I	301	NAD	1	0
2	J	301	NAD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	272/277 (98%)	-0.38	4 (1%)	76 76	19, 31, 57, 85	0
1	B	272/277 (98%)	-0.19	9 (3%)	50 47	19, 33, 62, 111	0
1	C	270/277 (97%)	-0.27	3 (1%)	82 82	19, 35, 64, 104	0
1	D	271/277 (97%)	0.10	20 (7%)	17 17	20, 44, 89, 112	0
1	E	272/277 (98%)	-0.12	8 (2%)	55 52	20, 34, 62, 113	0
1	F	272/277 (98%)	-0.44	3 (1%)	82 82	20, 31, 57, 109	0
1	G	268/277 (96%)	0.53	39 (14%)	3 3	21, 54, 100, 141	1 (0%)
1	H	272/277 (98%)	-0.14	13 (4%)	34 32	20, 39, 69, 109	0
1	I	272/277 (98%)	-0.22	10 (3%)	45 43	20, 34, 68, 105	0
1	J	272/277 (98%)	-0.13	13 (4%)	34 32	20, 38, 74, 117	0
All	All	2713/2770 (97%)	-0.13	122 (4%)	37 35	19, 36, 81, 141	1 (0%)

The worst 5 of 122 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	8	ALA	10.2
1	J	3	ILE	7.5
1	H	3	ILE	7.0
1	G	48	PRO	6.8
1	E	4	ILE	6.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MPO	B	302	13/13	0.91	0.13	3.31	41,49,66,67	0
3	MPO	J	302	13/13	0.94	0.15	2.57	40,44,59,60	0
3	MPO	C	302	13/13	0.94	0.14	2.43	38,47,58,60	0
3	MPO	F	302	13/13	0.94	0.11	1.92	30,39,50,52	0
3	MPO	D	302	13/13	0.93	0.14	1.63	42,49,73,74	0
3	MPO	G	302	13/13	0.91	0.17	1.46	50,64,79,81	0
2	NAD	A	301	44/44	0.91	0.15	1.40	30,43,50,56	0
2	NAD	G	301	44/44	0.85	0.25	1.38	49,75,84,92	0
2	NAD	F	301	44/44	0.90	0.14	1.35	31,42,53,56	0
2	NAD	D	301	44/44	0.88	0.19	1.23	41,59,71,74	0
3	MPO	I	302	13/13	0.95	0.11	1.23	36,40,53,55	0
2	NAD	J	301	44/44	0.88	0.19	1.06	33,53,70,74	0
3	MPO	E	302	13/13	0.91	0.14	0.90	40,48,58,58	0
2	NAD	C	301	44/44	0.92	0.14	0.76	37,47,57,61	0
2	NAD	H	301	44/44	0.91	0.16	0.61	30,48,62,65	0
2	NAD	E	301	44/44	0.87	0.16	0.58	35,56,65,75	0
3	MPO	A	302	13/13	0.96	0.10	0.42	33,42,48,48	0
2	NAD	I	301	44/44	0.92	0.12	0.25	29,42,50,50	0
3	MPO	H	302	13/13	0.95	0.09	-0.13	37,39,48,50	0
2	NAD	B	301	44/44	0.92	0.12	-0.14	30,47,58,61	0
4	CL	A	303	1/1	0.97	0.09	-0.89	39,39,39,39	0
4	CL	H	303	1/1	0.97	0.06	-0.93	45,45,45,45	0
4	CL	C	303	1/1	0.98	0.07	-1.03	36,36,36,36	0
4	CL	B	303	1/1	0.99	0.04	-1.84	33,33,33,33	0
4	CL	J	303	1/1	0.98	0.07	-1.98	39,39,39,39	0
4	CL	D	303	1/1	0.98	0.05	-2.15	37,37,37,37	0
4	CL	F	303	1/1	0.99	0.07	-2.34	37,37,37,37	0
4	CL	I	303	1/1	0.99	0.04	-2.44	40,40,40,40	0
4	CL	E	303	1/1	0.99	0.05	-2.85	34,34,34,34	0
4	CL	G	303	1/1	0.98	0.04	-3.24	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.