



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:19 PM GMT

PDB ID : 4BUR  
Title : Crystal structure of the reduced human Apoptosis inducing factor complexed with NAD  
Authors : Martinez-Julvez, M.; Herguedas, B.; Hermoso, J.A.; Ferreira, P.; Villanueva, R.; Medina, M.  
Deposited on : 2013-06-23  
Resolution : 2.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1945 (2.90-2.86)
Clashscore	102246	2202 (2.90-2.86)
Ramachandran outliers	100387	2149 (2.90-2.86)
Sidechain outliers	100360	2152 (2.90-2.86)
RSRZ outliers	91569	1950 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	 75% 12% • 12%
1	B	511	 74% 13% • 12%
1	C	511	 74% 13% • 13%
1	D	511	 2% 71% 13% • 15%

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 14362 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APOPTOSIS INDUCING FACTOR 1, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	450	Total 3475	C 2208	N 618	O 638	S 11	0	0	0
1	B	450	Total 3471	C 2205	N 616	O 639	S 11	0	0	0
1	C	446	Total 3452	C 2194	N 613	O 634	S 11	0	0	0
1	D	435	Total 3371	C 2147	N 599	O 614	S 11	0	0	0

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	A	1	Total 44	C 21	N 7	O 14	P 2	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	9	Total	O	0	0
			9	9		
5	B	8	Total	O	0	0
			8	8		
5	C	7	Total	O	0	0
			7	7		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: APOPTOSIS INDUCING FACTOR 1, MITOCHONDRIAL



- Molecule 1: APOPTOSIS INDUCING FACTOR 1, MITOCHONDRIAL



- Molecule 1: APOPTOSIS INDUCING FACTOR 1, MITOCHONDRIAL





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.77Å 120.77Å 343.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.88 77.21 – 2.88	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.00-2.88) 99.7 (77.21-2.88)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.81 (at 2.86Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.176 , 0.230 0.174 , 0.226	Depositor DCC
$R_{free}$ test set	4785 reflections (7.77%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.4	Xtriage
Anisotropy	0.530	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 38.5	EDS
Estimated twinning fraction	0.044 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 66422 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	14362	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAD, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	3/3546 (0.1%)	0.71	2/4794 (0.0%)
1	B	0.55	2/3540 (0.1%)	0.67	0/4785
1	C	0.55	5/3522 (0.1%)	0.68	2/4760 (0.0%)
1	D	0.54	3/3440 (0.1%)	0.64	0/4648
All	All	0.56	13/14048 (0.1%)	0.67	4/18987 (0.0%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	477	TRP	CD2-CE2	5.73	1.48	1.41
1	B	483	TRP	CD2-CE2	5.60	1.48	1.41
1	D	351	TRP	CD2-CE2	5.49	1.48	1.41
1	C	351	TRP	CD2-CE2	5.30	1.47	1.41
1	D	483	TRP	CD2-CE2	5.28	1.47	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	400	LEU	CA-CB-CG	6.94	131.27	115.30
1	A	400	LEU	CA-CB-CG	5.32	127.54	115.30
1	C	400	LEU	CB-CG-CD2	-5.10	102.33	111.00
1	A	400	LEU	CB-CG-CD2	-5.08	102.36	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3475	0	3509	33	0
1	B	3471	0	3505	44	0
1	C	3452	0	3487	25	0
1	D	3371	0	3418	36	0
2	A	88	0	52	0	0
2	B	88	0	52	2	0
2	C	88	0	52	0	0
2	D	88	0	52	1	0
3	A	53	0	31	4	0
3	B	53	0	31	5	0
3	C	53	0	31	2	0
3	D	53	0	31	5	0
4	D	5	0	0	0	0
5	A	9	0	0	1	0
5	B	8	0	0	0	0
5	C	7	0	0	1	0
All	All	14362	0	14251	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:589:ARG:HH11	1:B:589:ARG:HG3	1.48	0.78
1:D:584:ARG:HB3	1:D:587:ILE:HD12	1.63	0.78
1:A:445:ILE:HG12	5:A:2008:HOH:O	1.84	0.77
1:A:610:ILE:HA	1:A:611:HIS:C	2.04	0.77
1:C:564:VAL:HG12	1:C:610:ILE:HD11	1.68	0.75

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/511 (87%)	417 (94%)	29 (6%)	0	100	100
1	B	446/511 (87%)	421 (94%)	22 (5%)	3 (1%)	26	61
1	C	442/511 (86%)	416 (94%)	25 (6%)	1 (0%)	52	83
1	D	431/511 (84%)	396 (92%)	34 (8%)	1 (0%)	52	83
All	All	1765/2044 (86%)	1650 (94%)	110 (6%)	5 (0%)	46	78

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	197	ASN
1	B	275	ALA
1	C	513	ALA
1	D	273	ALA
1	B	285	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/419 (88%)	347 (95%)	20 (5%)	27	60
1	B	366/419 (87%)	348 (95%)	18 (5%)	31	65
1	C	365/419 (87%)	343 (94%)	22 (6%)	24	55
1	D	357/419 (85%)	331 (93%)	26 (7%)	17	43
All	All	1455/1676 (87%)	1369 (94%)	86 (6%)	24	55

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	127	LYS
1	C	272	ARG
1	D	433	ILE
1	C	187	VAL
1	C	201	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	432	ASN
1	B	454	HIS
1	D	222	ASN
1	B	250	GLN
1	B	403	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FAD	A	1612	-	48,58,58	2.16	14 (29%)	54,89,89	2.67	14 (25%)
2	NAD	A	700	-	38,48,48	2.08	8 (21%)	47,73,73	2.60	5 (10%)
2	NAD	A	701	-	38,48,48	1.82	6 (15%)	47,73,73	2.79	8 (17%)
3	FAD	B	1612	-	48,58,58	2.18	11 (22%)	54,89,89	2.67	12 (22%)
2	NAD	B	700	-	38,48,48	1.68	7 (18%)	47,73,73	2.69	8 (17%)
2	NAD	B	701	-	38,48,48	1.92	7 (18%)	47,73,73	2.34	5 (10%)
3	FAD	C	1612	-	48,58,58	2.06	12 (25%)	54,89,89	2.40	9 (16%)
2	NAD	C	700	-	38,48,48	1.84	6 (15%)	47,73,73	2.67	8 (17%)
2	NAD	C	701	-	38,48,48	2.03	7 (18%)	47,73,73	2.28	3 (6%)
3	FAD	D	1612	-	48,58,58	2.14	9 (18%)	54,89,89	2.47	15 (27%)
4	SO4	D	1613	-	4,4,4	0.63	0	6,6,6	0.16	0
2	NAD	D	700	-	38,48,48	1.83	6 (15%)	47,73,73	2.55	6 (12%)
2	NAD	D	701	-	38,48,48	1.85	7 (18%)	47,73,73	2.95	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	1612	-	-	0/30/50/50	0/6/6/6
2	NAD	A	700	-	-	0/22/62/62	0/5/5/5
2	NAD	A	701	-	-	0/22/62/62	0/5/5/5
3	FAD	B	1612	-	-	0/30/50/50	0/6/6/6
2	NAD	B	700	-	-	0/22/62/62	0/5/5/5
2	NAD	B	701	-	-	0/22/62/62	0/5/5/5
3	FAD	C	1612	-	-	0/30/50/50	0/6/6/6
2	NAD	C	700	-	-	0/22/62/62	0/5/5/5
2	NAD	C	701	-	-	0/22/62/62	0/5/5/5
3	FAD	D	1612	-	-	0/30/50/50	0/6/6/6
4	SO4	D	1613	-	-	0/0/0/0	0/0/0/0
2	NAD	D	700	-	-	0/22/62/62	0/5/5/5
2	NAD	D	701	-	-	0/22/62/62	0/5/5/5

The worst 5 of 100 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	700	NAD	C3N-C7N	-7.18	1.39	1.50
2	C	701	NAD	C3N-C7N	-7.17	1.39	1.50
2	D	700	NAD	C3N-C7N	-6.18	1.40	1.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	700	NAD	C3N-C7N	-6.15	1.40	1.50
2	B	701	NAD	C3N-C7N	-6.09	1.40	1.50

The worst 5 of 103 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	700	NAD	N3A-C2A-N1A	-15.13	117.31	128.89
2	A	701	NAD	N3A-C2A-N1A	-15.04	117.38	128.89
2	D	700	NAD	N3A-C2A-N1A	-14.70	117.64	128.89
2	B	700	NAD	N3A-C2A-N1A	-14.60	117.72	128.89
2	D	701	NAD	N3A-C2A-N1A	-14.51	117.78	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1612	FAD	4	0
3	B	1612	FAD	5	0
2	B	700	NAD	1	0
2	B	701	NAD	1	0
3	C	1612	FAD	2	0
3	D	1612	FAD	5	0
2	D	700	NAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	450/511 (88%)	0.12	2 (0%) 93 92	27, 42, 74, 111	0
1	B	450/511 (88%)	0.19	2 (0%) 93 92	30, 51, 81, 116	0
1	C	446/511 (87%)	0.17	1 (0%) 95 95	30, 52, 84, 100	0
1	D	435/511 (85%)	0.40	10 (2%) 64 60	39, 67, 104, 123	0
All	All	1781/2044 (87%)	0.22	15 (0%) 87 86	27, 53, 91, 123	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	179	LEU	2.7
1	A	128	ALA	2.6
1	D	325	ALA	2.5
1	A	611	HIS	2.4
1	D	326	LEU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAD	A	701	44/44	0.93	0.25	0.92	71,87,104,112	0
2	NAD	B	701	44/44	0.92	0.28	0.77	74,102,110,116	0
2	NAD	C	701	44/44	0.96	0.22	0.55	61,70,81,86	0
2	NAD	D	701	44/44	0.94	0.25	0.09	71,97,111,112	0
2	NAD	C	700	44/44	0.98	0.20	-0.52	30,41,56,60	0
4	SO4	D	1613	5/5	0.90	0.17	-0.55	72,74,86,89	0
3	FAD	C	1612	53/53	0.98	0.20	-0.55	30,39,50,54	0
3	FAD	B	1612	53/53	0.98	0.20	-0.56	29,43,50,52	0
3	FAD	A	1612	53/53	0.98	0.20	-0.62	22,33,40,43	0
2	NAD	A	700	44/44	0.98	0.19	-0.67	23,33,47,47	0
3	FAD	D	1612	53/53	0.97	0.21	-0.81	38,53,63,67	0
2	NAD	B	700	44/44	0.98	0.19	-0.82	27,41,49,49	0
2	NAD	D	700	44/44	0.96	0.19	-0.83	36,57,79,82	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.