



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:14 PM GMT

PDB ID : 4BWV  
Title : Structure of Neurospora crassa PAN3 pseudokinase mutant  
Authors : Christie, M.; Boland, A.; Huntzinger, E.; Weichenrieder, O.; Izaurralde, E.  
Deposited on : 2013-07-04  
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2228 (2.90-2.82)
Clashscore	102246	2499 (2.90-2.82)
Ramachandran outliers	100387	2439 (2.90-2.82)
Sidechain outliers	100360	2442 (2.90-2.82)
RSRZ outliers	91569	2236 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	<div> <div>3%</div> <div>82%</div> <div>10%</div> <div>8%</div> </div>
1	B	429	<div> <div>3%</div> <div>82%</div> <div>11%</div> <div>7%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	0	0
			3224	2055	573	583	13			
1	B	397	Total	C	N	O	S	0	0	0
			3236	2058	578	587	13			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	228	GLY	-	EXPRESSION TAG	UNP Q7SDP4
A	229	PRO	-	EXPRESSION TAG	UNP Q7SDP4
A	230	HIS	-	EXPRESSION TAG	UNP Q7SDP4
A	231	MET	-	EXPRESSION TAG	UNP Q7SDP4
A	232	LEU	-	EXPRESSION TAG	UNP Q7SDP4
A	233	GLU	-	EXPRESSION TAG	UNP Q7SDP4
A	510	MET	LEU	ENGINEERED MUTATION	UNP Q7SDP4
A	511	PRO	GLY	ENGINEERED MUTATION	UNP Q7SDP4
A	512	MET	GLY	ENGINEERED MUTATION	UNP Q7SDP4
A	514	GLY	THR	ENGINEERED MUTATION	UNP Q7SDP4
A	515	ALA	THR	ENGINEERED MUTATION	UNP Q7SDP4
A	516	ARG	HIS	ENGINEERED MUTATION	UNP Q7SDP4
A	517	PHE	LEU	ENGINEERED MUTATION	UNP Q7SDP4
B	228	GLY	-	EXPRESSION TAG	UNP Q7SDP4
B	229	PRO	-	EXPRESSION TAG	UNP Q7SDP4
B	230	HIS	-	EXPRESSION TAG	UNP Q7SDP4
B	231	MET	-	EXPRESSION TAG	UNP Q7SDP4
B	232	LEU	-	EXPRESSION TAG	UNP Q7SDP4
B	233	GLU	-	EXPRESSION TAG	UNP Q7SDP4
B	510	MET	LEU	ENGINEERED MUTATION	UNP Q7SDP4
B	511	PRO	GLY	ENGINEERED MUTATION	UNP Q7SDP4
B	512	MET	GLY	ENGINEERED MUTATION	UNP Q7SDP4
B	514	GLY	THR	ENGINEERED MUTATION	UNP Q7SDP4
B	515	ALA	THR	ENGINEERED MUTATION	UNP Q7SDP4

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	516	ARG	HIS	ENGINEERED MUTATION	UNP Q7SDP4
B	517	PHE	LEU	ENGINEERED MUTATION	UNP Q7SDP4

- Molecule 2 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (three-letter code: AGS) (formula:  $C_{10}H_{16}N_5O_{12}P_3S$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN-3



- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN-3



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.31Å 90.31Å 230.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.16 – 2.85 45.16 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (45.16-2.85) 99.9 (45.16-2.85)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.62 (at 2.86Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.211 , 0.242 0.217 , 0.245	Depositor DCC
$R_{free}$ test set	1256 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.4	Xtriage
Anisotropy	0.652	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 51.4	EDS
Estimated twinning fraction	0.066 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 24755 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6524	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AGS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/3295	0.41	0/4457
1	B	0.22	0/3306	0.41	0/4471
All	All	0.22	0/6601	0.41	0/8928

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3224	0	3215	29	0
1	B	3236	0	3224	30	0
2	A	31	0	12	2	0
2	B	31	0	12	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
All	All	6524	0	6463	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:543:ARG:HH21	1:B:246:LEU:HD13	1.48	0.79
1:A:510:MET:HG2	1:B:515:ALA:HA	1.81	0.62
1:B:413:LYS:NZ	2:B:1647:AGS:O1A	2.31	0.61
1:B:543:ARG:NH1	1:B:607:ASP:OD2	2.34	0.61
1:A:462:THR:HG21	1:A:475:ALA:HB1	1.82	0.61
1:B:277:ASN:OD1	1:B:279:THR:OG1	2.19	0.59
1:A:511:PRO:HB3	1:B:511:PRO:HB3	1.85	0.56
1:B:538:GLU:O	1:B:541:ASN:ND2	2.37	0.56
1:A:568:GLU:HG3	1:A:569:THR:HG23	1.89	0.55
1:B:458:LEU:HD23	1:B:467:LEU:HD21	1.89	0.55
1:B:618:SER:OG	1:B:620:ASP:O	2.22	0.54
1:B:462:THR:HG21	1:B:475:ALA:HB1	1.89	0.54
1:A:620:ASP:O	1:A:622:ALA:N	2.41	0.54
1:A:359:THR:HG23	2:A:1647:AGS:H3'	1.89	0.54
1:A:357:SER:O	2:A:1647:AGS:O2'	2.26	0.54
1:A:382:GLU:HG2	1:A:386:TRP:CD1	2.43	0.53
1:B:568:GLU:HG3	1:B:569:THR:HG23	1.91	0.53
1:A:234:ILE:N	1:A:537:ARG:HH11	2.06	0.53
1:B:407:ARG:NH1	1:B:430:ASP:OD2	2.42	0.53
1:A:543:ARG:NH1	1:A:607:ASP:OD2	2.43	0.52
1:B:273:LYS:HE2	1:B:275:THR:HB	1.91	0.52
1:A:286:TRP:HB2	1:A:303:ARG:HB3	1.93	0.51
1:A:515:ALA:HA	1:B:510:MET:HE2	1.91	0.50
1:B:334:HIS:HB2	1:B:350:VAL:HG13	1.94	0.49
1:A:536:MET:HB3	1:B:246:LEU:HD21	1.96	0.48
1:A:615:LEU:HD21	1:A:626:VAL:HG22	1.96	0.48
1:A:235:PRO:HD3	1:A:534:HIS:CE1	2.49	0.47
1:B:311:ASN:ND2	1:B:313:LYS:HB2	2.29	0.47
1:B:548:MET:SD	1:B:603:LEU:HD21	2.55	0.47
1:A:244:LYS:HB3	1:A:356:LEU:HD11	1.97	0.46
1:A:566:TRP:O	1:A:572:ARG:NE	2.41	0.46
1:B:408:CYS:SG	1:B:413:LYS:HG3	2.57	0.45
1:B:472:ILE:O	1:B:476:LEU:HB2	2.17	0.44
1:A:571:GLU:OE2	1:A:638:ARG:NH1	2.50	0.43
1:A:323:LYS:HG2	1:A:333:VAL:O	2.19	0.43
1:A:568:GLU:O	1:A:573:LEU:HD13	2.19	0.43
1:A:553:VAL:HG22	1:B:583:PHE:HB3	2.01	0.42
1:B:379:LYS:HD3	1:B:482:ARG:HD2	2.00	0.42
1:B:286:TRP:HB2	1:B:303:ARG:HB3	2.00	0.42
1:B:505:THR:O	1:B:508:THR:OG1	2.34	0.41
1:A:382:GLU:OE2	1:A:484:SER:HB3	2.21	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:273:LYS:HG2	1:B:275:THR:H	1.86	0.41
1:A:526:GLN:HG3	1:B:328:ALA:HB2	2.02	0.41
1:A:587:ASP:HA	1:A:593:ARG:HD2	2.03	0.41
1:A:615:LEU:HD23	1:A:615:LEU:HA	1.86	0.41
1:A:246:LEU:HD21	1:B:536:MET:HB3	2.03	0.41
1:B:462:THR:HG22	1:B:478:THR:OG1	2.21	0.41
1:A:548:MET:HE2	1:B:548:MET:HB3	2.02	0.40
1:A:499:ALA:HA	1:A:502:GLU:O	2.20	0.40
1:B:380:ILE:HD12	1:B:460:LEU:HB3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/429 (91%)	376 (97%)	12 (3%)	1 (0%)	46	76
1	B	391/429 (91%)	376 (96%)	15 (4%)	0	100	100
All	All	780/858 (91%)	752 (96%)	27 (4%)	1 (0%)	56	85

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	621	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	351/380 (92%)	349 (99%)	2 (1%)	90	97
1	B	353/380 (93%)	349 (99%)	4 (1%)	80	94
All	All	704/760 (93%)	698 (99%)	6 (1%)	84	95

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	453	PHE
1	A	517	PHE
1	B	279	THR
1	B	453	PHE
1	B	481	PRO
1	B	517	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	534	HIS
1	A	555	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	AGS	A	1647	3	24,33,33	1.91	6 (25%)	28,52,52	3.63	15 (53%)
2	AGS	B	1647	3	24,33,33	1.91	6 (25%)	28,52,52	3.60	15 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AGS	A	1647	3	-	0/15/38/38	0/3/3/3
2	AGS	B	1647	3	-	0/15/38/38	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1647	AGS	O4'-C4'	-2.31	1.39	1.45
2	B	1647	AGS	O4'-C4'	-2.25	1.39	1.45
2	A	1647	AGS	O3'-C3'	-2.12	1.37	1.43
2	B	1647	AGS	O3'-C3'	-2.09	1.38	1.43
2	A	1647	AGS	C2'-C3'	-2.04	1.47	1.53
2	B	1647	AGS	C2'-C3'	-2.00	1.47	1.53
2	A	1647	AGS	C2-N3	2.59	1.36	1.32
2	B	1647	AGS	C2-N3	2.61	1.36	1.32
2	B	1647	AGS	C6-N6	2.89	1.43	1.34
2	A	1647	AGS	C6-N6	2.92	1.44	1.34
2	A	1647	AGS	PG-S1G	6.51	2.03	1.90
2	B	1647	AGS	PG-S1G	6.56	2.03	1.90

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1647	AGS	N3-C2-N1	-11.39	120.17	128.89
2	B	1647	AGS	N3-C2-N1	-11.31	120.23	128.89
2	A	1647	AGS	O2A-PA-O1A	-5.60	82.18	112.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1647	AGS	O2A-PA-O1A	-5.59	82.20	112.53
2	B	1647	AGS	O2B-PB-O1B	-5.43	83.09	112.53
2	A	1647	AGS	O2B-PB-O1B	-5.22	84.23	112.53
2	A	1647	AGS	PB-O3B-PG	-5.05	115.73	132.67
2	A	1647	AGS	O2A-PA-O5'	-4.93	83.58	108.46
2	B	1647	AGS	O2A-PA-O5'	-4.90	83.74	108.46
2	B	1647	AGS	O2A-PA-O3A	-4.49	84.73	105.09
2	A	1647	AGS	O2A-PA-O3A	-4.39	85.18	105.09
2	B	1647	AGS	PB-O3B-PG	-4.18	118.66	132.67
2	A	1647	AGS	PA-O3A-PB	-4.17	121.01	132.73
2	B	1647	AGS	PA-O3A-PB	-4.17	121.03	132.73
2	B	1647	AGS	C4-C5-N7	-2.58	107.10	109.48
2	A	1647	AGS	C4-C5-N7	-2.53	107.15	109.48
2	A	1647	AGS	O5'-PA-O1A	2.04	117.53	109.62
2	B	1647	AGS	O5'-PA-O1A	2.07	117.64	109.62
2	B	1647	AGS	O4'-C4'-C3'	2.12	109.42	105.15
2	A	1647	AGS	O4'-C4'-C3'	2.28	109.75	105.15
2	B	1647	AGS	O2G-PG-O3B	2.70	117.35	105.09
2	A	1647	AGS	O3G-PG-O3B	2.81	117.83	105.09
2	A	1647	AGS	O2B-PB-O3B	2.86	118.05	105.09
2	B	1647	AGS	O2B-PB-O3A	2.94	118.44	105.09
2	A	1647	AGS	O2G-PG-O3B	2.99	118.67	105.09
2	A	1647	AGS	O2B-PB-O3A	3.04	118.90	105.09
2	B	1647	AGS	O2B-PB-O3B	3.20	119.61	105.09
2	B	1647	AGS	O3G-PG-O3B	3.25	119.85	105.09
2	B	1647	AGS	O3A-PA-O5'	5.73	118.13	102.94
2	A	1647	AGS	O3A-PA-O5'	6.05	118.98	102.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1647	AGS	2	0
2	B	1647	AGS	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	395/429 (92%)	0.12	14 (3%) 48 40	42, 83, 149, 207	0
1	B	397/429 (92%)	0.14	11 (2%) 56 51	54, 82, 139, 186	0
All	All	792/858 (92%)	0.13	25 (3%) 51 44	42, 83, 147, 207	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	471	ASN	4.7
1	B	469	PHE	3.8
1	A	279	THR	3.7
1	B	569	THR	3.5
1	B	619	ARG	3.5
1	A	470	ASN	3.5
1	B	501	GLY	3.3
1	B	568	GLU	3.1
1	B	437	PRO	2.9
1	A	236	LYS	2.8
1	A	568	GLU	2.7
1	B	256	SER	2.7
1	A	469	PHE	2.6
1	A	588	ALA	2.6
1	A	278	SER	2.6
1	A	434	HIS	2.6
1	B	617	THR	2.5
1	A	619	ARG	2.5
1	A	566	TRP	2.3
1	A	646	GLU	2.3
1	A	569	THR	2.3
1	B	503	THR	2.2
1	A	499	ALA	2.2
1	B	472	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	500	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	AGS	A	1647	31/31	0.95	0.17	-0.50	48,88,214,296	0
2	AGS	B	1647	31/31	0.95	0.18	-0.56	44,74,219,275	0
3	MG	B	1648	1/1	0.97	0.14	-0.99	69,69,69,69	0
3	MG	A	1648	1/1	0.97	0.13	-2.35	104,104,104,104	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.