



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:43 AM GMT

PDB ID : 3BYV
Title : Crystal structure of Toxoplasma gondii specific rhoptry antigen kinase domain
Authors : Wernimont, A.K.; Lunin, V.V.; Yang, C.; Lew, J.; Kozieradzki, I.; Lin, Y.H.; Sun, X.; Khuu, C.; Zhao, Y.; Schapira, M.; Arrowsmith, C.H.; Edwards, A.M.; Weigelt, J.; Bochkarev, A.; Hui, R.; Sibley, D.; Qiu, W.; Structural Genomics Consortium (SGC)
Deposited on : 2008-01-16
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	<div> <div>21%</div> <div>68%</div> <div>20%</div> <div>• 8%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3023 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhoptyr kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	8	0
			2846	1839	498	500	9			


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	192	MET	-	EXPRESSION TAG	PDB 3BYV
A	193	HIS	-	EXPRESSION TAG	PDB 3BYV
A	194	HIS	-	EXPRESSION TAG	PDB 3BYV
A	195	HIS	-	EXPRESSION TAG	PDB 3BYV
A	196	HIS	-	EXPRESSION TAG	PDB 3BYV
A	197	HIS	-	EXPRESSION TAG	PDB 3BYV
A	198	HIS	-	EXPRESSION TAG	PDB 3BYV
A	199	SER	-	EXPRESSION TAG	PDB 3BYV
A	200	SER	-	EXPRESSION TAG	PDB 3BYV
A	201	GLY	-	EXPRESSION TAG	PDB 3BYV
A	202	ARG	-	EXPRESSION TAG	PDB 3BYV
A	203	GLU	-	EXPRESSION TAG	PDB 3BYV
A	204	ASN	-	EXPRESSION TAG	PDB 3BYV
A	205	LEU	-	EXPRESSION TAG	PDB 3BYV
A	206	TYR	-	EXPRESSION TAG	PDB 3BYV
A	207	PHE	-	EXPRESSION TAG	PDB 3BYV
A	208	GLN	-	EXPRESSION TAG	PDB 3BYV
A	209	GLY	-	EXPRESSION TAG	PDB 3BYV

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	164	Total	O	0	0
			164	164		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rhoptry kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.07Å 69.82Å 85.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.44 – 1.80 36.43 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.44-1.80) 99.9 (36.43-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.202 , 0.233 0.245 , 0.274	Depositor DCC
R_{free} test set	1692 reflections (5.08%)	DCC
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 48.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 33423 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3023	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.33	10/2935 (0.3%)	1.25	26/3976 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	354	VAL	C-O	10.18	1.42	1.23
1	A	412	SER	CB-OG	-6.94	1.33	1.42
1	A	280	TYR	CD1-CE1	6.37	1.49	1.39
1	A	422	TYR	CE1-CZ	5.82	1.46	1.38
1	A	422	TYR	CG-CD2	5.67	1.46	1.39
1	A	467[A]	ARG	CG-CD	5.61	1.66	1.51
1	A	467[B]	ARG	CG-CD	5.61	1.66	1.51
1	A	419	VAL	CB-CG2	5.47	1.64	1.52
1	A	441	PHE	CD2-CE2	5.42	1.50	1.39
1	A	534	PHE	CD1-CE1	5.02	1.49	1.39

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	477	ARG	NE-CZ-NH1	12.00	126.30	120.30
1	A	260	GLY	N-CA-C	-10.35	87.23	113.10
1	A	477	ARG	NE-CZ-NH2	-8.92	115.84	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	476	ASP	CB-CG-OD1	8.31	125.78	118.30
1	A	345	ASP	CB-CG-OD2	7.77	125.29	118.30
1	A	212	ASP	CB-CG-OD2	-7.27	111.76	118.30
1	A	281	LEU	CA-CB-CG	7.15	131.75	115.30
1	A	502	LEU	CB-CG-CD2	6.44	121.94	111.00
1	A	486	ASP	CB-CG-OD2	-6.38	112.56	118.30
1	A	345	ASP	CB-CG-OD1	-6.15	112.76	118.30
1	A	480	LEU	CB-CG-CD2	-6.15	100.54	111.00
1	A	539	LYS	CD-CE-NZ	-6.13	97.59	111.70
1	A	558	ARG	NE-CZ-NH2	-6.03	117.28	120.30
1	A	492	LEU	CB-CG-CD1	-5.84	101.07	111.00
1	A	476	ASP	CB-CG-OD2	-5.79	113.09	118.30
1	A	501	ASP	CB-CG-OD1	5.68	123.41	118.30
1	A	475	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	A	558	ARG	CG-CD-NE	5.61	123.57	111.80
1	A	322	ARG	NE-CZ-NH1	5.55	123.08	120.30
1	A	502	LEU	CA-CB-CG	5.49	127.92	115.30
1	A	459	PHE	CB-CG-CD1	-5.44	116.99	120.80
1	A	486	ASP	CB-CG-OD1	5.36	123.13	118.30
1	A	218	LEU	CA-CB-CG	-5.31	103.09	115.30
1	A	335	ARG	CG-CD-NE	-5.09	101.10	111.80
1	A	402	LEU	CB-CG-CD2	-5.07	102.38	111.00
1	A	383	VAL	CG1-CB-CG2	-5.04	102.84	110.90

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	259	LEU	Peptide
1	A	292	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2846	0	2861	53	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	12	0	18	1	0
4	A	164	0	0	6	2
All	All	3023	0	2879	53	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:390:THR:HG22	1:A:391:HIS:CD2	1.65	1.32
1:A:467[A]:ARG:HD3	4:A:836:HOH:O	1.30	1.28
1:A:390:THR:CG2	1:A:391:HIS:CD2	2.28	1.17
1:A:390:THR:HG22	1:A:391:HIS:HD2	0.92	1.07
1:A:303:PRO:HD2	1:A:362[A]:TRP:HE3	1.22	1.03
1:A:303:PRO:HD2	1:A:362[A]:TRP:CE3	1.94	1.02
1:A:433:ARG:HG2	4:A:829:HOH:O	1.61	0.99
1:A:217:GLU:OE2	1:A:221:ARG:HD2	1.62	0.97
1:A:390:THR:CG2	1:A:391:HIS:HD2	1.71	0.93
1:A:303:PRO:CD	1:A:362[A]:TRP:CE3	2.57	0.87
1:A:558:ARG:NH2	4:A:810:HOH:O	2.08	0.86
1:A:303:PRO:CD	1:A:362[A]:TRP:HE3	1.95	0.79
1:A:231:SER:CB	4:A:858:HOH:O	2.30	0.79
1:A:312:LYS:HD2	1:A:342:LEU:HD23	1.67	0.77
1:A:312:LYS:HD2	1:A:342:LEU:CD2	2.16	0.76
1:A:404:LEU:HD12	1:A:561:LEU:HD23	1.71	0.73
1:A:306:ASN:O	1:A:310:GLN:HG2	1.91	0.70
1:A:301:GLU:HA	1:A:362[B]:TRP:CZ3	2.32	0.64
1:A:390:THR:HG23	1:A:391:HIS:CD2	2.27	0.64
1:A:408:ARG:HH21	1:A:558:ARG:HH12	1.48	0.60
1:A:217:GLU:OE2	1:A:221:ARG:CD	2.46	0.59
1:A:309:LYS:HE2	1:A:313:GLU:OE1	2.04	0.58
1:A:247:TRP:HZ2	1:A:367:PHE:CZ	2.22	0.58
1:A:264:ARG:NH2	1:A:285:ASP:OD2	2.37	0.57
1:A:404:LEU:CD1	1:A:561:LEU:HD23	2.35	0.56
1:A:537[B]:TYR:HB3	1:A:538:PRO:HD3	1.87	0.56
1:A:247:TRP:CZ2	1:A:367:PHE:CZ	2.93	0.55
1:A:303:PRO:HD3	1:A:362[A]:TRP:CE3	2.41	0.55
1:A:238:ALA:HB2	1:A:274:GLN:HE21	1.75	0.52
1:A:401:GLN:OE1	1:A:561:LEU:HD21	2.10	0.51
1:A:315:VAL:HG22	1:A:339:PRO:HG2	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:ARG:HG2	1:A:281:LEU:HD23	1.93	0.50
1:A:245:SER:HB3	1:A:246:LEU:HG	1.93	0.49
1:A:238:ALA:HB2	1:A:274:GLN:NE2	2.27	0.48
1:A:241:LEU:O	1:A:245:SER:HB2	2.14	0.48
1:A:353:ARG:HG3	1:A:362[A]:TRP:NE1	2.28	0.48
1:A:525:GLN:HG3	1:A:526:PRO:HD3	1.96	0.48
1:A:562[B]:SER:HA	1:A:565:LEU:HD22	1.96	0.47
1:A:562[A]:SER:HA	1:A:565:LEU:HD22	1.97	0.47
1:A:217:GLU:HG3	1:A:221:ARG:HD3	1.98	0.46
1:A:495:TYR:CD2	1:A:503:PRO:HD3	2.51	0.45
1:A:500:ALA:O	1:A:501:ASP:HB2	2.15	0.45
1:A:518:ARG:NH2	4:A:844:HOH:O	2.49	0.45
1:A:467[A]:ARG:NH1	4:A:804:HOH:O	2.51	0.44
1:A:536:ARG:HG2	1:A:538:PRO:HD2	1.98	0.44
1:A:467[A]:ARG:HG3	1:A:467[A]:ARG:NH1	2.33	0.43
1:A:467[A]:ARG:NH1	1:A:467[A]:ARG:CG	2.80	0.43
1:A:278:TYR:OH	3:A:604:EDO:H22	2.18	0.43
1:A:240:SER:O	1:A:244:THR:HG23	2.19	0.42
1:A:232:GLU:O	1:A:234:MET:N	2.52	0.42
1:A:385:LEU:HD22	1:A:500:ALA:HB3	2.02	0.41
1:A:297:PRO:HG2	1:A:364:LEU:HB2	2.03	0.41
1:A:239:ASP:HA	1:A:271:VAL:HG21	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:740:HOH:O	4:A:855:HOH:O[4_455]	1.99	0.21
4:A:819:HOH:O	4:A:855:HOH:O[4_455]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/377 (90%)	330 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/336 (92%)	294 (95%)	16 (5%)	29	12

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	218	LEU
1	A	236	SER
1	A	245	SER
1	A	247	TRP
1	A	259	LEU
1	A	265	THR
1	A	270	THR
1	A	281	LEU
1	A	312	LYS
1	A	367	PHE
1	A	390	THR
1	A	454	SER
1	A	465	GLU
1	A	504	ILE
1	A	525	GLN
1	A	565	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	274	GLN
1	A	391	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	602	-	3,3,3	0.78	0	2,2,2	0.28	0
3	EDO	A	603	-	3,3,3	0.68	0	2,2,2	0.36	0
3	EDO	A	604	-	3,3,3	0.64	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	602	-	-	0/1/1/1	0/0/0/0
3	EDO	A	603	-	-	0/1/1/1	0/0/0/0
3	EDO	A	604	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	604	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/377 (91%)	1.11	79 (22%) ⓘ ⓘ	2, 12, 23, 33	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	362[A]	TRP	7.1
1	A	248	ASN	6.7
1	A	236	SER	6.3
1	A	244	THR	6.1
1	A	240	SER	6.0
1	A	430[A]	LEU	5.4
1	A	346	PRO	5.0
1	A	301	GLU	4.6
1	A	300	THR	4.6
1	A	361	MET	4.3
1	A	429	VAL	4.2
1	A	354	VAL	4.1
1	A	305	SER	4.1
1	A	377	LEU	4.0
1	A	428	ILE	4.0
1	A	265	THR	3.9
1	A	410	LEU	3.9
1	A	260	GLY	3.8
1	A	235	GLN	3.7
1	A	268	ARG	3.7
1	A	287	GLU	3.6
1	A	402	LEU	3.6
1	A	237	ALA	3.5
1	A	336	PHE	3.5
1	A	493	VAL	3.4
1	A	353	ARG	3.4
1	A	245	SER	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	406	VAL	3.4
1	A	438	LEU	3.4
1	A	251	GLN	3.3
1	A	243	SER	3.2
1	A	389	SER	3.2
1	A	436	VAL	3.2
1	A	407	ILE	3.2
1	A	423	LEU	3.1
1	A	490	LEU	3.1
1	A	519	SER	3.1
1	A	239	ASP	3.1
1	A	347	GLN	3.0
1	A	250	GLY	3.0
1	A	380	PHE	3.0
1	A	288	THR	2.9
1	A	298	TYR	2.9
1	A	269	GLY	2.9
1	A	518	ARG	2.9
1	A	337	ILE	2.8
1	A	426	VAL	2.8
1	A	345	ASP	2.8
1	A	284	THR	2.7
1	A	309	LYS	2.6
1	A	511	GLY	2.6
1	A	334	LEU	2.6
1	A	467[A]	ARG	2.6
1	A	520	CYS	2.5
1	A	215[A]	ILE	2.5
1	A	306	ASN	2.5
1	A	409	LEU	2.5
1	A	437	PHE	2.5
1	A	420	HIS	2.5
1	A	513	SER	2.5
1	A	413	LEU	2.5
1	A	304	PRO	2.5
1	A	238	ALA	2.4
1	A	521	LYS	2.4
1	A	267	VAL	2.4
1	A	439	THR	2.3
1	A	441	PHE	2.3
1	A	403	THR	2.3
1	A	421	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	352	ILE	2.3
1	A	261	GLU	2.3
1	A	234	MET	2.3
1	A	254	ARG	2.2
1	A	537[A]	TYR	2.1
1	A	259	LEU	2.1
1	A	351	MET	2.1
1	A	252	PRO	2.1
1	A	271	VAL	2.1
1	A	489	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	EDO	A	602	4/4	0.93	0.18	0.30	5,6,7,8	0
3	EDO	A	604	4/4	0.78	0.14	-0.19	32,34,35,37	0
3	EDO	A	603	4/4	0.90	0.12	-0.28	15,17,25,27	0
2	MG	A	601	1/1	0.97	0.09	-	8,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.