



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2016 – 11:53 AM EDT

PDB ID : 5BY7  
Title : AbyA1 - tetronic acid condensing enzyme  
Authors : Byrne, M.J.; Race, P.R.  
Deposited on : 2015-06-10  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	<div> <div>3%</div> <div>84% 10% 6%</div> </div>
1	B	360	<div> <div>3%</div> <div>82% 7% 11%</div> </div>
1	C	360	<div> <div>3%</div> <div>80% 6% 14%</div> </div>
1	D	360	<div> <div>3%</div> <div>79% . . 16%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9602 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-ACP synthase III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	337	Total	C	N	O	S	0	0	0
			2441	1536	439	461	5			
1	B	321	Total	C	N	O	S	0	0	0
			2341	1470	418	448	5			
1	C	310	Total	C	N	O	S	0	0	0
			2211	1386	392	429	4			
1	D	302	Total	C	N	O	S	0	0	0
			2129	1338	376	411	4			

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-37	MET	-	initiating methionine	UNP F4F7F9
A	-36	ALA	-	expression tag	UNP F4F7F9
A	-35	HIS	-	expression tag	UNP F4F7F9
A	-34	HIS	-	expression tag	UNP F4F7F9
A	-33	HIS	-	expression tag	UNP F4F7F9
A	-32	HIS	-	expression tag	UNP F4F7F9
A	-31	HIS	-	expression tag	UNP F4F7F9
A	-30	HIS	-	expression tag	UNP F4F7F9
A	-29	SER	-	expression tag	UNP F4F7F9
A	-28	SER	-	expression tag	UNP F4F7F9
A	-27	GLY	-	expression tag	UNP F4F7F9
A	-26	LEU	-	expression tag	UNP F4F7F9
A	-25	GLU	-	expression tag	UNP F4F7F9
A	-24	VAL	-	expression tag	UNP F4F7F9
A	-23	LEU	-	expression tag	UNP F4F7F9
A	-22	PHE	-	expression tag	UNP F4F7F9
A	-21	GLN	-	expression tag	UNP F4F7F9
A	-20	GLY	-	expression tag	UNP F4F7F9
A	-19	PRO	-	expression tag	UNP F4F7F9
B	-37	MET	-	initiating methionine	UNP F4F7F9
B	-36	ALA	-	expression tag	UNP F4F7F9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-35	HIS	-	expression tag	UNP F4F7F9
B	-34	HIS	-	expression tag	UNP F4F7F9
B	-33	HIS	-	expression tag	UNP F4F7F9
B	-32	HIS	-	expression tag	UNP F4F7F9
B	-31	HIS	-	expression tag	UNP F4F7F9
B	-30	HIS	-	expression tag	UNP F4F7F9
B	-29	SER	-	expression tag	UNP F4F7F9
B	-28	SER	-	expression tag	UNP F4F7F9
B	-27	GLY	-	expression tag	UNP F4F7F9
B	-26	LEU	-	expression tag	UNP F4F7F9
B	-25	GLU	-	expression tag	UNP F4F7F9
B	-24	VAL	-	expression tag	UNP F4F7F9
B	-23	LEU	-	expression tag	UNP F4F7F9
B	-22	PHE	-	expression tag	UNP F4F7F9
B	-21	GLN	-	expression tag	UNP F4F7F9
B	-20	GLY	-	expression tag	UNP F4F7F9
B	-19	PRO	-	expression tag	UNP F4F7F9
C	-37	MET	-	initiating methionine	UNP F4F7F9
C	-36	ALA	-	expression tag	UNP F4F7F9
C	-35	HIS	-	expression tag	UNP F4F7F9
C	-34	HIS	-	expression tag	UNP F4F7F9
C	-33	HIS	-	expression tag	UNP F4F7F9
C	-32	HIS	-	expression tag	UNP F4F7F9
C	-31	HIS	-	expression tag	UNP F4F7F9
C	-30	HIS	-	expression tag	UNP F4F7F9
C	-29	SER	-	expression tag	UNP F4F7F9
C	-28	SER	-	expression tag	UNP F4F7F9
C	-27	GLY	-	expression tag	UNP F4F7F9
C	-26	LEU	-	expression tag	UNP F4F7F9
C	-25	GLU	-	expression tag	UNP F4F7F9
C	-24	VAL	-	expression tag	UNP F4F7F9
C	-23	LEU	-	expression tag	UNP F4F7F9
C	-22	PHE	-	expression tag	UNP F4F7F9
C	-21	GLN	-	expression tag	UNP F4F7F9
C	-20	GLY	-	expression tag	UNP F4F7F9
C	-19	PRO	-	expression tag	UNP F4F7F9
D	-37	MET	-	initiating methionine	UNP F4F7F9
D	-36	ALA	-	expression tag	UNP F4F7F9
D	-35	HIS	-	expression tag	UNP F4F7F9
D	-34	HIS	-	expression tag	UNP F4F7F9
D	-33	HIS	-	expression tag	UNP F4F7F9
D	-32	HIS	-	expression tag	UNP F4F7F9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-31	HIS	-	expression tag	UNP F4F7F9
D	-30	HIS	-	expression tag	UNP F4F7F9
D	-29	SER	-	expression tag	UNP F4F7F9
D	-28	SER	-	expression tag	UNP F4F7F9
D	-27	GLY	-	expression tag	UNP F4F7F9
D	-26	LEU	-	expression tag	UNP F4F7F9
D	-25	GLU	-	expression tag	UNP F4F7F9
D	-24	VAL	-	expression tag	UNP F4F7F9
D	-23	LEU	-	expression tag	UNP F4F7F9
D	-22	PHE	-	expression tag	UNP F4F7F9
D	-21	GLN	-	expression tag	UNP F4F7F9
D	-20	GLY	-	expression tag	UNP F4F7F9
D	-19	PRO	-	expression tag	UNP F4F7F9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	180	Total	O	0	0
			180	180		
2	B	75	Total	O	0	0
			75	75		
2	C	129	Total	O	0	0
			129	129		
2	D	96	Total	O	0	0
			96	96		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: 3-oxoacyl-ACP synthase III




#### • Molecule 1: 3-oxoacyl-ACP synthase III




#### • Molecule 1: 3-oxoacyl-ACP synthase III



#### • Molecule 1: 3-oxoacyl-ACP synthase III





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	176.71Å 58.69Å 128.80Å 90.00° 95.43° 90.00°	Depositor
Resolution (Å)	55.67 – 1.80 54.31 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (55.67-1.80) 100.0 (54.31-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.24 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, $R_{free}$	0.185 , 0.221 0.198 , 0.199	Depositor DCC
$R_{free}$ test set	6080 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.6	Xtriage
Anisotropy	0.411	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9602	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.01	0/2476	1.04	10/3374 (0.3%)
1	B	1.01	3/2370 (0.1%)	1.06	7/3221 (0.2%)
1	C	1.02	5/2235 (0.2%)	1.01	5/3045 (0.2%)
1	D	0.98	0/2151	1.01	11/2933 (0.4%)
All	All	1.01	8/9232 (0.1%)	1.03	33/12573 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	3
All	All	0	4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	102	SER	CB-OG	6.40	1.50	1.42
1	B	132	ASP	CB-CG	5.69	1.63	1.51
1	C	98	TYR	CG-CD1	-5.38	1.32	1.39
1	C	102	SER	CB-OG	5.33	1.49	1.42
1	B	198	TRP	CB-CG	-5.25	1.40	1.50
1	C	180	SER	CB-OG	-5.25	1.35	1.42
1	C	132	ASP	CB-CG	5.23	1.62	1.51
1	C	321	GLY	C-O	5.11	1.31	1.23

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	219	ARG	NE-CZ-NH1	9.57	125.09	120.30
1	B	219	ARG	NE-CZ-NH2	-8.96	115.82	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	207	ARG	NE-CZ-NH2	-8.71	115.95	120.30
1	A	124	ARG	NE-CZ-NH1	7.88	124.24	120.30
1	D	190	ARG	NE-CZ-NH2	-7.62	116.49	120.30
1	D	204	ASP	CB-CG-OD1	7.57	125.11	118.30
1	D	190	ARG	NE-CZ-NH1	7.35	123.97	120.30
1	C	168	ARG	NE-CZ-NH1	7.22	123.91	120.30
1	A	132	ASP	CB-CG-OD1	-6.87	112.12	118.30
1	B	168	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	D	291	ARG	NE-CZ-NH1	6.46	123.53	120.30
1	A	219	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	1	ASP	CB-CG-OD1	5.94	123.65	118.30
1	D	26	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	D	45	ASP	CB-CG-OD2	-5.79	113.09	118.30
1	B	79	MET	CA-CB-CG	5.76	123.10	113.30
1	D	79	MET	CA-CB-CG	5.75	123.07	113.30
1	A	244	ASP	CB-CG-OD1	-5.58	113.28	118.30
1	A	124	ARG	NE-CZ-NH2	-5.57	117.51	120.30
1	C	168	ARG	NE-CZ-NH2	-5.57	117.51	120.30
1	D	190	ARG	CG-CD-NE	5.52	123.39	111.80
1	A	179	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	C	207	ARG	NE-CZ-NH2	-5.44	117.58	120.30
1	D	204	ASP	CB-CG-OD2	-5.36	113.48	118.30
1	D	93	ASP	CB-CG-OD1	5.32	123.08	118.30
1	B	136	ARG	NE-CZ-NH2	-5.29	117.66	120.30
1	C	287	LEU	CB-CG-CD1	5.25	119.93	111.00
1	B	1	ASP	CB-CG-OD2	-5.24	113.59	118.30
1	A	213	ASP	CB-CG-OD2	-5.21	113.61	118.30
1	D	179	ARG	NE-CZ-NH1	5.15	122.88	120.30
1	C	79	MET	CB-CG-SD	-5.10	97.11	112.40
1	A	287	LEU	CB-CG-CD1	5.07	119.62	111.00
1	A	20	ASP	CB-CG-OD1	5.05	122.84	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	120	GLY	Peptide
1	D	120	GLY	Peptide
1	D	121	GLY	Peptide
1	D	122	THR	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2441	0	2413	18	0
1	B	2341	0	2325	11	0
1	C	2211	0	2173	8	0
1	D	2129	0	2088	6	0
2	A	180	0	0	0	0
2	B	75	0	0	1	0
2	C	129	0	0	0	0
2	D	96	0	0	1	0
All	All	9602	0	8999	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:ARG:CG	1:A:168:ARG:HH11	1.77	0.96
1:C:312:TRP:CZ3	1:C:312:TRP:CZ2	2.47	0.96
1:A:168:ARG:CG	1:A:168:ARG:NH1	2.43	0.82
1:A:129:LEU:HD22	1:A:164:ALA:CB	2.14	0.77
1:A:129:LEU:HD22	1:A:164:ALA:HB2	1.75	0.68
1:A:128:VAL:C	1:A:129:LEU:HD23	2.14	0.66
1:D:122:THR:HA	2:D:484:HOH:O	1.96	0.65
1:A:128:VAL:O	1:A:129:LEU:HD23	1.98	0.64
1:A:250:GLN:HB3	1:A:274:VAL:HB	1.80	0.63
1:B:267:GLN:CB	1:B:267:GLN:CD	2.68	0.61
1:A:154:ASN:OD1	1:A:214:TYR:OH	2.19	0.61
1:B:207:ARG:NH2	2:B:402:HOH:O	2.34	0.61
1:A:-1:THR:HG22	1:A:29:HIS:CE1	2.41	0.56
1:C:250:GLN:HB3	1:C:274:VAL:HB	1.89	0.55
1:A:115:ARG:NH1	1:A:178:LEU:O	2.40	0.54
1:A:129:LEU:HD22	1:A:164:ALA:HB1	1.93	0.51
1:B:250:GLN:HB3	1:B:274:VAL:HB	1.93	0.51
1:B:203:GLU:OE2	1:B:203:GLU:CB	2.60	0.49
1:C:291:ARG:O	1:C:295:ARG:HD3	2.13	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203:GLU:CD	1:B:203:GLU:CB	2.83	0.47
1:B:88:ASP:OD1	1:C:188:LEU:HD12	2.14	0.47
1:D:190:ARG:HD3	1:D:312:TRP:CD1	2.50	0.46
1:A:267:GLN:CB	1:A:267:GLN:CD	2.84	0.45
1:D:250:GLN:HB3	1:D:274:VAL:HB	1.99	0.44
1:A:47:ALA:HB3	1:A:86:VAL:HG21	2.00	0.44
1:B:18:TRP:C	1:B:20:ASP:N	2.71	0.43
1:C:138:TYR:CE2	1:C:158:PHE:CE1	3.05	0.43
1:B:135:ALA:HA	1:B:138:TYR:CE2	2.53	0.43
1:B:250:GLN:NE2	1:B:273:CYS:H	2.16	0.43
1:A:-15:THR:HG21	1:A:171:GLU:HG2	2.00	0.43
1:D:79:MET:HA	1:D:80:PRO:C	2.40	0.42
1:B:4:THR:HG22	1:B:34:LEU:CD2	2.48	0.42
1:C:240:ARG:NH1	1:C:263:LEU:O	2.51	0.42
1:B:-15:THR:HG21	1:B:171:GLU:HG3	2.02	0.42
1:A:135:ALA:HA	1:A:138:TYR:CE2	2.55	0.42
1:A:183:THR:HG22	1:A:315:SER:HB2	2.02	0.42
1:A:240:ARG:NH1	1:A:263:LEU:O	2.53	0.41
1:D:250:GLN:NE2	1:D:273:CYS:H	2.18	0.41
1:D:222:PRO:HG3	1:D:255:MET:CE	2.51	0.41
1:C:250:GLN:NE2	1:C:273:CYS:H	2.18	0.41
1:A:250:GLN:NE2	1:A:273:CYS:H	2.19	0.40
1:C:135:ALA:HA	1:C:138:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/360 (93%)	331 (99%)	4 (1%)	0	100	100
1	B	315/360 (88%)	309 (98%)	6 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	304/360 (84%)	299 (98%)	5 (2%)	0	100	100
1	D	294/360 (82%)	289 (98%)	5 (2%)	0	100	100
All	All	1248/1440 (87%)	1228 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/276 (85%)	233 (99%)	2 (1%)	84	80
1	B	229/276 (83%)	227 (99%)	2 (1%)	84	80
1	C	214/276 (78%)	213 (100%)	1 (0%)	92	91
1	D	201/276 (73%)	200 (100%)	1 (0%)	92	91
All	All	879/1104 (80%)	873 (99%)	6 (1%)	88	86

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	ARG
1	A	158	PHE
1	B	28	ARG
1	B	311	LYS
1	C	28	ARG
1	D	28	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	HIS
1	A	250	GLN
1	B	29	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	250	GLN
1	C	29	HIS
1	C	250	GLN
1	D	29	HIS
1	D	250	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	337/360 (93%)	-0.18	2 (0%)	90 88	5, 13, 29, 43	0
1	B	321/360 (89%)	-0.15	2 (0%)	90 88	4, 12, 36, 55	0
1	C	310/360 (86%)	-0.09	10 (3%)	51 45	5, 14, 35, 50	0
1	D	302/360 (83%)	0.02	10 (3%)	50 44	6, 17, 37, 51	0
All	All	1270/1440 (88%)	-0.10	24 (1%)	70 66	4, 14, 35, 55	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	9	LEU	4.3
1	D	3	ALA	3.8
1	D	6	GLY	3.6
1	D	5	LEU	3.6
1	C	9	LEU	3.4
1	D	122	THR	3.3
1	C	155	TYR	3.1
1	C	6	GLY	3.1
1	C	37	GLY	2.9
1	D	27	THR	2.8
1	D	138	TYR	2.8
1	B	37	GLY	2.6
1	C	23	ILE	2.5
1	D	1	ASP	2.4
1	C	139	ASP	2.4
1	A	206	ASN	2.4
1	D	206	ASN	2.3
1	C	122	THR	2.3
1	A	230	GLU	2.2
1	C	3	ALA	2.2
1	B	12	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	204	ASP	2.1
1	C	35	ASP	2.0
1	C	153	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.