



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:44 AM GMT

PDB ID : 3C16
Title : Complex of GS-Alpha with the Catalytic Domains of Mammalian Adenylyl Cyclase: Complex with Adenosine-5'-Triphosphate and Ca
Authors : Mou, T.-C.; Sprang, S.R.
Deposited on : 2008-01-22
Resolution : 2.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtrriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1945 (2.90-2.86)
Clashscore	102246	2202 (2.90-2.86)
Ramachandran outliers	100387	2149 (2.90-2.86)
Sidechain outliers	100360	2152 (2.90-2.86)
RSRZ outliers	91569	1950 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	 3% 70% 12% • 16%
2	B	212	 2% 69% 17% • 12%
3	C	402	 % 68% 14% 18%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 5739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylate cyclase type 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	190	1484	933	260	274	17	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	MET	-	EXPRESSION TAG	UNP P30803
A	357	HIS	-	EXPRESSION TAG	UNP P30803
A	358	HIS	-	EXPRESSION TAG	UNP P30803
A	359	HIS	-	EXPRESSION TAG	UNP P30803
A	360	HIS	-	EXPRESSION TAG	UNP P30803
A	361	HIS	-	EXPRESSION TAG	UNP P30803
A	362	HIS	-	EXPRESSION TAG	UNP P30803
A	476	MET	VAL	ENGINEERED	UNP P30803

- Molecule 2 is a protein called Adenylate cyclase type 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	186	1443	921	236	276	10	0	0	0

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	331	2710	1718	472	507	13	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	373	GLY	-	EXPRESSION TAG	UNP P04896

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	374	GLY	-	EXPRESSION TAG	UNP P04896
C	375	HIS	-	EXPRESSION TAG	UNP P04896
C	376	HIS	-	EXPRESSION TAG	UNP P04896
C	377	HIS	-	EXPRESSION TAG	UNP P04896
C	378	HIS	-	EXPRESSION TAG	UNP P04896
C	379	HIS	-	EXPRESSION TAG	UNP P04896
C	380	HIS	-	EXPRESSION TAG	UNP P04896

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Cl 1 1	0	0


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total Mg 1 1	0	0

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Ca 2 2	0	0

- Molecule 7 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: C₁₀H₁₆N₅O₁₃P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
9	A	1	31	10	5	13	3	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	6	Total	O	0	0
			6	6		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	118.33Å 133.53Å 70.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.87 24.64 – 2.87	Depositor EDS
% Data completeness (in resolution range)	85.9 (15.00-2.87) 86.1 (24.64-2.87)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.89Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.249 , 0.299 0.246 , 0.297	Depositor DCC
R_{free} test set	1151 reflections (5.42%)	DCC
Wilson B-factor (Å ²)	43.0	Xtrriage
Anisotropy	0.129	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 31.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Outliers	1 of 22537 reflections (0.004%)	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	5739	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSP, MG, CL, CA, ATP, FOK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1512	0.49	0/2038
2	B	0.35	0/1468	0.50	0/1982
3	C	0.35	0/2767	0.50	0/3744
All	All	0.35	0/5747	0.50	0/7764

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1484	0	1454	18	0
2	B	1443	0	1441	23	0
3	C	2710	0	2657	31	0
4	C	1	0	0	1	0
5	C	1	0	0	0	0
6	A	2	0	0	0	0
7	C	32	0	12	2	0
8	A	29	0	34	3	0
9	A	31	0	12	0	0
10	A	6	0	0	0	0
All	All	5739	0	5610	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:62:VAL:HG12	3:C:63:ASN:H	1.46	0.79
2:B:890:ALA:HB1	2:B:996:LEU:HD11	1.65	0.78
8:A:101:FOK:H201	8:A:101:FOK:H173	1.69	0.74
3:C:79:GLU:HA	3:C:82:GLU:HG2	1.70	0.73
8:A:101:FOK:H202	8:A:101:FOK:H193	1.73	0.70
3:C:62:VAL:HG12	3:C:63:ASN:N	2.10	0.67
2:B:891:SER:HA	2:B:943:THR:HA	1.80	0.64
2:B:1059:ARG:HH11	2:B:1062:ILE:HD11	1.66	0.61
3:C:129:LYS:O	3:C:133:GLU:HG2	2.01	0.60
1:A:397:ILE:HD12	1:A:440:ASP:HB2	1.84	0.59
2:B:918:ILE:HG12	2:B:986:ILE:HG13	1.85	0.58
1:A:426:LEU:HD22	1:A:465:ASP:HB3	1.85	0.58
2:B:1062:ILE:HD12	2:B:1072:THR:HG21	1.87	0.57
3:C:144:SER:HA	3:C:147:TYR:CE2	2.39	0.57
3:C:121:PRO:HB2	3:C:123:GLU:OE2	2.05	0.57
3:C:66:LYS:HE3	3:C:175:LEU:HD11	1.88	0.56
2:B:900:THR:HB	2:B:905:ASN:ND2	2.22	0.55
1:A:407:CYS:HB2	1:A:411:GLU:HB3	1.91	0.53
2:B:977:TYR:O	2:B:980:VAL:HB	2.09	0.52
2:B:900:THR:HB	2:B:905:ASN:HD21	1.73	0.52
1:A:526:ILE:HB	1:A:564:ILE:HD12	1.91	0.51
2:B:904:VAL:HG11	3:C:185:ILE:HB	1.93	0.51
3:C:187:GLU:OE2	3:C:198:HIS:ND1	2.44	0.51
3:C:344:ALA:HB3	7:C:395:GSP:N7	2.26	0.51
3:C:255:TRP:CE2	3:C:327:SER:HA	2.46	0.51
3:C:62:VAL:CG1	3:C:63:ASN:H	2.19	0.50
3:C:79:GLU:HA	3:C:82:GLU:CG	2.41	0.50
2:B:921:ASP:HB3	2:B:982:LYS:HD2	1.93	0.50
3:C:77:LEU:HD21	3:C:160:ILE:HG12	1.93	0.50
3:C:270:ASN:HA	3:C:342:THR:O	2.11	0.50
1:A:530:LYS:HD2	1:A:531:ALA:N	2.27	0.49
3:C:78:LYS:HA	3:C:124:PHE:HZ	1.77	0.49
3:C:70:VAL:HA	3:C:73:ILE:HD12	1.94	0.49
1:A:388:ASP:OD1	1:A:491:ARG:HB3	2.13	0.49
1:A:401:THR:HG21	2:B:1066:GLY:HA3	1.93	0.48
3:C:177:ARG:HA	7:C:395:GSP:O2'	2.12	0.48
2:B:891:SER:HB3	2:B:943:THR:HG23	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:995:LYS:HB2	2:B:1036:LEU:HD22	1.96	0.48
3:C:105:ARG:O	3:C:109:ILE:HG12	2.13	0.47
2:B:940:ILE:HB	2:B:943:THR:HB	1.96	0.47
8:A:101:FOK:H201	8:A:101:FOK:C17	2.42	0.46
2:B:892:ILE:HD12	2:B:942:SER:HB2	1.98	0.46
1:A:388:ASP:HB3	1:A:389:ASN:ND2	2.31	0.46
3:C:235:ILE:HG12	3:C:241:THR:O	2.15	0.45
2:B:1049:LEU:HB3	2:B:1054:TYR:HB2	1.99	0.45
3:C:144:SER:HB2	3:C:149:LEU:HD23	1.97	0.45
3:C:62:VAL:CG1	3:C:63:ASN:N	2.79	0.45
1:A:438:LEU:HB2	1:A:441:CYS:HB3	1.98	0.45
3:C:206:ARG:HG2	3:C:209:ARG:NH1	2.32	0.45
1:A:488:HIS:CD2	1:A:489:SER:O	2.70	0.45
3:C:275:LEU:O	3:C:279:VAL:HG23	2.17	0.44
1:A:488:HIS:CG	1:A:489:SER:N	2.86	0.43
3:C:43:LEU:O	3:C:224:PHE:HA	2.17	0.43
3:C:109:ILE:HG13	3:C:131:LEU:CD1	2.48	0.43
3:C:227:ALA:HB3	4:C:397:CL:CL	2.56	0.43
1:A:536:LEU:HD11	1:A:540:TYR:CD1	2.54	0.42
3:C:205:GLN:O	3:C:209:ARG:HG3	2.20	0.42
2:B:1059:ARG:HG3	2:B:1074:PHE:CZ	2.54	0.41
3:C:342:THR:HG22	3:C:353:VAL:HG21	2.03	0.41
1:A:383:TYR:C	1:A:384:ILE:HG13	2.39	0.41
2:B:995:LYS:HD2	2:B:1037:ASP:OD2	2.20	0.41
1:A:420:PHE:HA	1:A:423:PHE:HB2	2.03	0.41
2:B:936:LYS:HD2	2:B:944:TYR:OH	2.20	0.41
2:B:1009:VAL:HG11	2:B:1015:PRO:HB3	2.01	0.41
3:C:296:TYR:OH	3:C:299:PRO:HD3	2.21	0.41
1:A:509:ASN:O	1:A:513:LEU:N	2.50	0.41
2:B:895:PHE:CZ	2:B:915:LEU:HB2	2.56	0.41
2:B:1040:GLN:HA	2:B:1073:TYR:O	2.21	0.41
1:A:507:TRP:O	1:A:508:SER:HB3	2.20	0.40
1:A:423:PHE:HA	1:A:426:LEU:HD12	2.03	0.40
2:B:1030:MET:O	2:B:1040:GLN:NE2	2.50	0.40
3:C:267:LEU:HB3	3:C:339:PRO:HA	2.04	0.40
1:A:551:ASN:HB3	1:A:554:LEU:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/225 (84%)	171 (91%)	17 (9%)	0	100	100
2	B	182/212 (86%)	174 (96%)	6 (3%)	2 (1%)	17	49
3	C	327/402 (81%)	306 (94%)	19 (6%)	2 (1%)	30	65
All	All	697/839 (83%)	651 (93%)	42 (6%)	4 (1%)	30	65

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	116	PRO
2	B	1065	LYS
3	C	153	ALA
2	B	1013	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/189 (84%)	151 (95%)	8 (5%)	30	64
2	B	160/184 (87%)	156 (98%)	4 (2%)	55	84
3	C	298/357 (84%)	286 (96%)	12 (4%)	38	73
All	All	617/730 (84%)	593 (96%)	24 (4%)	39	74

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	382	ILE
1	A	388	ASP
1	A	398	GLU
1	A	401	THR
1	A	416	LEU
1	A	491	ARG
1	A	507	TRP
1	A	530	LYS
2	B	900	THR
2	B	1059	ARG
2	B	1061	ILE
2	B	1077	THR
3	C	39	ARG
3	C	68	THR
3	C	117	ASP
3	C	154	GLN
3	C	159	LYS
3	C	185	ILE
3	C	207	ASP
3	C	225	VAL
3	C	262	THR
3	C	334	ARG
3	C	348	GLU
3	C	359	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	385	GLN
1	A	537	ASN
2	B	880	GLN
2	B	1001	ASN
2	B	1050	GLN
2	B	1076	ASN
3	C	75	ASN
3	C	245	GLN
3	C	349	ASN
3	C	355	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	ATP	A	1	6	24,33,33	1.45	3 (12%)	31,52,52	2.08	5 (16%)
8	FOK	A	101	-	26,31,31	1.04	1 (3%)	35,54,54	1.75	7 (20%)
7	GSP	C	395	5	25,34,34	2.12	2 (8%)	31,54,54	1.82	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	ATP	A	1	6	-	0/18/38/38	0/3/3/3
8	FOK	A	101	-	-	0/7/80/80	0/3/3/3
7	GSP	C	395	5	-	0/15/38/38	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	395	GSP	PG-S1G	-9.50	1.72	1.90
9	A	1	ATP	PG-O2G	2.30	1.63	1.54
7	C	395	GSP	C6-N1	2.49	1.37	1.33
9	A	1	ATP	PG-O1G	3.69	1.63	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	1	ATP	O4'-C1'	3.79	1.46	1.41
8	A	101	FOK	O4-C21	4.48	1.45	1.35

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	1	ATP	N3-C2-N1	-8.96	122.03	128.89
7	C	395	GSP	N3-C2-N1	-4.74	120.22	127.44
7	C	395	GSP	PA-O3A-PB	-4.48	120.14	132.73
8	A	101	FOK	C7-O4-C21	-4.37	111.49	117.83
7	C	395	GSP	PB-O3B-PG	-3.57	120.71	132.67
7	C	395	GSP	C2'-C1'-N9	-3.27	109.29	114.29
9	A	1	ATP	PA-O3A-PB	-3.12	123.96	132.73
9	A	1	ATP	PB-O3B-PG	-2.99	122.64	132.67
9	A	1	ATP	C4'-O4'-C1'	-2.90	106.53	109.72
7	C	395	GSP	C5-C6-N1	-2.61	120.01	123.59
9	A	1	ATP	C4-C5-N7	-2.57	107.12	109.48
8	A	101	FOK	O7-C11-C12	-2.02	117.20	121.99
8	A	101	FOK	C12-C11-C9	2.18	120.59	115.09
8	A	101	FOK	C3-C4-C5	2.43	111.61	107.73
8	A	101	FOK	O4-C7-C6	2.57	112.84	107.92
7	C	395	GSP	C6-N1-C2	2.66	119.62	115.94
8	A	101	FOK	O4-C21-C22	3.47	117.65	111.10
8	A	101	FOK	C13-O1-C8	4.59	124.85	120.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	101	FOK	3	0
7	C	395	GSP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	190/225 (84%)	0.04	6 (3%) 51 45	18, 51, 82, 86	0
2	B	186/212 (87%)	-0.22	4 (2%) 65 62	16, 31, 71, 85	0
3	C	331/402 (82%)	-0.21	5 (1%) 76 74	17, 34, 66, 86	0
All	All	707/839 (84%)	-0.15	15 (2%) 67 63	16, 37, 73, 86	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	475	GLU	4.1
1	A	480	ASN	4.0
1	A	376	ASP	3.9
1	A	479	VAL	3.4
2	B	1012	ALA	3.4
2	B	1063	ASN	3.3
3	C	365	HIS	3.2
3	C	364	MET	2.8
2	B	1065	LYS	2.7
3	C	332	ASP	2.6
1	A	377	MET	2.3
3	C	63	ASN	2.2
1	A	565	LEU	2.2
2	B	1064	VAL	2.2
3	C	362	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CL	C	397	1/1	0.98	0.22	1.09	30,30,30,30	0
8	FOK	A	101	29/29	0.94	0.18	0.84	16,33,49,50	0
7	GSP	C	395	32/32	0.93	0.16	-0.04	17,22,32,41	0
9	ATP	A	1	31/31	0.91	0.17	-0.19	31,51,64,67	0
6	CA	A	29	1/1	0.89	0.06	-	49,49,49,49	0
6	CA	A	30	1/1	0.95	0.05	-	38,38,38,38	0
5	MG	C	398	1/1	0.97	0.18	-	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.