



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:18 PM GMT

PDB ID : 4C2E
Title : Crystal structure of the protease CtpB(S309A) present in a resting state
Authors : Mastny, M.; Heuck, A.; Kurzbauer, R.; Clausen, T.
Deposited on : 2013-08-17
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	<div> <div>2%</div> <div>88%</div> <div>9%</div> <div>.</div> </div>
1	B	446	<div> <div>2%</div> <div>87%</div> <div>10%</div> <div>..</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7779 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CARBOXY-TERMINAL PROCESSING PROTEASE CTPB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	0	1
			3397	2144	580	663	10			
1	B	436	Total	C	N	O	S	0	0	1
			3397	2144	580	663	10			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	MET	-	INITIATING METHIONINE	UNP O35002
A	481	LEU	-	EXPRESSION TAG	UNP O35002
A	482	GLU	-	EXPRESSION TAG	UNP O35002
A	483	HIS	-	EXPRESSION TAG	UNP O35002
A	484	HIS	-	EXPRESSION TAG	UNP O35002
A	485	HIS	-	EXPRESSION TAG	UNP O35002
A	486	HIS	-	EXPRESSION TAG	UNP O35002
A	487	HIS	-	EXPRESSION TAG	UNP O35002
A	488	HIS	-	EXPRESSION TAG	UNP O35002
A	227	THR	ALA	CONFLICT	UNP O35002
A	257	ILE	LEU	CONFLICT	UNP O35002
A	309	ALA	SER	ENGINEERED MUTATION	UNP O35002
A	444	ILE	VAL	CONFLICT	UNP O35002
B	43	MET	-	INITIATING METHIONINE	UNP O35002
B	481	LEU	-	EXPRESSION TAG	UNP O35002
B	482	GLU	-	EXPRESSION TAG	UNP O35002
B	483	HIS	-	EXPRESSION TAG	UNP O35002
B	484	HIS	-	EXPRESSION TAG	UNP O35002
B	485	HIS	-	EXPRESSION TAG	UNP O35002
B	486	HIS	-	EXPRESSION TAG	UNP O35002
B	487	HIS	-	EXPRESSION TAG	UNP O35002
B	488	HIS	-	EXPRESSION TAG	UNP O35002
B	227	THR	ALA	CONFLICT	UNP O35002
B	257	ILE	LEU	CONFLICT	UNP O35002
B	309	ALA	SER	ENGINEERED MUTATION	UNP O35002

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	444	ILE	VAL	CONFLICT	UNP O35002


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	447	Total O 447 447	0	0
2	B	538	Total O 538 538	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CARBOXY-TERMINAL PROCESSING PROTEASE CTPB



• Molecule 1: CARBOXY-TERMINAL PROCESSING PROTEASE CTPB



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.05Å 72.69Å 79.57Å 117.09° 90.45° 102.71°	Depositor
Resolution (Å)	20.00 – 1.80 19.86 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.7 (20.00-1.80) 80.0 (19.86-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.97 (at 1.80Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.182 , 0.191 0.181 , 0.189	Depositor DCC
R_{free} test set	4708 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.525	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 93488 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7779	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3452	0.55	0/4636
1	B	0.39	0/3452	0.57	0/4636
All	All	0.38	0/6904	0.56	0/9272

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3397	0	3446	37	0
1	B	3397	0	3446	41	0
2	A	447	0	0	10	0
2	B	538	0	0	16	0
All	All	7779	0	6892	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:ASP:OD2	1:B:72:ARG:NH2	1.86	1.08

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:LYS:HB2	1:A:399:GLU:HG2	1.36	1.02
1:B:367:LYS:NZ	2:B:2414:HOH:O	2.08	0.87
1:B:370:GLU:OE2	2:B:2395:HOH:O	1.98	0.81
1:B:196:GLU:N	1:B:196:GLU:OE1	2.11	0.78
1:A:212:HIS:ND1	1:A:243:GLU:OE1	2.21	0.73
1:B:464:GLU:OE2	2:B:2526:HOH:O	2.05	0.73
1:A:73:GLU:O	1:A:77:GLU:HG3	1.89	0.72
1:A:236:GLU:O	1:A:240:LYS:HD3	1.89	0.72
1:A:142:LYS:HE3	1:A:382:SER:HA	1.73	0.70
1:A:121:GLU:O	1:A:124:LYS:HG2	1.91	0.70
2:A:2020:HOH:O	1:B:280:ARG:NH1	2.08	0.69
1:A:72:ARG:HH11	1:B:47:GLU:HG3	1.58	0.67
1:A:73:GLU:OE1	2:A:2043:HOH:O	2.13	0.67
1:B:124:LYS:NZ	2:B:2135:HOH:O	2.29	0.65
1:B:280:ARG:NH2	2:B:2345:HOH:O	2.18	0.64
1:A:394:VAL:HG21	1:A:427:LYS:HD2	1.79	0.64
1:B:236:GLU:O	1:B:240:LYS:HD2	2.00	0.61
1:A:186:LYS:O	2:A:2228:HOH:O	2.16	0.61
1:A:147:ILE:HA	1:A:179:ILE:HD13	1.82	0.60
1:A:186:LYS:HB2	1:A:399:GLU:CG	2.23	0.60
1:B:285:LYS:HE3	1:B:287:TYR:OH	2.02	0.59
1:B:45:ASP:OD2	1:B:48:ARG:HB2	2.03	0.58
1:B:286:ARG:NE	2:B:2341:HOH:O	2.36	0.58
1:B:186:LYS:HG2	1:B:187:GLN:N	2.19	0.58
1:B:436:GLN:NE2	2:B:2489:HOH:O	2.18	0.56
1:B:444:ILE:O	1:B:444:ILE:HG13	2.04	0.56
1:A:240:LYS:N	1:A:240:LYS:HD3	2.20	0.56
1:A:399:GLU:HG3	2:A:2399:HOH:O	2.05	0.55
1:B:367:LYS:HE2	2:B:2417:HOH:O	2.05	0.55
1:A:98:GLN:HB3	2:A:2092:HOH:O	2.08	0.54
1:A:364:ILE:HG13	2:A:2369:HOH:O	2.09	0.53
1:A:444:ILE:HD11	2:A:2418:HOH:O	2.07	0.53
1:A:394:VAL:CG2	1:A:427:LYS:HB2	2.39	0.53
1:B:261:GLU:O	1:B:265:LYS:HG3	2.09	0.52
1:B:465:LYS:HG2	2:B:2273:HOH:O	2.09	0.52
1:B:376:LYS:HE3	2:B:2389:HOH:O	2.09	0.52
1:A:72:ARG:NH1	1:B:47:GLU:HG3	2.25	0.51
1:A:124:LYS:HG3	1:A:126:ILE:HG23	1.92	0.51
1:A:264:LEU:CD2	1:A:314:LEU:HD12	2.41	0.50
1:B:198:PRO:HD2	2:B:2254:HOH:O	2.12	0.50
1:A:125:ILE:HD13	1:A:163:ALA:HB2	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:VAL:HG21	1:B:167:ILE:CD1	2.43	0.49
1:A:264:LEU:HD21	1:A:314:LEU:HD12	1.93	0.49
1:B:441:LYS:HB2	1:B:441:LYS:NZ	2.29	0.48
1:B:212:HIS:ND1	1:B:243:GLU:OE1	2.47	0.48
1:A:394:VAL:HG22	1:A:427:LYS:HB2	1.97	0.47
1:A:240:LYS:N	1:A:240:LYS:CD	2.77	0.47
1:A:97:LYS:HE2	2:A:2045:HOH:O	2.15	0.47
1:A:72:ARG:HH11	1:B:47:GLU:CG	2.27	0.46
1:A:92:SER:HA	1:A:350:LYS:O	2.17	0.45
1:A:435:ASP:OD1	1:A:441:LYS:HE2	2.17	0.44
1:B:49:ASP:HB3	2:B:2002:HOH:O	2.17	0.44
1:B:364:ILE:HG13	2:B:2411:HOH:O	2.17	0.44
1:B:120:MET:HE1	1:B:159:ASP:HA	2.00	0.43
1:B:116:ALA:HB2	1:B:136:ALA:CB	2.47	0.43
1:A:375:ILE:HD11	1:A:468:LEU:HD21	2.00	0.43
1:B:285:LYS:CE	1:B:287:TYR:OH	2.65	0.43
1:B:285:LYS:HE3	1:B:287:TYR:CZ	2.55	0.42
1:B:394:VAL:CG1	1:B:427:LYS:HB2	2.50	0.42
1:B:63:SER:CB	1:B:72:ARG:HH11	2.33	0.41
1:B:270:LYS:NZ	2:B:2334:HOH:O	2.39	0.41
1:B:418:GLU:HG2	2:B:2465:HOH:O	2.19	0.41
1:B:92:SER:HA	1:B:350:LYS:O	2.20	0.41
1:A:98:GLN:HG2	2:A:2097:HOH:O	2.21	0.41
1:A:186:LYS:HE2	1:A:400:ASP:OD1	2.21	0.41
1:B:115:GLY:HA3	1:B:134:SER:HB2	2.01	0.41
1:A:350:LYS:HB3	1:A:350:LYS:HZ2	1.85	0.41
1:A:402:LYS:HG3	1:A:418:GLU:O	2.20	0.41
1:B:427:LYS:NZ	2:B:2474:HOH:O	2.54	0.41
1:A:177:MET:HE3	1:A:192:ILE:HD12	2.02	0.41
1:A:132:LYS:HG2	2:A:2162:HOH:O	2.21	0.41
1:B:118:VAL:HG21	1:B:167:ILE:HD12	2.02	0.40
1:B:260:VAL:CG2	1:B:314:LEU:HB2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/446 (97%)	425 (98%)	9 (2%)	0	100	100
1	B	434/446 (97%)	428 (99%)	6 (1%)	0	100	100
All	All	868/892 (97%)	853 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/381 (97%)	365 (99%)	5 (1%)	74	65
1	B	370/381 (97%)	363 (98%)	7 (2%)	65	52
All	All	740/762 (97%)	728 (98%)	12 (2%)	70	59

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	GLN
1	A	132	LYS
1	A	184	THR
1	A	209	VAL
1	A	422	PHE
1	B	45	ASP
1	B	117	GLU
1	B	189	SER
1	B	240	LYS
1	B	367	LYS
1	B	422	PHE
1	B	444	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/446 (97%)	-0.13	9 (2%) 67 62	15, 27, 48, 71	0
1	B	436/446 (97%)	-0.22	11 (2%) 61 56	14, 23, 43, 78	0
All	All	872/892 (97%)	-0.18	20 (2%) 64 59	14, 25, 46, 78	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	44	ALA	6.1
1	A	44	ALA	5.6
1	A	479	VAL	4.5
1	A	183	GLY	4.3
1	B	108	ASP	3.6
1	B	185	LYS	3.3
1	B	389	LYS	3.1
1	B	45	ASP	2.9
1	A	185	LYS	2.8
1	A	389	LYS	2.8
1	A	45	ASP	2.5
1	B	46	SER	2.4
1	A	401	VAL	2.4
1	A	210	GLN	2.1
1	B	107	LEU	2.1
1	B	313	ILE	2.1
1	B	109	SER	2.1
1	B	62	ILE	2.1
1	A	122	ASP	2.1
1	B	122	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.