



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 21, 2016 – 05:31 AM EDT

PDB ID : 5C21  
Title : Crystal structure of native HlyD from E. coli  
Authors : Ha, N.C.; Kim, J.S.; Yoon, B.Y.  
Deposited on : 2015-06-15  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0122  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>...</div> </div> </div>
1	B	279	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>12%</div> <div>6%</div> </div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chromosomal hemolysin D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2171	1364	379	426	2			
1	B	261	Total	C	N	O	S	0	0	0
			2116	1328	371	415	2			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	MET	-	expression tag	UNP O87505
A	95	ALA	-	expression tag	UNP O87505
B	94	MET	-	expression tag	UNP O87505
B	95	ALA	-	expression tag	UNP O87505


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Chromosomal hemolysin D



#### • Molecule 1: Chromosomal hemolysin D



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.61Å 93.17Å 163.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.88 – 2.50 19.88 – 2.19	Depositor EDS
% Data completeness (in resolution range)	97.3 (19.88-2.50) 94.7 (19.88-2.19)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.80 (at 2.19Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.235 , 0.274 0.237 , 0.279	Depositor DCC
$R_{free}$ test set	1382 reflections (3.41%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.8	Xtriage
Anisotropy	0.352	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 52.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	5 of 58644 reflections (0.009%)	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4287	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 19.98 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.6057e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/2190	0.44	0/2945
1	B	0.28	0/2132	0.50	1/2865 (0.0%)
All	All	0.27	0/4322	0.47	1/5810 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	244	LEU	CA-CB-CG	-6.04	101.41	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2171	0	2259	16	1
1	B	2116	0	2216	24	1
All	All	4287	0	4475	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:GLU:HB2	1:B:168:LEU:HA	1.53	0.90
1:A:125:LEU:HD13	1:A:127:LEU:HD13	1.71	0.73
1:B:240:PHE:HB3	1:B:254:VAL:HG22	1.78	0.65
1:A:189:SER:O	1:A:193:GLU:HG3	1.98	0.64
1:A:180:SER:OG	1:A:183:GLU:OE1	2.16	0.64
1:B:180:SER:OG	1:B:182:GLU:HG2	2.00	0.61
1:B:301:LEU:HA	1:B:304:LEU:HB2	1.83	0.60
1:B:152:ARG:HH21	1:B:194:GLN:HE22	1.49	0.60
1:A:107:VAL:HA	1:A:127:LEU:HD12	1.84	0.59
1:A:162:LEU:HB3	1:A:164:LYS:HG2	1.85	0.58
1:A:350:VAL:HG21	1:A:356:LEU:HD21	1.87	0.56
1:A:231:ARG:NH2	1:B:206:GLU:OE2	2.42	0.53
1:B:126:LYS:HG2	1:B:329:VAL:HG12	1.93	0.51
1:B:180:SER:HB3	1:B:183:GLU:OE1	2.13	0.49
1:A:203:TYR:CE1	1:B:236:ARG:HD2	2.47	0.49
1:B:150:GLN:O	1:B:154:GLN:HG3	2.14	0.47
1:B:125:LEU:HD21	1:B:356:LEU:HD13	1.97	0.47
1:B:134:ALA:HA	1:B:137:LEU:HD12	1.95	0.47
1:A:240:PHE:HB3	1:A:254:VAL:HG22	1.96	0.47
1:B:244:LEU:HD13	1:B:249:ILE:O	2.15	0.46
1:B:244:LEU:HD12	1:B:244:LEU:HA	1.61	0.45
1:A:126:LYS:HG2	1:A:329:VAL:HG12	1.99	0.45
1:B:96:SER:HB3	1:B:360:VAL:HG12	1.99	0.45
1:A:233:GLU:OE1	1:A:257:GLN:NE2	2.33	0.44
1:A:322:GLU:O	1:A:325:GLN:HG2	2.17	0.44
1:B:311:ILE:HD13	1:B:311:ILE:HA	1.85	0.44
1:A:170:LEU:HD11	1:A:184:VAL:HG22	2.00	0.44
1:B:107:VAL:HB	1:B:345:THR:O	2.18	0.43
1:B:167:GLU:HB2	1:B:168:LEU:CA	2.36	0.43
1:A:181:GLU:N	1:A:181:GLU:OE1	2.52	0.42
1:A:226:TYR:CZ	1:A:267:GLU:HG2	2.53	0.42
1:B:231:ARG:HA	1:B:231:ARG:HD2	1.92	0.42
1:B:304:LEU:HD23	1:B:304:LEU:HA	1.81	0.42
1:A:216:ARG:NH1	1:A:279:GLU:OE2	2.41	0.41
1:B:167:GLU:HB3	1:B:169:LYS:H	1.85	0.41
1:B:128:THR:HG22	1:B:327:ALA:HA	2.02	0.41
1:B:167:GLU:CB	1:B:168:LEU:HA	2.38	0.41
1:B:189:SER:O	1:B:193:GLU:HG3	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:THR:O	1:B:100:LYS:NZ[1_544]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/279 (94%)	260 (99%)	3 (1%)	0	100	100
1	B	257/279 (92%)	254 (99%)	3 (1%)	0	100	100
All	All	520/558 (93%)	514 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/257 (96%)	243 (99%)	3 (1%)	78	93
1	B	240/257 (93%)	238 (99%)	2 (1%)	86	96
All	All	486/514 (95%)	481 (99%)	5 (1%)	82	95

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	127	LEU
1	A	231	ARG
1	A	363	ASP

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	137	LEU
1	B	165	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	267/279 (95%)	0.18	5 (1%) 70 73	36, 61, 92, 111	0
1	B	261/279 (93%)	0.21	10 (3%) 44 49	40, 67, 94, 110	0
All	All	528/558 (94%)	0.19	15 (2%) 56 61	36, 64, 93, 111	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	178	ASN	5.4
1	B	301	LEU	3.1
1	A	177	GLN	2.9
1	A	344	HIS	2.8
1	A	94	MET	2.7
1	B	252	HIS	2.6
1	B	179	VAL	2.6
1	B	316	LEU	2.4
1	A	301	LEU	2.3
1	B	344	HIS	2.3
1	B	124	LEU	2.2
1	B	305	ARG	2.2
1	B	189	SER	2.2
1	B	182	GLU	2.2
1	B	324	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.