



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:14 AM GMT

PDB ID : 2C49
Title : CRYSTAL STRUCTURE OF METHANOCALDOCOCUS JANNASCHII
NUCLEOSIDE KINASE - AN ARCHAEAL MEMBER OF THE RIBOKI-
NASE FAMILY
Authors : Arnfors, L.; Hansen, T.; Meining, W.; Schoenheit, P.; Ladenstein, R.
Deposited on : 2005-10-17
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5832 (1.94-1.90)
Clashscore	102246	6540 (1.94-1.90)
Ramachandran outliers	100387	6464 (1.94-1.90)
Sidechain outliers	100360	6465 (1.94-1.90)
RSRZ outliers	91569	5846 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	<div> <div>11%</div> <div>81%</div> <div>15%</div> <div>..</div> </div>
1	B	302	<div> <div>12%</div> <div>75%</div> <div>21%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ADN	B	1301	-	-	-	X
4	ANP	A	1304	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 5004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUGAR KINASE MJ0406.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	0	0
			2371	1521	390	451	9			
1	B	298	Total	C	N	O	S	0	2	0
			2367	1520	387	451	9			

- Molecule 2 is ADENOSINE (three-letter code: ADN) (formula: C₁₀H₁₃N₅O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	10	5	4		
2	B	1	Total	C	N	O	0	0
			19	10	5	4		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Mg 2 2	0	0
3	A	2	Total Mg 2 2	0	0

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 31 10 6 12 3	0	0
4	B	1	Total C N O P 31 10 6 12 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	80	Total O 80 80	0	0
5	B	82	Total O 82 82	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SUGAR KINASE MJ0406



• Molecule 1: SUGAR KINASE MJ0406



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.01Å 83.13Å 146.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.55 – 1.92 48.24 – 1.92	Depositor EDS
% Data completeness (in resolution range)	97.1 (72.55-1.92) 97.0 (48.24-1.92)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.243 , 0.281 0.240 , 0.279	Depositor DCC
R_{free} test set	2955 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 58475 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5004	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.03 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.9426e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADN, MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/2425	0.82	3/3277 (0.1%)
1	B	0.70	0/2429	0.81	7/3284 (0.2%)
All	All	0.71	0/4854	0.81	10/6561 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	ASP	CB-CG-OD2	6.54	124.19	118.30
1	B	68	ASP	CB-CG-OD2	6.54	124.19	118.30
1	A	202	ASP	CB-CG-OD2	6.12	123.80	118.30
1	B	140	ASP	CB-CG-OD2	6.08	123.78	118.30
1	B	291	ASP	CB-CG-OD2	5.93	123.63	118.30
1	A	140	ASP	CB-CG-OD2	5.78	123.50	118.30
1	B	104	ASP	CB-CG-OD1	5.61	123.35	118.30
1	B	262	ASP	CB-CG-OD1	5.36	123.12	118.30
1	B	202	ASP	CB-CG-OD2	5.06	122.86	118.30
1	B	250	ARG	NE-CZ-NH1	-5.05	117.77	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	69	PHE	Peptide
1	B	70	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2371	0	2342	35	0
1	B	2367	0	2336	44	0
2	A	19	0	13	3	0
2	B	19	0	13	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	31	0	13	1	0
4	B	31	0	13	4	0
5	A	80	0	0	2	0
5	B	82	0	0	1	0
All	All	5004	0	4730	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:ARG:HG3	1:A:192:ARG:HH11	1.03	1.10
1:B:22:VAL:HG21	1:B:32:ILE:HD12	1.32	1.09
1:B:22:VAL:HG21	1:B:32:ILE:CD1	1.99	0.92
1:B:5:MET:HG3	1:B:7:LYS:HB3	1.55	0.86
2:A:1301:ADN:HN62	1:B:33:GLN:HE21	1.20	0.85
1:B:157:VAL:H	1:B:181:ASN:HD22	1.23	0.85
1:B:5:MET:SD	1:B:7:LYS:HD3	2.17	0.83
1:A:192:ARG:HG3	1:A:192:ARG:NH1	1.83	0.82
1:A:192:ARG:HH11	1:A:192:ARG:CG	1.88	0.82
1:A:163:GLN:HE21	1:A:163:GLN:H	1.23	0.81
1:A:157:VAL:H	1:A:181:ASN:HD22	1.31	0.79
1:A:300:HIS:CD2	1:A:301:ARG:H	2.02	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:HIS:O	1:A:192:ARG:HG2	1.86	0.74
1:B:5:MET:O	1:B:132:GLU:HG3	1.87	0.74
1:B:233:CYS:SG	5:B:2064:HOH:O	2.46	0.73
1:B:5:MET:HG3	1:B:7:LYS:CB	2.18	0.72
1:B:22:VAL:CG2	1:B:32:ILE:HD12	2.17	0.69
1:A:178:GLU:HG2	1:A:207:ARG:HG2	1.76	0.67
1:A:145:LEU:HD11	1:A:175:GLU:HG2	1.77	0.66
1:A:163:GLN:H	1:A:163:GLN:NE2	1.97	0.61
1:A:300:HIS:CG	1:A:301:ARG:H	2.18	0.60
1:A:84:ASN:HD22	1:A:84:ASN:C	2.05	0.60
1:B:103:THR:HB	1:B:109:GLN:HG3	1.84	0.59
1:B:19:ILE:HD13	1:B:101:ILE:HB	1.86	0.58
1:B:7:LYS:HG2	1:B:131:THR:HG22	1.86	0.57
1:B:70:LYS:HE3	1:B:90:TYR:HD2	1.69	0.57
1:A:119:LYS:HE3	1:A:120:HIS:CE1	2.40	0.57
1:B:240:ILE:HD12	1:B:279:GLU:O	2.05	0.56
1:B:159:PHE:CD2	1:B:176:ILE:HD12	2.40	0.55
1:B:298:GLU:OE2	1:B:299:LYS:HD3	2.06	0.55
1:B:216:GLY:O	4:B:1304:ANP:H2'	2.06	0.55
1:B:175:GLU:O	1:B:179:HIS:HD2	1.90	0.55
1:A:277:VAL:CG2	1:A:287:LEU:HD23	2.37	0.55
1:A:5:MET:O	1:A:132:GLU:HG3	2.08	0.54
1:B:215:LYS:HE3	1:B:221[B]:VAL:HG23	1.90	0.54
1:B:178:GLU:HG2	1:B:207:ARG:HG2	1.89	0.54
1:B:242:PRO:HA	1:B:278:VAL:HG12	1.91	0.52
1:B:70:LYS:HG2	1:B:90:TYR:CD2	2.44	0.52
1:B:243:THR:HA	4:B:1304:ANP:O1G	2.11	0.51
1:B:243:THR:HG23	4:B:1304:ANP:O1G	2.11	0.51
1:A:192:ARG:NH1	1:A:192:ARG:CG	2.53	0.50
1:A:84:ASN:HD22	1:A:86:SER:H	1.60	0.50
1:A:277:VAL:HG21	1:A:287:LEU:HD23	1.93	0.49
1:B:7:LYS:HA	1:B:58:ASN:O	2.12	0.48
1:B:289:THR:OG1	1:B:292:LYS:HD2	2.12	0.48
1:A:114:LEU:HD12	1:B:34:ILE:HG12	1.95	0.48
1:A:84:ASN:ND2	1:A:86:SER:H	2.12	0.47
1:B:21:ASN:HD21	1:B:38:ARG:HH12	1.62	0.47
1:B:216:GLY:HA2	4:B:1304:ANP:H8	1.97	0.47
1:A:60:GLU:HB3	1:A:84:ASN:HB3	1.97	0.46
1:B:84:ASN:ND2	1:B:86:SER:H	2.14	0.46
1:B:66:GLY:O	1:B:70:LYS:HG3	2.16	0.45
1:A:175:GLU:O	1:A:179:HIS:HD2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:SER:HB2	1:B:120:HIS:CD2	2.51	0.45
1:B:46:ALA:HA	1:B:61:LEU:HD21	1.99	0.45
1:A:119:LYS:O	1:A:122:LYS:HG2	2.17	0.45
1:A:234:ILE:HD13	1:A:272:ALA:HB2	1.99	0.44
1:A:185:MET:CE	1:A:190:PHE:HD1	2.29	0.44
2:B:1301:ADN:H2'	2:B:1301:ADN:H8	1.80	0.43
1:B:57:VAL:HG23	1:B:259:LYS:HE2	1.99	0.43
1:A:246:GLY:O	1:A:250:ARG:HD3	2.18	0.43
1:A:268:LEU:HB3	1:A:297:LEU:HD21	2.01	0.43
1:A:109:GLN:HE21	4:A:1304:ANP:PG	2.41	0.43
1:A:287:LEU:HA	1:A:288:PRO:HD3	1.90	0.43
1:A:72:SER:CB	5:A:2020:HOH:O	2.67	0.42
1:B:70:LYS:HE3	1:B:90:TYR:CD2	2.51	0.42
1:B:141:PRO:C	1:B:172:MET:HE1	2.39	0.42
2:A:1301:ADN:HN61	1:B:31:SER:HB3	1.85	0.42
1:B:70:LYS:HG2	1:B:90:TYR:CE2	2.55	0.42
1:B:159:PHE:CZ	1:B:161:PRO:HB3	2.55	0.41
1:A:172:MET:HE3	1:A:172:MET:HB3	1.96	0.41
1:B:46:ALA:O	1:B:50:VAL:HG23	2.20	0.41
1:A:300:HIS:CG	1:A:301:ARG:N	2.86	0.41
1:A:207:ARG:HD2	1:A:207:ARG:HH11	1.75	0.41
1:B:54:LYS:HE3	1:B:287:LEU:O	2.21	0.41
2:A:1301:ADN:HN62	1:B:33:GLN:NE2	2.01	0.40
1:B:70:LYS:HG2	1:B:90:TYR:HD2	1.87	0.40
1:A:6:GLU:HG2	1:A:258:VAL:HG13	2.04	0.40
1:A:28:PRO:HD3	5:A:2003:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/302 (98%)	283 (95%)	13 (4%)	1 (0%)	46	34
1	B	298/302 (99%)	290 (97%)	7 (2%)	1 (0%)	46	34
All	All	595/604 (98%)	573 (96%)	20 (3%)	2 (0%)	46	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	238	LYS
1	B	6	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/257 (99%)	235 (92%)	20 (8%)	16	6
1	B	256/257 (100%)	236 (92%)	20 (8%)	16	6
All	All	511/514 (99%)	471 (92%)	40 (8%)	16	6

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LYS
1	A	23	GLU
1	A	24	LYS
1	A	30	THR
1	A	38	ARG
1	A	70	LYS
1	A	72	SER
1	A	74	TYR
1	A	76	ARG
1	A	84	ASN
1	A	119	LYS
1	A	122	LYS
1	A	155	ASN
1	A	156	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	163	GLN
1	A	172	MET
1	A	192	ARG
1	A	205	LEU
1	A	225	LYS
1	A	297	LEU
1	B	4	LYS
1	B	34	ILE
1	B	70	LYS
1	B	72	SER
1	B	74	TYR
1	B	76	ARG
1	B	84	ASN
1	B	103	THR
1	B	109	GLN
1	B	123	GLU
1	B	142	GLU
1	B	176	ILE
1	B	186[A]	ASN
1	B	186[B]	ASN
1	B	198	ASN
1	B	225	LYS
1	B	227	LYS
1	B	292	LYS
1	B	298	GLU
1	B	299	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	84	ASN
1	A	109	GLN
1	A	128	ASN
1	A	144	ASN
1	A	163	GLN
1	A	179	HIS
1	A	181	ASN
1	A	286	ASN
1	A	300	HIS
1	B	21	ASN
1	B	33	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	84	ASN
1	B	144	ASN
1	B	179	HIS
1	B	181	ASN
1	B	198	ASN
1	B	286	ASN
1	B	300	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADN	A	1301	-	16,21,21	1.72	1 (6%)	16,31,31	3.13	5 (31%)
4	ANP	A	1304	-	27,33,33	3.95	9 (33%)	30,52,52	2.74	8 (26%)
2	ADN	B	1301	-	16,21,21	1.89	2 (12%)	16,31,31	2.69	6 (37%)
4	ANP	B	1304	-	27,33,33	3.79	7 (25%)	30,52,52	2.47	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADN	A	1301	-	-	0/2/22/22	0/3/3/3
4	ANP	A	1304	-	-	0/12/38/38	0/3/3/3
2	ADN	B	1301	-	-	0/2/22/22	0/3/3/3
4	ANP	B	1304	-	-	0/12/38/38	0/3/3/3

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1304	ANP	C2-N3	2.27	1.36	1.32
4	B	1304	ANP	PA-O1A	2.28	1.59	1.51
4	A	1304	ANP	C5-C4	2.38	1.45	1.40
4	A	1304	ANP	PA-O1A	2.42	1.60	1.51
2	B	1301	ADN	C5-C4	2.43	1.46	1.40
4	B	1304	ANP	C5-C4	2.76	1.46	1.40
4	A	1304	ANP	PB-O3A	4.17	1.64	1.59
4	A	1304	ANP	PB-N3B	4.35	1.74	1.63
4	B	1304	ANP	PG-N3B	4.39	1.75	1.63
4	B	1304	ANP	PB-N3B	4.60	1.75	1.63
4	A	1304	ANP	PG-N3B	4.76	1.75	1.63
2	A	1301	ADN	C4-N3	5.64	1.44	1.35
4	B	1304	ANP	C4-N3	6.07	1.44	1.35
2	B	1301	ADN	C4-N3	6.24	1.44	1.35
4	A	1304	ANP	C4-N3	7.19	1.46	1.35
4	B	1304	ANP	PG-O1G	11.64	1.59	1.46
4	A	1304	ANP	PB-O1B	11.71	1.59	1.46
4	A	1304	ANP	PG-O1G	11.93	1.59	1.46
4	B	1304	ANP	PB-O1B	11.97	1.59	1.46

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1301	ADN	N3-C2-N1	-10.12	121.14	128.89
4	A	1304	ANP	N3-C2-N1	-9.02	121.99	128.89
4	A	1304	ANP	C2'-C1'-N9	-8.08	101.95	114.29
2	B	1301	ADN	N3-C2-N1	-7.90	122.85	128.89
4	B	1304	ANP	N3-C2-N1	-7.70	123.00	128.89
4	B	1304	ANP	C2'-C1'-N9	-6.64	104.15	114.29
4	B	1304	ANP	PA-O3A-PB	-4.64	117.12	132.67
4	A	1304	ANP	PA-O3A-PB	-4.24	118.43	132.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1304	ANP	C4-C5-N7	-3.84	105.95	109.48
2	B	1301	ADN	C4-C5-N7	-3.83	105.95	109.48
2	A	1301	ADN	C4-C5-N7	-3.62	106.15	109.48
2	B	1301	ADN	C2'-C1'-N9	-2.81	110.00	114.29
4	A	1304	ANP	O3G-PG-O1G	-2.34	107.28	113.49
2	A	1301	ADN	C1'-N9-C4	-2.27	123.52	126.94
4	B	1304	ANP	O1G-PG-N3B	-2.12	108.65	111.90
4	B	1304	ANP	C4'-O4'-C1'	2.13	112.06	109.72
4	A	1304	ANP	O3'-C3'-C4'	2.25	117.81	111.05
2	A	1301	ADN	C2-N1-C6	2.32	122.91	118.77
4	A	1304	ANP	O2B-PB-O3A	2.32	115.62	105.09
2	B	1301	ADN	C2-N1-C6	2.39	123.03	118.77
4	A	1304	ANP	O3A-PB-N3B	2.63	113.68	106.44
4	B	1304	ANP	C2-N1-C6	2.84	123.85	118.77
2	B	1301	ADN	C4'-O4'-C1'	2.93	112.94	109.72
4	A	1304	ANP	C2-N1-C6	2.98	124.09	118.77
2	A	1301	ADN	C5'-C4'-C3'	3.11	122.45	115.08
2	B	1301	ADN	O4'-C1'-N9	3.11	114.62	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1301	ADN	3	0
4	A	1304	ANP	1	0
2	B	1301	ADN	1	0
4	B	1304	ANP	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/302 (99%)	0.83	32 (10%) 8 9	14, 19, 28, 42	24 (8%)
1	B	298/302 (98%)	0.92	36 (12%) 6 6	13, 19, 28, 43	32 (10%)
All	All	597/604 (98%)	0.88	68 (11%) 7 7	13, 19, 28, 43	56 (9%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	ASN	8.9
1	B	278	VAL	8.5
1	B	239	VAL	7.4
1	B	240	ILE	7.1
1	A	301	ARG	6.3
1	B	277	VAL	6.0
1	B	243	THR	5.3
1	B	274	ALA	5.2
1	A	5	MET	5.1
1	A	239	VAL	4.9
1	B	5	MET	4.8
1	B	238	LYS	4.6
1	B	242	PRO	4.5
1	B	294	VAL	4.5
1	B	245	ALA	4.4
1	B	258	VAL	4.3
1	A	237	GLY	4.2
1	B	3	GLY	4.0
1	A	240	ILE	3.9
1	B	4	LYS	3.9
1	B	282	GLY	3.7
1	A	70	LYS	3.6
1	B	236	ALA	3.5
1	A	3	GLY	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	234	ILE	3.4
1	A	76	ARG	3.3
1	B	300	HIS	3.3
1	B	234	ILE	3.3
1	A	300	HIS	3.2
1	A	245	ALA	3.1
1	B	297	LEU	3.1
1	B	279	GLU	3.1
1	B	287	LEU	3.0
1	A	4	LYS	3.0
1	B	288	PRO	2.9
1	B	71	ASN	2.9
1	B	296	ARG	2.9
1	A	217	SER	2.8
1	A	248	SER	2.8
1	B	237	GLY	2.8
1	A	107	ASN	2.7
1	B	298	GLU	2.7
1	B	244	GLY	2.7
1	A	268	LEU	2.7
1	A	244	GLY	2.6
1	B	241	ASP	2.6
1	A	277	VAL	2.6
1	A	30	THR	2.6
1	A	72	SER	2.5
1	A	279	GLU	2.5
1	B	280	ALA	2.4
1	A	298	GLU	2.3
1	A	280	ALA	2.3
1	B	72	SER	2.3
1	B	20	PHE	2.3
1	B	293	VAL	2.3
1	A	276	PHE	2.3
1	B	290	TRP	2.3
1	B	283	CYS	2.2
1	A	252	GLY	2.2
1	A	238	LYS	2.2
1	A	88	LEU	2.2
1	B	106	ASP	2.2
1	A	273	THR	2.2
1	A	269	ILE	2.1
1	B	113	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	6	GLU	2.0
1	A	221	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ADN	B	1301	19/19	0.55	0.53	12.32	43,46,49,51	11
4	ANP	A	1304	31/31	0.84	0.33	2.55	35,41,45,47	7
4	ANP	B	1304	31/31	0.76	0.37	1.53	54,59,68,69	7
3	MG	B	1302	1/1	0.76	0.16	0.45	54,54,54,54	0
2	ADN	A	1301	19/19	0.95	0.09	-0.95	20,24,27,28	0
3	MG	A	1302	1/1	0.94	0.08	-1.85	30,30,30,30	0
3	MG	B	1303	1/1	0.82	0.23	-	42,42,42,42	0
3	MG	A	1303	1/1	0.97	0.07	-	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.