



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:15 AM GMT

PDB ID : 2C8T
Title : THE 3.0 Å RESOLUTION STRUCTURE OF CASEINOLYTIC CLP PROTEASE 1 FROM MYCOBACTERIUM TUBERCULOSIS
Authors : Ingvarsson, H.; Högbohm, M.; Jones, T.A.; Unge, T.
Deposited on : 2005-12-07
Resolution : 3.00 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	 2% 63% 17% 17%
1	B	206	 2% 65% 15% 17%
1	C	206	 2% 65% 15% 17%
1	D	206	 2% 64% 15% 17%
1	E	206	 2% 62% 17% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	206	<p>2% 62% 18% • 17%</p>
1	G	206	<p>2% 63% 16% • 17%</p>
1	H	206	<p>2% 63% 17% • 17%</p>
1	I	206	<p>0% 64% 15% 5% 17%</p>
1	J	206	<p>0% 63% 17% • 17%</p>
1	K	206	<p>0% 66% 14% • 17%</p>
1	L	206	<p>0% 65% 16% • 17%</p>
1	M	206	<p>0% 63% 17% • 17%</p>
1	N	206	<p>2% 67% 13% • 17%</p>

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 18482 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUB-UNIT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	171	1322	839	224	250	9	0	0	0
1	B	171	1320	839	223	249	9	0	0	0
1	C	171	1320	839	223	249	9	0	0	0
1	D	171	1320	839	223	249	9	0	0	0
1	E	171	1320	839	223	249	9	0	0	0
1	F	171	1320	839	223	249	9	0	0	0
1	G	171	1320	839	223	249	9	0	0	0
1	H	171	1320	839	223	249	9	0	0	0
1	I	171	1320	839	223	249	9	0	0	0
1	J	171	1320	839	223	249	9	0	0	0
1	K	171	1320	839	223	249	9	0	0	0
1	L	171	1320	839	223	249	9	0	0	0
1	M	171	1320	839	223	249	9	0	0	0
1	N	171	1320	839	223	249	9	0	0	0

There are 98 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP P0A526
A	1	ALA	-	EXPRESSION TAG	UNP P0A526
A	201	GLY	-	EXPRESSION TAG	UNP P0A526
A	202	HIS	-	EXPRESSION TAG	UNP P0A526
A	203	HIS	-	EXPRESSION TAG	UNP P0A526
A	204	HIS	-	EXPRESSION TAG	UNP P0A526
A	205	HIS	-	EXPRESSION TAG	UNP P0A526
B	0	MET	-	EXPRESSION TAG	UNP P0A526
B	1	ALA	-	EXPRESSION TAG	UNP P0A526
B	201	GLY	-	EXPRESSION TAG	UNP P0A526
B	202	HIS	-	EXPRESSION TAG	UNP P0A526
B	203	HIS	-	EXPRESSION TAG	UNP P0A526
B	204	HIS	-	EXPRESSION TAG	UNP P0A526
B	205	HIS	-	EXPRESSION TAG	UNP P0A526
C	0	MET	-	EXPRESSION TAG	UNP P0A526
C	1	ALA	-	EXPRESSION TAG	UNP P0A526
C	201	GLY	-	EXPRESSION TAG	UNP P0A526
C	202	HIS	-	EXPRESSION TAG	UNP P0A526
C	203	HIS	-	EXPRESSION TAG	UNP P0A526
C	204	HIS	-	EXPRESSION TAG	UNP P0A526
C	205	HIS	-	EXPRESSION TAG	UNP P0A526
D	0	MET	-	EXPRESSION TAG	UNP P0A526
D	1	ALA	-	EXPRESSION TAG	UNP P0A526
D	201	GLY	-	EXPRESSION TAG	UNP P0A526
D	202	HIS	-	EXPRESSION TAG	UNP P0A526
D	203	HIS	-	EXPRESSION TAG	UNP P0A526
D	204	HIS	-	EXPRESSION TAG	UNP P0A526
D	205	HIS	-	EXPRESSION TAG	UNP P0A526
E	0	MET	-	EXPRESSION TAG	UNP P0A526
E	1	ALA	-	EXPRESSION TAG	UNP P0A526
E	201	GLY	-	EXPRESSION TAG	UNP P0A526
E	202	HIS	-	EXPRESSION TAG	UNP P0A526
E	203	HIS	-	EXPRESSION TAG	UNP P0A526
E	204	HIS	-	EXPRESSION TAG	UNP P0A526
E	205	HIS	-	EXPRESSION TAG	UNP P0A526
F	0	MET	-	EXPRESSION TAG	UNP P0A526
F	1	ALA	-	EXPRESSION TAG	UNP P0A526
F	201	GLY	-	EXPRESSION TAG	UNP P0A526
F	202	HIS	-	EXPRESSION TAG	UNP P0A526
F	203	HIS	-	EXPRESSION TAG	UNP P0A526
F	204	HIS	-	EXPRESSION TAG	UNP P0A526
F	205	HIS	-	EXPRESSION TAG	UNP P0A526
G	0	MET	-	EXPRESSION TAG	UNP P0A526

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	1	ALA	-	EXPRESSION TAG	UNP P0A526
G	201	GLY	-	EXPRESSION TAG	UNP P0A526
G	202	HIS	-	EXPRESSION TAG	UNP P0A526
G	203	HIS	-	EXPRESSION TAG	UNP P0A526
G	204	HIS	-	EXPRESSION TAG	UNP P0A526
G	205	HIS	-	EXPRESSION TAG	UNP P0A526
H	0	MET	-	EXPRESSION TAG	UNP P0A526
H	1	ALA	-	EXPRESSION TAG	UNP P0A526
H	201	GLY	-	EXPRESSION TAG	UNP P0A526
H	202	HIS	-	EXPRESSION TAG	UNP P0A526
H	203	HIS	-	EXPRESSION TAG	UNP P0A526
H	204	HIS	-	EXPRESSION TAG	UNP P0A526
H	205	HIS	-	EXPRESSION TAG	UNP P0A526
I	0	MET	-	EXPRESSION TAG	UNP P0A526
I	1	ALA	-	EXPRESSION TAG	UNP P0A526
I	201	GLY	-	EXPRESSION TAG	UNP P0A526
I	202	HIS	-	EXPRESSION TAG	UNP P0A526
I	203	HIS	-	EXPRESSION TAG	UNP P0A526
I	204	HIS	-	EXPRESSION TAG	UNP P0A526
I	205	HIS	-	EXPRESSION TAG	UNP P0A526
J	0	MET	-	EXPRESSION TAG	UNP P0A526
J	1	ALA	-	EXPRESSION TAG	UNP P0A526
J	201	GLY	-	EXPRESSION TAG	UNP P0A526
J	202	HIS	-	EXPRESSION TAG	UNP P0A526
J	203	HIS	-	EXPRESSION TAG	UNP P0A526
J	204	HIS	-	EXPRESSION TAG	UNP P0A526
J	205	HIS	-	EXPRESSION TAG	UNP P0A526
K	0	MET	-	EXPRESSION TAG	UNP P0A526
K	1	ALA	-	EXPRESSION TAG	UNP P0A526
K	201	GLY	-	EXPRESSION TAG	UNP P0A526
K	202	HIS	-	EXPRESSION TAG	UNP P0A526
K	203	HIS	-	EXPRESSION TAG	UNP P0A526
K	204	HIS	-	EXPRESSION TAG	UNP P0A526
K	205	HIS	-	EXPRESSION TAG	UNP P0A526
L	0	MET	-	EXPRESSION TAG	UNP P0A526
L	1	ALA	-	EXPRESSION TAG	UNP P0A526
L	201	GLY	-	EXPRESSION TAG	UNP P0A526
L	202	HIS	-	EXPRESSION TAG	UNP P0A526
L	203	HIS	-	EXPRESSION TAG	UNP P0A526
L	204	HIS	-	EXPRESSION TAG	UNP P0A526
L	205	HIS	-	EXPRESSION TAG	UNP P0A526
M	0	MET	-	EXPRESSION TAG	UNP P0A526

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	1	ALA	-	EXPRESSION TAG	UNP P0A526
M	201	GLY	-	EXPRESSION TAG	UNP P0A526
M	202	HIS	-	EXPRESSION TAG	UNP P0A526
M	203	HIS	-	EXPRESSION TAG	UNP P0A526
M	204	HIS	-	EXPRESSION TAG	UNP P0A526
M	205	HIS	-	EXPRESSION TAG	UNP P0A526
N	0	MET	-	EXPRESSION TAG	UNP P0A526
N	1	ALA	-	EXPRESSION TAG	UNP P0A526
N	201	GLY	-	EXPRESSION TAG	UNP P0A526
N	202	HIS	-	EXPRESSION TAG	UNP P0A526
N	203	HIS	-	EXPRESSION TAG	UNP P0A526
N	204	HIS	-	EXPRESSION TAG	UNP P0A526
N	205	HIS	-	EXPRESSION TAG	UNP P0A526


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1




- Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1



- Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1



- Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1





• Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1




• Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1




• Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1



• Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1



• Molecule 1: ATP-DEPENDENT CLP PROTEASE PROTEOLYTIC SUBUNIT 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.72Å 168.22Å 103.76Å 90.00° 114.60° 90.00°	Depositor
Resolution (Å)	19.87 – 3.00 19.87 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.87-3.00) 100.0 (19.87-3.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 2.98Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.213 , 0.238 0.215 , 0.237	Depositor DCC
R_{free} test set	3036 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	53.2	Xtrriage
Anisotropy	0.116	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.1	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	0 of 60123 reflections	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	18482	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/1344	0.66	1/1815 (0.1%)
1	B	0.60	0/1342	0.70	1/1812 (0.1%)
1	C	0.68	0/1342	0.75	1/1812 (0.1%)
1	D	0.72	0/1342	0.74	1/1812 (0.1%)
1	E	0.66	0/1342	0.73	1/1812 (0.1%)
1	F	0.76	0/1342	0.79	2/1812 (0.1%)
1	G	0.68	0/1342	0.70	1/1812 (0.1%)
1	H	0.70	0/1342	0.74	1/1812 (0.1%)
1	I	0.76	0/1342	0.81	2/1812 (0.1%)
1	J	0.62	0/1342	0.71	2/1812 (0.1%)
1	K	0.60	0/1342	0.72	1/1812 (0.1%)
1	L	0.72	0/1342	0.77	1/1812 (0.1%)
1	M	0.59	0/1342	0.68	1/1812 (0.1%)
1	N	0.58	0/1342	0.70	1/1812 (0.1%)
All	All	0.66	0/18790	0.73	17/25371 (0.1%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	121	LEU	CA-CB-CG	7.06	131.55	115.30
1	K	121	LEU	CA-CB-CG	6.73	130.78	115.30
1	N	121	LEU	CA-CB-CG	6.71	130.74	115.30
1	A	121	LEU	CA-CB-CG	6.57	130.41	115.30
1	B	121	LEU	CA-CB-CG	6.46	130.17	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1322	0	1316	41	0
1	B	1320	0	1315	37	0
1	C	1320	0	1315	33	0
1	D	1320	0	1315	37	0
1	E	1320	0	1315	39	0
1	F	1320	0	1315	40	0
1	G	1320	0	1315	44	0
1	H	1320	0	1315	31	0
1	I	1320	0	1315	34	0
1	J	1320	0	1315	41	0
1	K	1320	0	1315	34	0
1	L	1320	0	1315	36	0
1	M	1320	0	1315	37	0
1	N	1320	0	1315	34	0
All	All	18482	0	18411	450	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 450 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:121:LEU:HD11	1:N:142:GLN:HE22	1.18	1.09
1:D:142:GLN:HE22	1:E:121:LEU:HD11	1.13	1.07
1:L:90:THR:HG22	1:L:104:LEU:HD12	1.34	1.06
1:E:90:THR:HG22	1:E:104:LEU:HD12	1.36	1.06
1:M:90:THR:HG22	1:M:104:LEU:HD12	1.37	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	B	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	C	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	D	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	E	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	F	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	G	167/206 (81%)	160 (96%)	7 (4%)	0	100	100
1	H	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	I	167/206 (81%)	162 (97%)	5 (3%)	0	100	100
1	J	167/206 (81%)	160 (96%)	7 (4%)	0	100	100
1	K	167/206 (81%)	161 (96%)	6 (4%)	0	100	100
1	L	167/206 (81%)	160 (96%)	7 (4%)	0	100	100
1	M	167/206 (81%)	159 (95%)	8 (5%)	0	100	100
1	N	167/206 (81%)	160 (96%)	7 (4%)	0	100	100
All	All	2338/2884 (81%)	2249 (96%)	89 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/161 (84%)	126 (93%)	10 (7%)	17	52
1	B	135/161 (84%)	125 (93%)	10 (7%)	17	52
1	C	135/161 (84%)	125 (93%)	10 (7%)	17	52
1	D	135/161 (84%)	125 (93%)	10 (7%)	17	52
1	E	135/161 (84%)	127 (94%)	8 (6%)	24	63
1	F	135/161 (84%)	125 (93%)	10 (7%)	17	52
1	G	135/161 (84%)	125 (93%)	10 (7%)	17	52
1	H	135/161 (84%)	122 (90%)	13 (10%)	10	38
1	I	135/161 (84%)	123 (91%)	12 (9%)	12	42
1	J	135/161 (84%)	127 (94%)	8 (6%)	24	63
1	K	135/161 (84%)	126 (93%)	9 (7%)	20	57
1	L	135/161 (84%)	128 (95%)	7 (5%)	29	68
1	M	135/161 (84%)	125 (93%)	10 (7%)	17	52
1	N	135/161 (84%)	126 (93%)	9 (7%)	20	57
All	All	1891/2254 (84%)	1755 (93%)	136 (7%)	18	53

5 of 136 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	119	ARG
1	H	158	THR
1	M	190	ILE
1	G	154	ASN
1	H	30	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	154	ASN
1	I	154	ASN
1	N	47	GLN
1	H	154	ASN
1	C	154	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	171/206 (83%)	-0.22	3 (1%) 71 43	12, 14, 20, 23	0
1	B	171/206 (83%)	-0.28	3 (1%) 71 43	12, 14, 20, 23	0
1	C	171/206 (83%)	-0.38	4 (2%) 64 33	12, 14, 20, 23	0
1	D	171/206 (83%)	-0.41	4 (2%) 64 33	12, 14, 20, 23	0
1	E	171/206 (83%)	-0.36	2 (1%) 81 55	12, 14, 20, 23	0
1	F	171/206 (83%)	-0.41	4 (2%) 64 33	12, 14, 20, 23	0
1	G	171/206 (83%)	-0.29	5 (2%) 55 26	12, 14, 20, 23	0
1	H	171/206 (83%)	-0.41	4 (2%) 64 33	12, 14, 20, 23	0
1	I	171/206 (83%)	-0.51	1 (0%) 90 73	12, 14, 20, 23	0
1	J	171/206 (83%)	-0.38	2 (1%) 81 55	12, 14, 20, 23	0
1	K	171/206 (83%)	-0.35	3 (1%) 71 43	12, 14, 20, 23	0
1	L	171/206 (83%)	-0.42	2 (1%) 81 55	12, 14, 20, 23	0
1	M	171/206 (83%)	-0.33	3 (1%) 71 43	12, 14, 20, 23	0
1	N	171/206 (83%)	-0.23	5 (2%) 55 26	12, 14, 20, 23	0
All	All	2394/2884 (83%)	-0.36	45 (1%) 70 41	12, 14, 20, 23	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	194	HIS	6.1
1	G	138	ILE	5.5
1	D	194	HIS	5.3
1	N	138	ILE	4.7
1	H	194	HIS	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.