



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:48 AM GMT

PDB ID : 3C98
Title : Revised structure of the munc18a-syntaxin1 complex
Authors : Hattendorf, D.A.; Misura, K.M.; Burkhardt, P.; Scheller, R.H.; Fasshauer, D.; Weis, W.I.
Deposited on : 2008-02-15
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	606	<div> <div></div> <div>78%</div> <div>14%</div> <div>8%</div> </div>
2	B	279	<div> <div>5%</div> <div>73%</div> <div>9%</div> <div>18%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Syntaxin-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	556	Total	C	N	O	S	0	0	0
			4457	2827	756	849	25			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP P61765
A	-10	ARG	-	EXPRESSION TAG	UNP P61765
A	-9	GLY	-	EXPRESSION TAG	UNP P61765
A	-8	SER	-	EXPRESSION TAG	UNP P61765
A	-7	HIS	-	EXPRESSION TAG	UNP P61765
A	-6	HIS	-	EXPRESSION TAG	UNP P61765
A	-5	HIS	-	EXPRESSION TAG	UNP P61765
A	-4	HIS	-	EXPRESSION TAG	UNP P61765
A	-3	HIS	-	EXPRESSION TAG	UNP P61765
A	-2	HIS	-	EXPRESSION TAG	UNP P61765
A	-1	GLY	-	EXPRESSION TAG	UNP P61765
A	0	SER	-	EXPRESSION TAG	UNP P61765

- Molecule 2 is a protein called Syntaxin-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	230	Total	C	N	O	S	0	0	0
			1870	1145	331	383	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-11	MET	-	EXPRESSION TAG	UNP P32851
B	-10	ARG	-	EXPRESSION TAG	UNP P32851
B	-9	GLY	-	EXPRESSION TAG	UNP P32851
B	-8	SER	-	EXPRESSION TAG	UNP P32851

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	HIS	-	EXPRESSION TAG	UNP P32851
B	-6	HIS	-	EXPRESSION TAG	UNP P32851
B	-5	HIS	-	EXPRESSION TAG	UNP P32851
B	-4	HIS	-	EXPRESSION TAG	UNP P32851
B	-3	HIS	-	EXPRESSION TAG	UNP P32851
B	-2	HIS	-	EXPRESSION TAG	UNP P32851
B	-1	GLY	-	EXPRESSION TAG	UNP P32851
B	0	SER	-	EXPRESSION TAG	UNP P32851

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total	O	0	0
			58	58		
3	B	11	Total	O	0	0
			11	11		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

Label	Value
ALA	1
PRO	1
GLY	1
GLU	1
TYR	1
R552	1
S533	1
G534	1
I538	1
I539	1
L547	1
M550	1
R551	1
C552	1
G568	1
L578	1
T588	1
D589	1
I592	1
SER	1
SER	1
S158	1
K161	1
M164	1
K165	1
M166	1
P167	1
M416	1
L423	1
K424	1
K425	1
L446	1
T455	1
R458	1
K461	1
E466	1
K467	1
I468	1
S469	1
I482	1
K483	1
D484	1
I485	1
M486	1
T489	1
I502	1
S503	1
T504	1
R505	1
SER	1
SER	1
ALA	1
SER	1
PHE	1
SER	1
SER	1
SER	1
THR	1
THR	1
ALA	1
VAL	1
SER	1
ALA	1
ALA	1
TRG	1
TYR	1
GLY	1
HIS	1
TRP	1
HIS	1
LYS	1
ASN	1
LYS	1
V284	1
S158	1
K161	1
M164	1
K165	1
M166	1
P167	1
E221	1
L230	1
P242	1
F249	1
N261	1
D262	1
G270	1
L271	1
V279	1
L291	1
H295	1
V299	1
V303	1
T304	1
R305	1
D309	1
K314	1
R315	1
M316	1
ASN	1
THR	1
GLY	1
GLU	1
LYS	1
THR	1
THR	1
M324	1
R325	1
D326	1
L327	1
S328	1
Q329	1
M330	1
L331	1
K332	1
L350	1
V384	1
I4	1
K13	1
D17	1
R39	1
M51	1
T52	1
P69	1
E72	1
E81	1
H85	1
P94	1
P95	1
R100	1
A101	1
C110	1
P111	1
L114	1
F115	1
V119	1
L130	1
I135	1
Y145	1
S152	1
F153	1
Q154	1
S155	1
F156	1
Y157	1

Chain B:

Sequence logo for Chain B. The y-axis represents information content in bits (0.00 to 0.15). The x-axis shows amino acid positions (MET, ARG, GLY, SER, HIS, HIS, HIS, HIS, GLY, SER, MET, K2, Q6, E7, L8, A9, THR, ALA, LYS, ASP, SER, ASP, ASP, ASP, ASP, VAL, THR, VAL, VAL, VAL, ASP, ARG, D27, Q36, I40, S64, P67, L75, M79, T122, K126, V130, Y141, R148). A horizontal bar at the top indicates the percentage of positions with conservation: 5% (red), 73% (green), 9% (yellow), and 18% (grey).

4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	158.07Å 158.07Å 81.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.88 – 2.60 33.88 – 2.60	Depositor EDS
% Data completeness (in resolution range)	90.3 (33.88-2.60) 90.3 (33.88-2.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.03 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.203 , 0.263 0.201 , 0.263	Depositor DCC
R_{free} test set	2369 reflections (8.03%)	DCC
Wilson B-factor (Å ²)	59.1	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 41.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 31369 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6396	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/4535	0.41	0/6121
2	B	0.23	0/1887	0.38	0/2521
All	All	0.23	0/6422	0.40	0/8642

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4457	0	4549	55	0
2	B	1870	0	1849	17	0
3	A	58	0	0	1	0
3	B	11	0	0	0	0
All	All	6396	0	6398	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:GLU:HG2	1:A:101:ALA:HB3	1.61	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:482:ILE:HD11	1:A:539:ILE:HG21	1.64	0.79
1:A:13:LYS:HD3	1:A:135:ILE:HD11	1.68	0.75
1:A:242:PRO:HB2	1:A:578:LEU:HD22	1.75	0.68
1:A:384:LYS:HD3	1:A:384:LYS:H	1.63	0.64
1:A:485:ILE:O	1:A:489:THR:HG22	1.97	0.64
1:A:483:LYS:HA	1:A:486:MET:HE3	1.84	0.60
1:A:157:TYR:CZ	1:A:446:LEU:HD13	2.38	0.58
2:B:141:TYR:HD1	2:B:205:LEU:HD12	1.67	0.58
2:B:191:ALA:O	2:B:195:ILE:HG13	2.04	0.57
1:A:384:LYS:HD3	1:A:384:LYS:N	2.20	0.56
1:A:324:MET:HB3	1:A:327:LEU:HB3	1.88	0.56
1:A:325:ARG:C	1:A:327:LEU:H	2.10	0.55
1:A:111:PRO:HG2	1:A:114:LEU:HD12	1.89	0.54
1:A:483:LYS:HA	1:A:486:MET:CE	2.37	0.53
1:A:416:ASN:HD22	1:A:461:LYS:HG2	1.74	0.53
1:A:51:MET:HB2	2:B:122:THR:HG21	1.91	0.53
1:A:331:LEU:HD12	1:A:331:LEU:H	1.74	0.53
2:B:64:SER:O	2:B:67:PRO:HD3	2.10	0.52
1:A:486:MET:CE	1:A:552:CYS:HB3	2.39	0.52
1:A:152:SER:HB3	1:A:156:PHE:CE2	2.45	0.52
1:A:305:ARG:O	1:A:305:ARG:HD2	2.11	0.51
1:A:261:ASN:O	1:A:262:ASP:HB2	2.11	0.51
1:A:482:ILE:O	1:A:486:MET:HG3	2.11	0.49
2:B:205:LEU:HD23	2:B:206:GLU:N	2.27	0.49
1:A:81:GLU:HG3	1:A:85:HIS:CE1	2.47	0.49
1:A:154:GLN:O	1:A:158:SER:HB3	2.13	0.49
1:A:145:TYR:CE1	1:A:568:GLY:HA3	2.49	0.48
1:A:69:PRO:O	1:A:100:ARG:HB2	2.13	0.48
2:B:165:LEU:HD23	2:B:165:LEU:O	2.13	0.48
1:A:52:THR:HG22	1:A:52:THR:O	2.12	0.48
1:A:115:PHE:HE1	1:A:130:LEU:HD21	1.79	0.47
2:B:230:ILE:O	2:B:230:ILE:HG12	2.14	0.47
1:A:13:LYS:O	1:A:17:ASP:HB2	2.15	0.47
1:A:329:GLN:CG	1:A:332:LYS:HE2	2.44	0.47
2:B:173:ASN:HB2	2:B:176:ILE:HB	1.97	0.46
2:B:148:ARG:O	2:B:152:GLN:HG2	2.16	0.46
1:A:468:ILE:O	1:A:469:SER:C	2.54	0.46
1:A:52:THR:O	1:A:52:THR:CG2	2.63	0.46
1:A:486:MET:HE1	1:A:552:CYS:HB3	1.99	0.45
1:A:329:GLN:HG3	1:A:332:LYS:HE2	1.97	0.45
1:A:408:ILE:O	1:A:412:ILE:HG13	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:LEU:HD11	1:A:303:VAL:HG12	1.99	0.45
1:A:455:THR:HA	1:A:458:ARG:HD2	1.99	0.45
1:A:324:MET:O	1:A:325:ARG:HG3	2.17	0.44
1:A:547:LEU:HA	1:A:550:MET:HG3	1.99	0.44
2:B:75:LEU:O	2:B:79:MET:HG3	2.17	0.44
1:A:166:ASN:N	1:A:167:PRO:HD2	2.32	0.44
2:B:126:LYS:O	2:B:130:VAL:HG23	2.18	0.43
1:A:161:LYS:HA	1:A:164:MET:CE	2.49	0.43
1:A:94:PRO:HB2	1:A:95:PRO:HD3	2.01	0.43
2:B:126:LYS:HE2	2:B:126:LYS:HA	2.01	0.43
1:A:271:ILE:HG22	1:A:271:ILE:O	2.18	0.43
1:A:115:PHE:CE1	1:A:130:LEU:HD21	2.53	0.43
1:A:295:HIS:O	1:A:299:VAL:HG23	2.18	0.43
1:A:486:MET:HE2	1:A:552:CYS:HB3	2.01	0.43
1:A:249:PHE:CZ	1:A:350:LEU:HD23	2.53	0.43
1:A:39:ARG:HD2	3:A:612:HOH:O	2.19	0.43
1:A:119:VAL:HG12	2:B:8:LEU:HD23	2.00	0.42
2:B:151:ARG:O	2:B:155:ILE:HG12	2.19	0.42
1:A:391:VAL:HB	1:A:392:PRO:HD3	2.01	0.42
2:B:241:VAL:HG22	2:B:242:ASP:N	2.35	0.41
2:B:36:GLN:O	2:B:40:ILE:HG13	2.20	0.41
1:A:230:LEU:HD12	1:A:538:ILE:HB	2.01	0.41
1:A:110:CYS:HA	1:A:111:PRO:HD2	1.83	0.41
1:A:588:THR:O	1:A:589:ASP:HB3	2.20	0.41
1:A:326:ASP:C	1:A:328:SER:H	2.22	0.41
1:A:486:MET:HE3	1:A:486:MET:HB2	1.94	0.40
2:B:2:LYS:HB2	2:B:2:LYS:HE3	1.84	0.40
1:A:314:LYS:O	1:A:315:ARG:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/606 (91%)	518 (94%)	27 (5%)	5 (1%)	21	42
2	B	226/279 (81%)	217 (96%)	8 (4%)	1 (0%)	39	65
All	All	776/885 (88%)	735 (95%)	35 (4%)	6 (1%)	24	46

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	534	GLY
1	A	469	SER
1	A	533	SER
1	A	466	GLU
1	A	502	ILE
2	B	174	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	503/543 (93%)	497 (99%)	6 (1%)	78	92
2	B	211/255 (83%)	210 (100%)	1 (0%)	92	98
All	All	714/798 (90%)	707 (99%)	7 (1%)	82	94

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	279	VAL
1	A	309	ASP
1	A	384	LYS
1	A	423	LEU
1	A	425	LYS
1	A	446	LEU
2	B	2	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	ASN
1	A	143	GLN
1	A	329	GLN
1	A	416	ASN
1	A	558	GLN
1	A	576	GLN
2	B	107	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	556/606 (91%)	-0.26	7 (1%) 79 75	40, 64, 109, 166	0
2	B	230/279 (82%)	-0.04	13 (5%) 27 20	50, 77, 139, 179	0
All	All	786/885 (88%)	-0.20	20 (2%) 61 54	40, 68, 122, 179	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	467	ARG	5.2
1	A	504	THR	5.2
1	A	466	GLU	4.8
2	B	183	MET	4.3
2	B	182	ILE	3.9
2	B	177	PHE	3.5
2	B	170	GLU	3.1
2	B	173	ASN	3.1
2	B	184	ASP	3.0
2	B	6	GLN	3.0
2	B	181	ILE	2.7
1	A	221	GLU	2.5
2	B	176	ILE	2.4
1	A	505	ARG	2.3
1	A	532	ARG	2.1
2	B	180	GLY	2.0
2	B	248	VAL	2.0
2	B	160	THR	2.0
1	A	270	GLY	2.0
2	B	247	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.