Full wwPDB X-ray Structure Validation Report (i)

Feb 1, 2016 – 03:22 PM GMT

PDB ID	:	4C9U
Title	:	Xenopus ZNRF3 ectodomain in complex with Xenopus RSPO2 Fu1-Fu2 crystal form II
Authors	:	Zebisch, M.; Jones, E.Y.
Deposited on	:	2013-10-03
Resolution	:	3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	81% 7% • 12%
1	C	182	77% 8% • 15%
2	B	121	3% 73% 12% • 15%
2	D	121	5% 62% 6% • 29%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 UBIQUITIN-PROTEIN LIGASE ZNRF3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	161	1231	771	216	237	7	0	0	0
			Total	C	N	O	S			
1	C	155	1188	747	210	224	7	0	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLU	-	EXPRESSION TAG	UNP Q08D68
A	23	THR	-	EXPRESSION TAG	UNP Q08D68
A	192	GLY	-	EXPRESSION TAG	UNP Q08D68
A	193	THR	-	EXPRESSION TAG	UNP Q08D68
A	194	HIS	-	EXPRESSION TAG	UNP Q08D68
A	195	HIS	-	EXPRESSION TAG	UNP Q08D68
A	196	HIS	-	EXPRESSION TAG	UNP Q08D68
A	197	HIS	-	EXPRESSION TAG	UNP Q08D68
A	198	HIS	-	EXPRESSION TAG	UNP Q08D68
A	199	HIS	-	EXPRESSION TAG	UNP Q08D68
A	200	HIS	-	EXPRESSION TAG	UNP Q08D68
A	201	HIS	-	EXPRESSION TAG	UNP Q08D68
A	202	HIS	-	EXPRESSION TAG	UNP Q08D68
A	203	HIS	-	EXPRESSION TAG	UNP Q08D68
C	22	GLU	-	EXPRESSION TAG	UNP Q08D68
C	23	THR	-	EXPRESSION TAG	UNP Q08D68
C	192	GLY	-	EXPRESSION TAG	UNP Q08D68
C	193	THR	-	EXPRESSION TAG	UNP Q08D68
C	194	HIS	-	EXPRESSION TAG	UNP Q08D68
C	195	HIS	-	EXPRESSION TAG	UNP Q08D68
C	196	HIS	-	EXPRESSION TAG	UNP Q08D68
C	197	HIS	-	EXPRESSION TAG	UNP Q08D68
C	198	HIS	-	EXPRESSION TAG	UNP Q08D68
C	199	HIS	-	EXPRESSION TAG	UNP Q08D68
C	200	HIS	-	EXPRESSION TAG	UNP Q08D68

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	201	HIS	-	EXPRESSION TAG	UNP Q08D68
C	202	HIS	-	EXPRESSION TAG	UNP Q08D68
C	203	HIS	-	EXPRESSION TAG	UNP Q08D68

- Molecule 2 is a protein called R-SPONDIN-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	103	810	498	145	148	19	0	0	0
			Total	C	N	O	S			
2	D	86	627	379	119	113	16	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	32	GLU	-	EXPRESSION TAG	UNP Q5M7L6
B	33	THR	-	EXPRESSION TAG	UNP Q5M7L6
B	34	GLY	-	EXPRESSION TAG	UNP Q5M7L6
B	145	THR	-	EXPRESSION TAG	UNP Q5M7L6
B	146	LYS	-	EXPRESSION TAG	UNP Q5M7L6
B	147	HIS	-	EXPRESSION TAG	UNP Q5M7L6
B	148	HIS	-	EXPRESSION TAG	UNP Q5M7L6
B	149	HIS	-	EXPRESSION TAG	UNP Q5M7L6
B	150	HIS	-	EXPRESSION TAG	UNP Q5M7L6
B	151	HIS	-	EXPRESSION TAG	UNP Q5M7L6
B	152	HIS	-	EXPRESSION TAG	UNP Q5M7L6
D	32	GLU	-	EXPRESSION TAG	UNP Q5M7L6
D	33	THR	-	EXPRESSION TAG	UNP Q5M7L6
D	34	GLY	-	EXPRESSION TAG	UNP Q5M7L6
D	145	THR	-	EXPRESSION TAG	UNP Q5M7L6
D	146	LYS	-	EXPRESSION TAG	UNP Q5M7L6
D	147	HIS	-	EXPRESSION TAG	UNP Q5M7L6
D	148	HIS	-	EXPRESSION TAG	UNP Q5M7L6
D	149	HIS	-	EXPRESSION TAG	UNP Q5M7L6
D	150	HIS	-	EXPRESSION TAG	UNP Q5M7L6
D	151	HIS	-	EXPRESSION TAG	UNP Q5M7L6
D	152	HIS	-	EXPRESSION TAG	UNP Q5M7L6

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: E3 UBIQUITIN-PROTEIN LIGASE ZNRF3

Chain A:



- Molecule 1: E3 UBIQUITIN-PROTEIN LIGASE ZNRF3

Chain C:

- Molecule 2: R-SPONDIN-2

Chain B:

- Molecule 2: R-SPONDIN-2

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α, β, γ	57.55Å 57.62Å 208.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.50 – 3.00 44.39 – 3.00	Depositor EDS
% Data completeness (in resolution range)	92.6 (50.50-3.00) 92.7 (44.39-3.00)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.72 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.227 , 0.310 0.229 , 0.305	Depositor DCC
R_{free} test set	677 reflections (5.26%)	DCC
Wilson B-factor (Å^2)	66.8	Xtriage
Anisotropy	0.147	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.27 , 35.8	EDS
Estimated twinning fraction	0.054 for k,h,-l	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	2 of 13539 reflections (0.015%)	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	3856	wwPDB-VP
Average B, all atoms (Å^2)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.66	0/1252	0.79	0/1694
1	C	0.63	0/1209	0.79	0/1635
2	B	0.66	0/828	0.83	0/1105
2	D	0.60	0/637	0.86	0/852
All	All	0.64	0/3926	0.81	0/5286

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1231	0	1218	8	0
1	C	1188	0	1181	9	0
2	B	810	0	751	5	0
2	D	627	0	558	7	0
All	All	3856	0	3708	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:GLN:HG2	2:D:48:LYS:HE3	1.77	0.66
1:A:98:LYS:NZ	1:A:133:ASP:OD2	2.28	0.66
2:B:133:ALA:HB1	2:B:134:PRO:HD2	1.75	0.66
1:A:101:GLN:HG2	2:D:48:LYS:CE	2.28	0.63
1:C:64:THR:HG22	1:C:64:THR:O	1.99	0.61
1:C:42:SER:HB2	1:C:44:ASN:HB3	1.83	0.60
2:D:40:CYS:SG	2:D:40:CYS:O	2.66	0.54
2:B:133:ALA:HB1	2:B:134:PRO:CD	2.39	0.52
2:B:138:THR:HG22	2:B:140:VAL:HG13	1.91	0.52
2:B:39:ILE:CG1	2:B:40:CYS:H	2.25	0.49
2:D:117:PHE:HB3	2:D:124:CYS:O	2.11	0.49
1:A:35:GLU:HG3	1:A:51:THR:CG2	2.41	0.49
2:D:120:HIS:N	2:D:123:GLN:O	2.35	0.49
1:A:35:GLU:HG3	1:A:51:THR:HG23	1.95	0.48
1:C:35:GLU:HG3	1:C:51:THR:HG23	1.96	0.47
1:C:130:ASP:C	1:C:130:ASP:OD1	2.53	0.47
1:C:35:GLU:HG3	1:C:51:THR:CG2	2.45	0.47
2:D:95:ARG:HG2	2:D:96:CYS:N	2.33	0.43
1:C:100:GLU:O	1:C:101:GLN:C	2.56	0.43
1:A:82:ASN:HB2	1:A:84:ASP:H	1.82	0.43
2:B:138:THR:O	2:B:139:MET:HB2	2.18	0.43
2:D:83:TYR:CZ	2:D:94:SER:HB3	2.53	0.43
1:A:82:ASN:O	1:A:83:ASN:HB2	2.19	0.43
1:C:103:GLU:H	1:C:103:GLU:CD	2.23	0.42
1:A:38:LEU:HD22	1:A:171:GLN:HG3	2.01	0.42
1:C:169:ASN:HB3	1:C:170:LYS:HG3	2.02	0.40
1:C:130:ASP:OD1	1:C:132:SER:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/182 (87%)	149 (94%)	10 (6%)	0	100	100
1	C	153/182 (84%)	144 (94%)	9 (6%)	0	100	100
2	B	101/121 (84%)	94 (93%)	7 (7%)	0	100	100
2	D	84/121 (69%)	77 (92%)	7 (8%)	0	100	100
All	All	497/606 (82%)	464 (93%)	33 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/155 (84%)	127 (97%)	4 (3%)	47	83
1	C	125/155 (81%)	121 (97%)	4 (3%)	46	82
2	B	92/107 (86%)	83 (90%)	9 (10%)	10	36
2	D	65/107 (61%)	60 (92%)	5 (8%)	16	50
All	All	413/524 (79%)	391 (95%)	22 (5%)	28	67

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	41	SER
1	A	48	LYS
1	A	82	ASN
2	B	39	ILE
2	B	65	ARG
2	B	86	ARG
2	B	89	ASP
2	B	94	SER
2	B	103	SER
2	B	107	ARG
2	B	128	CYS
2	B	132	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	48	LYS
1	C	147	ASP
1	C	163	LYS
1	C	169	ASN
2	D	40	CYS
2	D	48	LYS
2	D	94	SER
2	D	96	CYS
2	D	104	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	ASN
2	B	100	ASN
1	C	166	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	161/182 (88%)	-0.53	0 100 100	32, 56, 94, 119	0
1	C	155/182 (85%)	-0.46	0 100 100	36, 62, 104, 134	0
2	B	103/121 (85%)	-0.19	4 (3%) 43 18	38, 65, 111, 150	0
2	D	86/121 (71%)	0.07	6 (6%) 19 7	44, 80, 149, 157	0
All	All	505/606 (83%)	-0.34	10 (1%) 68 39	32, 64, 119, 157	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	120	HIS	4.5
2	B	138	THR	4.4
2	B	137	ASP	3.6
2	D	102	ASP	2.8
2	B	89	ASP	2.3
2	D	121	LYS	2.3
2	B	136	ASP	2.2
2	D	113	CYS	2.1
2	D	122	GLY	2.1
2	D	111	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.