



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 10:14 PM GMT

PDB ID : 5CA1
Title : Crystal structure of T2R-TTL-Nocodazole complex
Authors : Wang, Y.; Yu, Y.; Chen, Q.; Yang, J.
Deposited on : 2015-06-29
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	<div> <div>90%</div> <div>7%</div> </div>
1	C	450	<div> <div>91%</div> <div>7%</div> </div>
2	B	445	<div> <div>4%</div> <div>87%</div> <div>9%</div> </div>
2	D	445	<div> <div>6%</div> <div>81%</div> <div>13%</div> <div>5%</div> </div>
3	E	143	<div> <div>7%</div> <div>74%</div> <div>10%</div> <div>15%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	384	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	GOL	A	504	-	-	-	X

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 17872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	0	0
			3416	2163	581	650	22			
1	C	440	Total	C	N	O	S	0	0	0
			3437	2175	584	656	22			

- Molecule 2 is a protein called Tubulin beta-2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	427	Total	C	N	O	S	0	0	0
			3361	2110	576	649	26			
2	D	421	Total	C	N	O	S	0	0	0
			3309	2080	562	640	27			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	121	Total	C	N	O	S	0	0	0
			1000	617	181	197	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	-	expression tag	UNP P63043
E	4	ALA	-	expression tag	UNP P63043


- Molecule 4 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	334	Total	C	N	O	S	0	0	0
			2744	1761	470	499	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	A	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		
7	C	1	Total	Ca	0	0
			1	1		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
9	D	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 10 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 11 is nocodazole (three-letter code: NZO) (formula: $C_{14}H_{11}N_3O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	B	1	Total	C	N	O	S	0	0
			21	14	3	3	1		
11	D	1	Total	C	N	O	S	0	0
			21	14	3	3	1		

- Molecule 12 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	F	1	Total	C	N	O	P	0	0
			31	11	5	12	3		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	101	Total	O	0	0
			101	101		
13	B	74	Total	O	0	0
			74	74		
13	C	131	Total	O	0	0
			131	131		
13	D	33	Total	O	0	0
			33	33		
13	E	13	Total	O	0	0
			13	13		
13	F	25	Total	O	0	0
			25	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha




- Molecule 1: Tubulin alpha




- Molecule 2: Tubulin beta-2 chain




- Molecule 2: Tubulin beta-2 chain






• Molecule 3: Stathmin-4



• Molecule 4: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.44Å 158.36Å 180.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.52 – 2.40 41.52 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.52-2.40) 99.8 (41.52-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.192 , 0.225 0.205 , 0.214	Depositor DCC
R_{free} test set	5924 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	41.3	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 118147 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	17872	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, GOL, MG, CA, GTP, ACP, MES, NZO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3494	0.41	0/4743
1	C	0.25	0/3515	0.42	0/4772
2	B	0.26	0/3436	0.44	0/4654
2	D	0.24	0/3382	0.42	0/4581
3	E	0.24	0/1008	0.37	0/1337
4	F	0.32	0/2806	0.56	0/3791
All	All	0.26	0/17641	0.44	0/23878

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	F	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	F	138	ARG	Peptide
4	F	142	ARG	Peptide
4	F	242	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3416	0	3330	20	0
1	C	3437	0	3348	19	0
2	B	3361	0	3238	26	0
2	D	3309	0	3189	38	0
3	E	1000	0	1018	10	0
4	F	2744	0	2709	57	0
5	A	32	0	12	1	0
5	C	32	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0
8	A	6	0	8	0	0
9	B	28	0	12	0	0
9	D	28	0	12	2	0
10	B	12	0	12	2	0
10	D	12	0	12	3	0
11	B	21	0	11	0	0
11	D	21	0	11	2	0
12	F	31	0	14	5	0
13	A	101	0	0	0	0
13	B	74	0	0	1	0
13	C	131	0	0	0	0
13	D	33	0	0	0	0
13	E	13	0	0	0	0
13	F	25	0	0	3	0
All	All	17872	0	16948	164	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 164 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:410:GLU:O	2:B:414:ASN:ND2	2.00	0.93
4:F:180:HIS:O	13:F:501:HOH:O	1.92	0.87
4:F:138:ARG:HE	4:F:143:GLU:HB2	1.46	0.78
4:F:136:ASN:O	4:F:140:GLU:N	2.13	0.78
2:B:35:SER:OG	2:B:58:LYS:NZ	2.16	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	435/450 (97%)	425 (98%)	10 (2%)	0	100	100
1	C	438/450 (97%)	427 (98%)	11 (2%)	0	100	100
2	B	425/445 (96%)	413 (97%)	11 (3%)	1 (0%)	52	69
2	D	417/445 (94%)	399 (96%)	18 (4%)	0	100	100
3	E	117/143 (82%)	114 (97%)	3 (3%)	0	100	100
4	F	324/384 (84%)	308 (95%)	13 (4%)	3 (1%)	21	30
All	All	2156/2317 (93%)	2086 (97%)	66 (3%)	4 (0%)	52	69

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	57	ASN
4	F	236	LYS
4	F	142	ARG
4	F	140	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/378 (97%)	367 (100%)	1 (0%)	94	98
1	C	371/378 (98%)	370 (100%)	1 (0%)	94	98
2	B	369/383 (96%)	366 (99%)	3 (1%)	86	94
2	D	364/383 (95%)	360 (99%)	4 (1%)	80	92
3	E	109/127 (86%)	109 (100%)	0	100	100
4	F	301/342 (88%)	292 (97%)	9 (3%)	48	70
All	All	1882/1991 (94%)	1864 (99%)	18 (1%)	82	93

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	205	GLU
4	F	12	SER
4	F	234	GLN
2	D	95	SER
2	D	180	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	137	HIS
2	B	414	ASN
4	F	243	HIS
4	F	260	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 5 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	GTP	A	501	6	26,34,34	0.96	1 (3%)	29,54,54	1.60	4 (13%)
8	GOL	A	504	-	5,5,5	0.37	0	5,5,5	0.22	0
9	GDP	B	501	6	24,30,30	1.11	2 (8%)	26,47,47	1.89	5 (19%)
10	MES	B	503	-	12,12,12	2.05	1 (8%)	15,16,16	2.48	7 (46%)
11	NZO	B	504	-	23,23,23	1.66	4 (17%)	18,32,32	2.85	4 (22%)
5	GTP	C	501	6	26,34,34	0.92	1 (3%)	29,54,54	1.55	3 (10%)
9	GDP	D	501	-	24,30,30	1.16	2 (8%)	26,47,47	1.85	5 (19%)
10	MES	D	502	-	12,12,12	2.17	1 (8%)	15,16,16	2.36	3 (20%)
11	NZO	D	503	-	23,23,23	1.71	4 (17%)	18,32,32	2.85	4 (22%)
12	ACP	F	401	-	29,33,33	1.57	5 (17%)	29,52,52	1.71	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	501	6	-	0/18/38/38	0/3/3/3
8	GOL	A	504	-	-	0/4/4/4	0/0/0/0
9	GDP	B	501	6	-	0/12/32/32	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	MES	B	503	-	-	0/6/14/14	0/1/1/1
11	NZO	B	504	-	-	0/8/14/14	0/3/3/3
5	GTP	C	501	6	-	0/18/38/38	0/3/3/3
9	GDP	D	501	-	-	0/12/32/32	0/3/3/3
10	MES	D	502	-	-	0/6/14/14	0/1/1/1
11	NZO	D	503	-	-	0/8/14/14	0/3/3/3
12	ACP	F	401	-	-	0/15/38/38	0/3/3/3

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	D	502	MES	C8-S	-7.26	1.66	1.77
10	B	503	MES	C8-S	-6.79	1.67	1.77
12	F	401	ACP	PG-O3G	-3.94	1.45	1.54
11	D	503	NZO	C4-S	-3.59	1.67	1.72
11	B	504	NZO	C4-S	-3.30	1.68	1.72

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	D	503	NZO	C2-C1-S	-7.45	104.98	113.23
11	B	504	NZO	C2-C1-S	-7.15	105.32	113.23
12	F	401	ACP	N3-C2-N1	-6.54	123.73	128.87
5	A	501	GTP	N3-C2-N1	-5.52	120.05	127.56
5	C	501	GTP	N3-C2-N1	-5.29	120.36	127.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	501	GTP	1	0
10	B	503	MES	2	0
9	D	501	GDP	2	0
10	D	502	MES	3	0
11	D	503	NZO	2	0
12	F	401	ACP	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	437/450 (97%)	-0.16	3 (0%) 89 88	24, 39, 62, 87	0
1	C	440/450 (97%)	-0.38	1 (0%) 95 95	19, 31, 53, 69	0
2	B	427/445 (95%)	0.03	16 (3%) 45 46	22, 38, 69, 103	0
2	D	421/445 (94%)	0.21	26 (6%) 24 25	26, 48, 78, 101	0
3	E	121/143 (84%)	0.34	10 (8%) 14 14	29, 53, 83, 94	0
4	F	334/384 (86%)	0.53	54 (16%) 3 2	32, 59, 116, 135	0
All	All	2180/2317 (94%)	0.04	110 (5%) 32 33	19, 42, 82, 135	0

The worst 5 of 110 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	142	ARG	7.5
4	F	173	ILE	7.4
4	F	176	GLN	6.4
4	F	132	LEU	6.1
4	F	135	TYR	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
8	GOL	A	504	6/6	0.91	0.21	2.68	45,54,59,59	0
11	NZO	B	504	21/21	0.97	0.21	0.31	14,29,35,50	0
5	GTP	A	501	32/32	0.98	0.17	0.01	21,29,33,36	0
5	GTP	C	501	32/32	0.99	0.15	-0.09	19,24,30,30	0
10	MES	B	503	12/12	0.95	0.16	-0.15	32,43,51,53	0
10	MES	D	502	12/12	0.92	0.18	-0.20	50,58,63,66	0
11	NZO	D	503	21/21	0.94	0.15	-0.49	25,40,50,69	0
9	GDP	B	501	28/28	0.99	0.13	-0.58	23,29,31,40	0
9	GDP	D	501	28/28	0.94	0.13	-0.67	37,43,57,63	0
6	MG	A	502	1/1	0.98	0.13	-0.80	28,28,28,28	0
12	ACP	F	401	31/31	0.92	0.12	-1.20	73,81,94,97	0
7	CA	C	503	1/1	0.93	0.07	-1.67	45,45,45,45	0
7	CA	A	503	1/1	0.96	0.05	-2.93	54,54,54,54	0
6	MG	C	502	1/1	1.00	0.17	-	24,24,24,24	0
6	MG	B	502	1/1	0.95	0.19	-	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.